

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 48/00	A2	(11) International Publication Number: WO 99/26663 (43) International Publication Date: 3 June 1999 (03.06.99)
(21) International Application Number: PCT/US98/24830 (22) International Filing Date: 20 November 1998 (20.11.98) (30) Priority Data: 60/067,087 20 November 1997 (20.11.97) US 60/079,914 30 March 1998 (30.03.98) US 60/100,820 15 September 1998 (15.09.98) US (71) Applicant (for all designated States except US): VICAL, INC. [US/US]; Suite 100, 9373 Towne Centre Drive, San Diego, CA 92121-3088 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): HORTON, Holly [US/US]; 11791B Spruce Run Drive, San Diego, CA 92131 (US). PARKER, Suzanne [US/US]; 12539 Cypress Woods Court, San Diego, CA 92131 (US). MANTHORPE, Marston [US/US]; 418 Kestrel Street, San Diego, CA 92129 (US). FELGNER, Philip [US/US]; P.O. Box 3392, Rancho Santa Fe, CA 92067 (US). (74) Agents: STEFFE, Eric, K. et al.; Sterne, Kessler, Goldstein & Fox P.L.L.C., Suite 600, 1100 New York Avenue, N.W., Washington, DC 20005-3934 (US).		(81) Designated States: CA, JP, US, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: TREATMENT OF CANCER USING CYTOKINE-EXPRESSING POLYNUCLEOTIDES AND COMPOSITIONS THERE- FOR		
(57) Abstract <p>The present invention provides a pharmaceutical composition, comprising a non-infectious, non-integrating polynucleotide construct comprising a polynucleotide encoding an interferon ω and one or more cationic compounds. The present invention also provides methods of treating cancer in a mammal, comprising administering into a tissue of the mammal a non-infectious, non-integrating polynucleotide construct comprising a polynucleotide encoding a cytokine. In addition, the present invention also relates to the methodology for selective transfection of malignant cells with polynucleotides expressing therapeutic or prophylactic molecules in intracavity tumor bearing mammals. More specifically, the present invention provides a methodology for the suppression of an intra-cavity dissemination of malignant cells, such as intraperitoneal dissemination.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

Treatment of Cancer Using Cytokine-Expressing Polynucleotides and Compositions Therefor

Background of the Invention

5

Field of the Invention

The present invention relates to treatment of cancer in mammals. Generally, the present invention provides methods of treating cancer in a mammal by administering a polynucleotide construct comprising a polynucleotide encoding a cytokine. In addition, the present invention relates to the methodology for selective transfection of malignant cells with polynucleotides expressing therapeutic or prophylactic molecules in intra-cavity tumor bearing mammals. More specifically, the present invention provides a methodology for the suppression of an intra-cavity dissemination of malignant cells, such as intraperitoneal dissemination.

10
15

Related Art

Cytokines have been demonstrated both in pre-clinical animal models as well as in humans to have potent anti-tumor effects. In particular IFN's have been tried for the treatment of a number of human concerns.

The interferons (IFNs) are a family of cytokines with potent anti-viral, antiproliferative, and immunomodulatory activities and play important roles in the body's defensive response to viruses, bacteria, and tumors (Baron, S. *et al.*, *JAMA* 266:1375 (1991)). On the basis of antigenicity, biochemical properties, and producer cell, the interferon's have been divided into two classes, type I interferon and type II interferon. IFN α , IFN β , IFN ω , and IFN τ are type I interferons, and bind to the same α/β receptor. IFN γ is a type II interferon, and binds to the γ receptor (Pestka, S., *Ann. Rev. Biochem.* 56:727 (1987)). IFN α and IFN β are naturally expressed in many cells upon viral infection. IFN γ is produced by activated T lymphocytes and natural killer (NK) cells.

20
25
30

IFN τ is believed to possess hormone activity, and plays an important role in pregnancy in cattle, sheep, and related ruminants (Imakawa, K. *et al.*, *Nature* 330:377 (1987); Stewart, H.J. *et al.*, *J. Endocrinology* 115:R13 (1987)). Due to the pleiotropic activities of IFNs, these cytokines have been studied for their therapeutic efficacy in a number of diseases, particularly cancers and viral infectious diseases.

IFN ω was discovered independently by three different groups in 1985 (Capon, D.J., *et al.*, *Molec. Cell. Biol.* 5: 768-779 (1985), Feinstein, S. *et al.*, *Molec. Cell. Biol.* 5:510 (1985); and Hauptmann and Swetly, *Nucl. Acids Res.* 13: 4739-4749 (1985)). Unlike IFN α , for which at least 14 different functional nonallelic genes have been identified in man, IFN ω is encoded by a single functional gene. IFN ω genes are believed to be present in most mammals, but have not been found in dogs, rats or mice. The mature IFN ω polypeptide is 172 amino acids and shares 60% sequence homology with the human IFN α 's. Due to the sequence similarity with IFN α , IFN ω was originally considered to be a member or a subfamily of IFN α , and was originally termed IFN α -II. IFN ω is a significant component ($\approx 10\%$) of human leukocyte-derived interferon, the natural mixture of interferon produced after viral infection (Adolf, G. *et al.*, *Virology* 175:410 (1990)). IFN ω has been demonstrated to bind to the same α/β receptor as IFN α (Flores, I. *et al.*, *J. Biol. Chem.* 266: 19875-19877 (1991)), and to share similar biological activities with IFN α , including anti-proliferative activity against tumor cells *in vitro* (Kubes, M. *et al.*, *J. Interferon Research* 14:57 (1994) and immunomodulatory activity (Nieroda *et al.*, *Molec. Cell. Differentiation* 4: 335-351 (1996)).

Recombinant IFN α polypeptide has been approved for use in humans for hairy cell leukemia, AIDS-related Kaposi's sarcoma, malignant melanoma, chronic hepatitis B and C, chronic myelogenous leukemia, and condylomata acuminata (Baron, S. *et al.*, *JAMA* 266:1375 (1991)). However, for each of these indications, IFN α polypeptide must be administered repeatedly, often on a daily basis, for extended periods of time to maintain effective serum levels

due to the short half-life (hours) of the polypeptide in the serum (Friedman, *Interferons: A Primer*, Academic Press, New York, pp. 104-107 (1981); Galvani and Cawley, *Cytokine Therapy*, Cambridge University Press, Cambridge, pp. 114-115 (1992)). Thus, in spite of producing clinical benefit
5 for many disease conditions, the use of IFN α polypeptide is associated with acute and chronic side effects in most patients (Jones, *Cancer* 57: 1709-1715 (1986); and Quesda *et al.*, *Blood* 68: 493-497 (1986)). The severity of the adverse reaction correlates with peak serum interferon levels.

Viral or plasmid vectors containing IFN α genes have been used in *ex vivo* therapy to treat mouse tumors. For example, tumor cells were transfected
10 *in vitro* with viral or plasmid vectors containing an IFN α gene, and the transfected tumor cells were injected into mice (Belldgrun, A., *et al.*, *J. Natl. Cancer Inst.* 85: 207-216 (1993); Ferrantini, M. *et al.*, *Cancer Research* 53: 1107-1112 (1993); Ferrantini, M. *et al.*, *J. Immunology* 153: 4604-4615
15 (1994); Kaido, T. *et al.*, *Int. J. Cancer* 60: 221-229 (1995); Ogura, H. *et al.*, *Cancer Research* 50: 5102-5106 (1990); Santodonato, L., *et al.*, *Human Gene Therapy* 7:1-10 (1996); Santodonato, L., *et al.*, *Gene Therapy* 4:1246-1255 (1997)). In another *ex vivo* study, cervical carcinoma and leukemia cells were transfected with a viral vector containing the interferon-consensus gene, and
20 the transfected cells were injected into mice (Zhang, J.-F. *et al.*, *Cancer Gene Therapy* 3: 31-38 (1996)). In all of these *ex vivo* studies, varying levels of anti-tumor efficacy, such as tumor regression and/or prolonged survival, have been observed.

Viral or plasmid vectors containing interferon genes have also been
25 used in *in vivo* therapy for tumor-bearing mice. For example, a viral vector containing the interferon-consensus gene was injected into mice bearing transplanted MDA-MB-435 breast cancer, hamster melanoma, or K562 leukemia, and tumor regression was reported (Zhang, J.-F. *et al.*, *Proc. Natl. Acad. Sci. USA* 93: 4513-4518 (1996)). In a similar study, a plasmid vector
30 containing human IFN β gene complexed with cationic lipid was injected intracranially into mice bearing a human glioma, and tumor regression was reported (Yagi, K. *et al.*, *Biochemistry and Molecular Biology International*

32: 167-171 (1994)). In a murine model of renal cell carcinoma the direct intratumoral injection of an IL-2 plasmid DNA : lipid complex has been shown to result in complete tumor regression and a significant induction of a tumor specific CTL response increase in survival (Saffran *et al.*, *Cancer Gene Therapy* 5: 321-330 (1998)).

Plasmid vectors containing cytokine gens have also been reported to result in systemic levels of the encoded cytokine and in some cases, biological effects characteristic of each cytokine in mice. For example, the intramuscular injection of plasmid DNA encoding either TGF β , IL-2, IL-4, IL-5, or IFN α resulted in physiologically significant amounts in the systemic circulation of the corresponding cytokine polypeptide (Raz, E. *et al.*, *Proc. Natl. Acad. Sci. USA* 90: 4523-4527 (1993); Raz, E. *et al.*, *Lupus* 4: 266-292 (1995); Tokui, M. *et al.*, *Biochem. Biophys. Res. Comm.* 233: 527-531 (1997); Lawson, C. *et al.*, *J. Interferon Cytokine Res.* 17: 255-261 (1997); Yeow, W.-S. *et al.*, *J. Immunol.* 160: 2932-2939 (1998)).

U.S. Patent No. 5,676,954 reports on the injection of genetic material, complexed with cationic liposomes carriers, into mice. U.S. Patent Nos. 4,897,355; 4,946,787; 5,049,386; 5,459,127; 5,589,466; 5,693,622; 5,580,859; 5,703,055; and International Patent Application No. PCT/US94/06069 (publication no. WO 04/9469) provide cationic lipids for use in transfecting DNA into cells and mammals. U.S. Patent Nos. 5,589,466, 5,693,622, 5,580,859, 5,703,055, and international patent application no. PCT/US94/06069 (publication no. WO 04/9469) provide methods for delivering DNA-cationic lipid complexes to mammals.

Even though some viral vectors used in *ex vivo* and *in vivo* cancer therapy in murine models showed anti-tumor efficacy, the use of viral vectors to deliver interferon-expressing genes *in vivo* could induce anti-viral immune responses or result in viral integration into host chromosomes, causing disruption of essential host genes or activation of oncogenes (Ross *et al.*, *Human Gene Therapy* 7: 1781-1790 (1996)).

For treatment of multiple metastatic carcinomas of a body cavity are treated using laparoscopy (Childers *et al.*, *Gynecol. Oncol.* 59: 25-33, (1995)),

catheterization (Naumann *et al*, *Gynecol. Oncol.* 50: 291-3, (1993)) or other access devices (Almadrones *et al*, *Semin. Oncol. Nurs.* 11: 194-202, (1995)). Treatment is usually by surgical removal of primary and large metastatic tumors and postoperative chemotherapy (Kigwawa *et al*, *Am. J. Clin. Oncol.* 17: 230-3, (1994); Markman *et al*, *J. Clin. Oncol.* 10: 1485-91, (1992)) or radiotherapy (Fjeld *et al*, *Acta. Obstet. Gynecol. Scand. Suppl.* 155: 105-11, (1992)). Tumor recurrence is monitored by magnetic resonance imaging (Forstner *et al*, *Radiology* 196: 715-20, (1995)), ascites cytology (Clement, *Am. J. Clin. Pathol.* 103: 673-6, (1995); Forstner *et al*, *Radiology* 196: 715-20, 1995) and blood analyses (Forstner *et al*, *Radiology* 196: 715-20, (1995)). Many intraperitoneal (i.p.) cancers, such as ovarian cancer, eventually metastasize via the lymphatic system to the lungs or other vital organs, and the prognosis for the patient is very poor (Kataoka *et al*, *Nippon Sanka Fujinka Gakkai Zasshi* 46: 337-44, 1994; Hamilton, *Curr. Probl. Cancer* 16: 1-57, (1992)).

Human ovarian cancer is often diagnosed at an advanced stage when the effectiveness of surgery and chemotherapy are limited. The lack of effective treatment options for late-stage patients warrants the development of new treatment modalities for this disease. There have been several attempts to develop an effective immunotherapy for the treatment of ovarian cancer.

The early work in this area involved mouse studies in which bacteria-derived immunostimulants, such as *Bacillus Calmette-Guerin* (BCG) and *Corynebacterium parvum*, were injected i.p. as non-specific activators of the immune system. (Knapp and Berkowitz, *Am. J. Obstet. Gynecol.*, 128: 782-786, (1977); Bast *et al.*, *J. Immunol.*, 123: 1945-1951, (1979); Vanhaelen, *et al.*, *Cancer Research*, 41: 980-983, (1981); and Berek, *et al.*, *Cancer Research*, 44, 1871-1875, (1984)). These studies generally resulted in a non-specific immune response that often did not prevent the growth of later tumors. In addition, if the bacterial antigens were injected more than 24 hours after tumor cell inoculation, there was minimal antitumor response, suggesting that treatment of late-stage ovarian cancer patients with this type of therapy would not be effective.

More recent studies in both mice and humans have involved the i.p. or intravenous (i.v.) administration of cytokine proteins as more specific activators of the immune response (Adachi, *et al.*, *Cancer Immunol. Immunother.* 37: 1-6, (1993); Lissoni, *et al.*, *Tumori* 78: 118-20, (1992)).

5 Treating murine ovarian tumors with a combination of recombinant IL-2 and GM-CSF proteins had some beneficial effect in inhibiting ascites production; however, IL-2 was only effective if it was combined with GM-CSF (Kikuchi, *et al.*, *Cancer Immunol. Immunother.*, 43: 257-261, (1996)). Similarly, a combination of IL-2 and lymphokine-activated killer (LAK) cells was able to
10 reduce i.p. sarcomas in mice, while IL-2 protein alone was not as effective (Ottow, *et al.*, *Cellular Immunology*, 104: 366-376, (1987)). Human clinical trials evaluating IL-2 protein therapy of ovarian cancer patients indicated some antitumor effects (Chapman *et al.*, *Investigational New Drugs*, 6:179-188, (1988); West *et al.*, *N. Engl. J. Med.* 316:898-905, 1987; Lotze *et al.*, *Arch.*
15 *Surg.* 121:1373-1379, 1986; Benedetti Panici *et al.*, *Cancer Treatment Review*, 16A:123-127, 1989; Beller *et al.*, *Gynecol. Oncol.*, 34:407-412, 1989; Urba *et al.*, *J. Natl. Cancer Inst.*, 81:602-611, 1989; Stewart *et al.*, *Cancer Res.*, 50:6302-6310, 1990; Steis *et al.*, *J. Clin. Oncol.*, 8:1618-1629, 1990; Lissoni *et al.*, *Tumori*, 78:118-120, 1992; Sparano *et al.*, *J. of Immunotherapy*, 16:216-
20 223, 1994; Freedman *et al.*, *J. of Immunotherapy*, 16:198-210, 1994; Edwards *et al.*, *J. Clin. Oncol.*, 15:3399-3407, 1997).

Recent studies in mice have involved the injection of DNA constructs encoding "suicide" genes followed by treatment with prodrugs. This approach has successfully caused regression of some small tumors but has been less
25 successful on larger tumor masses. (Szala, *et al.* *Gene Therapy* 3: 1025-1031, 1996; Sugaya, *et al.* *Hum Gene Ther* 6: 317-323 (1996)). In another study, liposome-mediated *E1A* gene therapy for mice bearing ovarian cancers that overexpress HER-2/*neu* resulted in reduced mortality among these tumor bearing mice. (Yu, *et al.* *Oncogene*, 11: 1383-1388 (1995)). Similarly, the
30 successful treatment of murine ovarian carcinoma (MOT) has been demonstrated using cisplatin-induced gene transfer of DNA constructs encoding IFN γ via i.p. injection. (Son, *Cancer Gene Therapy* 4: 391-396

(1997)). However, this study demonstrated that tumors were poorly responsive to either the IFN γ gene or cisplatin alone, suggesting that the effectiveness of the cisplatin-based gene therapy protocol was mainly due to enhanced sensitization of cisplatin-exposed tumor cells to transfection by the IFN γ gene. (Son, *Cancer Gene Therapy* 4: 391-396, 1997).

Clearly, there is a need for superior therapeutic compositions and methods for treating mammalian cancer. Further, there is a need for an *in vivo* delivery system for IFN ω . The present invention provides a simple and safe yet effective compositions and methods for treatment of mammalian cancer.

The present invention also solves the problems inherent in prior attempts to treat body cavity malignancies. The inventors show herein that the malignant cell dissemination into body cavities, such as into the peritoneal cavity during late stage ovarian cancer, can be suppressed simply by administering as few as two to six doses of a polynucleotide formulation directly into the body cavity. This treatment results in selective transfection of malignant cells, and subsequent long-term local production of an effective amount of therapeutic molecules.

Summary of the Invention

The present invention is broadly directed to treatment of cancer by administering *in vivo*, into a tissue of a mammal suffering from cancer, a polynucleotide construct comprising a polynucleotide encoding a cytokine. The polynucleotide construct is incorporated into the cells of the mammal *in vivo*, and a therapeutically effective amount of a cytokine is produced *in vivo*, and delivered to tumor cells. Combinations of cytokine-encoding polynucleotides can be administered.

The present invention provides a pharmaceutical composition comprising about 1 ng to 20 mg of a non-infectious, non-integrating polynucleotide construct comprising a polynucleotide selected from the group

consisting of (a) a polynucleotide that hybridizes under stringent conditions to the nucleotide sequence of SEQ ID No. 7 or the complement thereof, wherein the polynucleotide sequence encodes a polypeptide that has antiproliferative activity when added to NIH-OVCAR3 cells *in vitro*; (b) a polynucleotide that encodes a polypeptide comprising an amino acid sequence which, except for at least one but not more than 20 amino acid substitutions, deletions, or insertions, is identical to amino acids -23 to 172 or 1 to 172 in SEQ ID No. 8, wherein the polypeptide has antiproliferative activity when added to NIH-OVCAR3 cells *in vitro*; and (c) a polynucleotide that encodes a polypeptide comprising amino acids 86-172 in SEQ ID No. 8, wherein the polypeptide has antiproliferative activity when added to NIH-OVCAR3 cells *in vitro*; and any of the above group complexed with one or more cationic compounds selected from the group consisting of cationic lipids, cationic peptides, cationic proteins, cationic polymers, and mixtures thereof.

The present invention also provides a pharmaceutical composition obtained by complexing a polynucleotide selected from the group consisting of (a) a polynucleotide that hybridizes under stringent conditions to the nucleotide sequence of SEQ ID No. 7 or the complement thereof, wherein the polynucleotide sequence encodes a polypeptide that has antiproliferative activity when added to NIH-OVCAR3 cells *in vitro*; (b) a polynucleotide that encodes a polypeptide comprising an amino acid sequence which, except for at least one but not more than 20 amino acid substitutions, deletions, or insertions, is identical to amino acids -23 to 172 or 1 to 172 in SEQ ID No. 8, wherein the polypeptide has antiproliferative activity when added to NIH-OVCAR3 cells *in vitro*; (c) a polynucleotide that encodes a polypeptide comprising amino acids 86-172 in SEQ ID No. 8, wherein the polypeptide has antiproliferative activity when added to NIH-OVCAR3 cells *in vitro*, with one or more cationic compounds selected from the group consisting of cationic lipids, cationic peptides, cationic proteins, cationic polymers, and mixtures thereof.

The present invention also provides a method of treating cancer in a mammal, comprising administering into a tissue of the mammal a non-

infectious, non-integrating polynucleotide construct comprising a polynucleotide encoding a cytokine, or an active fragment thereof, such that the polynucleotide is expressed *in vivo*, and such that the cytokine, or active fragment thereof, is delivered systemically to a tumor tissue in an amount effective to treat the cancer.

The present invention also provides a method of treating cancer in a mammal, comprising administering into a tissue of the mammal a non-infectious, non-integrating polynucleotide construct comprising a polynucleotide encoding a cytokine selected from the group consisting of interferon- ω , interferon- α , and a combination thereof, such that the polynucleotide or an active fragment thereof is expressed, and such that the cytokine is delivered locally to a tumor tissue in an amount effective to treat the cancer. Preferably, the polynucleotide construct is complexed with a cationic vehicle, more preferably, the cationic vehicle may be a cationic lipid, and most preferably, the cationic lipid may be mixed with a neutral lipid.

Another object of the invention is to provide a method of selectively transfecting malignant cells in a body cavity of a tumor-bearing mammal, comprising administering into the body cavity at least one non-infectious, non-integrating polynucleotide complexed with a cationic vehicle, such that the polynucleotide is expressed substantially in the malignant cells of the body cavity. Preferably, the cationic vehicle comprises one or more cationic lipids, and more preferably, the cationic vehicle comprises a cationic and neutral lipid mixture. In a preferred embodiment, the present invention is used to suppress peritoneal dissemination of malignant cells in a tumor-bearing mammal. In particular, the mammal may have ovarian cancer, or metastasis of ovarian cancer. Preferred polynucleotides may encode cytokines, or active fragments thereof. Most preferably, the polynucleotide may encode IL-2, or an active fragment thereof.

Compared to injection of recombinant cytokine polypeptides, the methods described herein have several important advantages. The present invention shows that *in vivo* transfection of cells with encoding polynucleotide, such as an IL-2 or IFN ω , results in serum levels of the

corresponding cytokine that have therapeutic effects, and yet are lower than the maximal serum levels typically required when cytokine polypeptides are injected. Further, injecting frequent high doses of cytokine polypeptides can produce debilitating side effects. The methods of the present invention provide cytokine therapy requiring less frequent injections of cytokine-encoding nucleic acids. The injection of polynucleotide constructs encoding cytokines produces sustained, low levels of biologically active cytokines that have beneficial effects, while minimizing adverse side effects.

Compared to the delivery of cytokine genes via a viral gene delivery vectors, the present method also has important advantages. Injection of non-viral vectors of the present method does not induce significant toxicity or pathological immune responses, as described, for example, in mice, pigs or monkeys (Parker, *et al.*, *Human Gene Therapy* 6: 575-590 (1995); and San, *et al.*, *Human Gene Therapy* 4: 781-788 (1993)). Thus, a non-viral vector is safer and can be repeatedly injected.

Brief Description of the Figures

A more complete appreciation of the invention and many of the attendant advantages thereof will be readily obtained as the same become better understood by reference to the following detailed description when considered in connection with the accompanying figures.

FIG. 1 shows the plasmid map of VR4151 (SEQ ID No. 4). The cytomegalovirus immediate-early gene promoter enhancer and 5' untranslated sequences (5'UTR + intron A) drive the expression of the human interferon ω coding sequence. The transcriptional terminator region includes polyadenylation and termination signals derived from the rabbit β -globin gene.

FIG. 2 shows the pharmacokinetics of hIFN ω in the serum of C57BL/6 mice (FIG. 2A) and nude mice (FIG. 2B) after a single intramuscular (i.m.) injection of hIFN ω plasmid DNA (VR4151). Mice were injected i.m. with 100 μ g of VR4151. Following the intramuscular

injection, mice were bled daily, and serum was collected and assayed for hIFN γ polypeptide using an ELISA. Each point represents an average of four mice. In C57BL/6 mice, the single i.m. injection resulted in peak serum levels of 254 pg/ml on day 6 after injection, and serum levels were still detectable 14 days after injection (50 pg/ml) (FIG. 2A). In nude mice, the single i.m. injection resulted in peak serum levels of 648 pg/ml on day 7, and serum levels were still detectable 14 days after injection (134 pg/ml) (FIG. 2B).

FIG. 3 shows that systemic mIFN α treatment reduces tumor volume (FIGs. 3A, 3C, and 3E) and increases survival (FIGs. 3B, 3D, and 3F) in three murine tumor models. C57BL/6 mice bearing subcutaneous B16F10 melanoma (FIGs. 3A and 3B), subcutaneous glioma 261 (FIGs. 3C and 3D), or DBA/2 mice bearing subcutaneous Cloudman melanoma (FIGs. 3E and 3F) were injected with 100 μ g either of VR4111 (mIFN α plasmid) or VR1055 (control plasmid), twice per week for three weeks, beginning on day 4 after tumor cell injection (n=8-10 mice per group).

FIG. 4 shows that systemic mIFN α , mIL-2 or mIL-12 plasmid DNA treatment reduces tumor volume (FIG. 4A) and mIFN α or mIL-12 plasmid DNA treatment increases survival (FIG. 4B) in the subcutaneous B16F10 melanoma model. C57BL/6 mice bearing subcutaneous B16F10 melanoma were injected with 100 μ g of VR4111 (mIFN α), VR4001 (mIL-12), VR1110 (mIL-2), or VR1012 (control plasmid) (n = 15-16 mice per group) twice per week for three weeks.

FIG. 5 shows that i.m. administration of hIFN γ pDNA reduces tumor volume (FIG. 5A) and increases survival (FIG. 5B) in nude mice bearing human A431 epidermoid carcinoma tumors. Mice bearing human A431 tumors between 30-80 mm³ were injected i.m. with 200 μ g of either VR4151 (hIFN γ plasmid) or VR1055 (control plasmid) twice per week for three weeks (n=15).

FIG. 6 shows that i.m. administration of mIFN α pDNA reduces B16F10 melanoma lung metastases in C57BL/6 mice. Mice bearing lung

metastases of B16F10 melanoma were injected i.m. with 100 µg of either VR4111 or VR1055 twice per week for three weeks, beginning on day 4 after tumor cell injection (n=10 mice per group). "TNTC" means too numerous to count as seen in the control group.

5 FIG. 7 shows that i.m. administration of mIFN α pDNA reduces intradermal M5076 primary tumor growth (FIG. 7A) as well as liver metastases (FIG. 7B) in C57BL/6 mice bearing murine M5076 reticulum cell sarcoma cells. Mice bearing M5076 tumors were injected i.m. with 100 µg of either VR4111 or VR1055 twice per week for three weeks, beginning on day 4
10 after tumor cell inoculation (n=10-13 mice per group).

 FIG. 8 shows a comparison of different dosages and frequencies of mIFN α pDNA administration in the subcutaneous B16F10 melanoma model. C57BL/6 mice bearing subcutaneous B16F10 melanoma were injected i.m. with 50 µg or 100 µg of either VR4111 or VR1055 twice a week for 3 weeks
15 beginning 4 days after tumor cell inoculation (n=10 mice per group). All groups treated with 100 µg of VR4111 showed significant reduction in tumor growth by day 21 (p=0.002) and significant enhancement in survival (p<0.008) with all treatments tested (FIGs. 8A and 8B). In mice treated with 50 µg VR4111, tumor growth was significantly reduced by day 21 (p=0.005),
20 and survival was significantly increased (p<0.003) in the groups of mice that were injected twice per week or once per week. The group injected every other week with 50 µg VR4111 was not significantly different from the mice that received the control plasmid (FIGs. 8C and 8D).

 FIG. 9 shows the results of experiments performed to determine the
25 role of NK and T cells in the antitumor response induced by mIFN α plasmid DNA. Nude mice (T cell deficient) (FIGs. 9A and 9B), and beige-nude mice (NK and T cell deficient) (FIGs. 9C and 9D) bearing subcutaneous B16F10 melanoma tumors were injected i.m. with 100 µg of either VR4111 or VR1055 twice per week for three weeks, beginning on day 4 after tumor cell
30 injection (n=15 mice per group). No significant reduction in tumor volume or increase in survival was found for nude or nude-beige mice treated with

VR4111, suggesting that T cells are involved in the mIFN α antitumor response.

FIG. 10 shows the results of experiments performed to evaluate the role of CD4⁺ and CD8⁺ T cells in the mIFN α DNA antitumor response. For depletion of CD4⁺ and CD8⁺ T cells, C57BL/6 mice bearing subcutaneous B16F10 melanoma tumors were injected i.p. with 500 μ g of either the anti-CD4 mAb (clone GK1.5, rat IgG) (ATCC, Rockville, MD) or anti-CD8 mAb (clone 2.43, rat IgG) (ATCC, Rockville, MD) one day after each i.m. injection of 100 μ g of either VR4111 or VR1055 twice per week for three weeks (n = 10 mice per group). The mIFN α plasmid DNA therapy significantly reduced tumor growth ($p \leq 0.002$) and enhanced survival ($p \leq 0.008$) of both normal mice and mice depleted of CD4⁺ T cells, suggesting that CD4⁺ T cells were not required for the response. In contrast, mice depleted of CD8⁺ T cells and injected with VR4111 had tumor volumes and survival that were not significantly different from mice treated with the control plasmid DNA, indicating a requirement for CD8⁺ T cells in the antitumor response.

FIG. 11 shows that intratumoral hIFN ω (VR4151) and hIFN α (VR4112) treatment reduces tumor volume in the human A375 melanoma model (FIG. 11A) and human NIH-OVCAR3 (FIG. 11B) in nude mice. Mice bearing subcutaneous tumor received direct intratumoral injections of a complex of DNA:DMRIE/DOPE (1:1 DNA:lipid mass ratio, 100 μ g of plasmid DNA) for 6 consecutive days followed by an additional 5 treatments every other day for a total of 11 injections (A375 melanoma model), or for every other day for a total of 11 injections (NIH-OVCAR3 ovarian cancer model).

FIG. 12 shows that intratumoral mIFN α (VR4101) plasmid DNA treatment reduces tumor volume (FIG. 12A) and increases survival (FIG. 12B) in the subcutaneous B16F10 melanoma model in C57BL/6 mice. Mice received a subcutaneous implantation of 10^4 B16F10 cells into the flank. Beginning at day 12 post tumor implant, mice received six consecutive

intratumoral injections of a complex of pDNA:DMRIE/DOPE (1:1 DNA:DMRIE mass ratio, 100 µg of plasmid DNA).

FIG. 13 shows luciferase activity in peritoneal tissues and MOT ascites in mice after i.p. injection of luciferase DNA:lipid complex. The results show high levels of reporter gene expression in ascites but low levels in peritoneal tissue. MOT tumor-bearing C3H/HeN mice received i.p. injections of a complex of pDNA:DMRIE/DOPE (1:1 DNA:DMRIE mass ratio, 100 µg of plasmid DNA) on days 5 and 6 after tumor cell implant. Tissues were collected 1 day (FIG. 13A) or 3 days (FIG. 13B) following the DNA:lipid injection.

FIG. 14 shows serum levels of IL-2 after i.p. injection of either IL-2 pDNA or protein in MOT tumor bearing mice. The serum levels of IL-2 were much lower than levels in ascites. Ascites and serum were collected at 4 hours and days 1, 2, 3, 6 and 10 post DNA or protein injection (5 mice for each time point), and analyzed for mIL-2 polypeptide using an ELISA.

FIG. 15 shows a significant reduction in MOT tumor growth ($p=0.01$) (FIG. 15A) and increased survival ($p=0.04$) (FIG. 15B) of mice treated with i.p. injection of IL-2 pDNA:lipid on days 5-10 after tumor cell injection. The DNA was complexed at either a 1:1 (15A and 15B) or 5:1 (FIGS. 15C and 15D) DNA:DMRIE mass ratio (100 µg pDNA). Plasmid DNA without lipid was not effective (FIGs. 15E and 15F).

FIG. 16 shows that i.p. mIL-2 plasmid DNA (VR1110):lipid treatment inhibits tumor growth (FIG. 16A) and enhances survival (FIG. 16B) in the MOT tumor model in C3H/HeN mice. MOT tumor-bearing mice received three alternative-day i.p. injections of a complex of pDNA:DMRIE/DOPE (1:1 DNA:DMRIE mass ratio, 100 µg of plasmid DNA).

FIG. 17 shows a significant reduction in MOT tumor growth and increased survival of mice treated with i.p. injection of IL-2 DNA:lipid followed by debulking of tumor ascites. MOT tumor-bearing mice received six consecutive intraperitoneal injections of a complex of pDNA:DMRIE/DOPE (1:1 DNA:DMRIE mass ratio, 100 µg of pDNA) and

debulked of 5 ml of tumor ascites 4 days after the last DNA:lipid injection (n=10).

FIG. 18 shows dose-response of mIL-2 pDNA (VR1110):lipid treatment in the MOT tumor model. C3H/HeN mice bearing MOT tumor were injected with 25, 50 or 100 μ g of VR1110:DMRIE/DOPE on days 5, 8 and 11 after MOT tumor cell injection. In mice treated with 50 or 100 μ g of VR1110, tumor growth was significantly reduced ($p=0.002$) and survival significantly enhanced ($p=0.01$) by day 15 post tumor cell inoculation compared to the control. Tumor-bearing mice treated with 25 μ g of VR1110:lipid were not significantly different from the control mice for either tumor volume or survival (n=15).

FIG. 19 shows the cytokine profile of ovarian tumor ascites in C3H/HeN mice MOT tumor model following mIL-2 pDNA (VR1110):lipid treatment. Mice received i.p. injections of a complex of pDNA/DMRIE/DOPE (1:1 DNA/DMRIE mass ratio, 100 μ g of plasmid DNA) on days 5, 8 and 11 after tumor cell implant. Two days after each injection, mice were sacrificed (5 mice for each time point), and the ascites were collected and analyzed for cytokine concentration. The level of IL-2 (days 7, 10 and 13) as well as IFN γ and GM-CSF (days 10 and 13) were markedly elevated suggesting that IL-2 upregulates IFN γ and GM-CSF production.

FIG. 20 shows that i.p. mIFN α pDNA (VR4111):lipid treatment enhances survival (FIG. 20B) in the MOT tumor model in C3H/HeN mice. MOT tumor-bearing mice received three alternative-day i.p. injections of a complex of pDNA:DMRIE/DOPE (1:1 DNA:DMRIE mass ratio, 100 μ g of plasmid DNA).

Detailed Description of the Preferred Embodiments

The present invention is broadly directed to treatment of cancer by administering *in vivo*, into a tissue of a mammal suffering from cancer, at least one polynucleotide construct comprising at least one polynucleotide encoding at least one cytokine, or at least one active fragment thereof. The polynucleotide construct is incorporated into the cells of the mammal *in vivo*, and a therapeutically effective amount of a cytokine is produced *in vivo*, and delivered to tumor cells. Combinations of cytokine-encoding polynucleotides can be administered.

The present invention provides a pharmaceutical composition comprising about 1 ng to 20 mg of a non-infectious, non-integrating polynucleotide construct comprising a polynucleotide selected from the group consisting of (a) a polynucleotide that hybridizes under stringent conditions to the nucleotide sequence of SEQ ID No. 7 or the complement thereof, wherein the polynucleotide sequence encodes a polypeptide that has antiproliferative activity when added to NIH-OVCAR3 cells *in vitro*; (b) a polynucleotide that encodes a polypeptide comprising an amino acid sequence which, except for at least one but not more than 20 amino acid substitutions, deletions, or insertions, is identical to amino acids -23 to 172 or 1 to 172 in SEQ ID No. 8, and wherein the polypeptide has antiproliferative activity when added to NIH-OVCAR3 cells *in vitro*; and (c) a polynucleotide that encodes a polypeptide comprising amino acids 86-172 in SEQ ID No. 8, wherein the polypeptide has antiproliferative activity when added to NIH-OVCAR3 cells *in vitro*; and one or more cationic compounds selected from the group consisting of cationic lipids, cationic peptides, cationic proteins, cationic polymers, and mixtures thereof. The pharmaceutical composition can be used to practice all of the methods of the present invention.

The present invention also provides a pharmaceutical composition obtained by complexing a polynucleotide selected from the group consisting of (a) a polynucleotide that hybridizes under stringent conditions to the nucleotide sequence of SEQ ID No. 7 or the complement thereof, wherein the polynucleotide sequence encodes a polypeptide that has antiproliferative activity when added to NIH-OVCAR3 cells *in vitro*; (b) a polynucleotide that

encodes a polypeptide comprising an amino acid sequence which, except for at least one but not more than 20 amino acid substitutions, deletions, or insertions, is identical to amino acids -23 to 172 or 1 to 172 in SEQ ID No. 8, wherein the polypeptide has antiproliferative activity when added to NIH-OVCAR3 cells *in vitro*; and (c) a polynucleotide that encodes a polypeptide comprising amino acids 86-172 in SEQ ID No. 8, wherein the polypeptide has antiproliferative activity when added to NIH-OVCAR3 cells *in vitro*, with one or more cationic compounds selected from the group consisting of cationic lipids, cationic peptides, cationic proteins, cationic polymers, and mixtures thereof.

The pharmaceutical composition of the present invention can be a polynucleotide construct comprising a polynucleotide that hybridizes under stringent conditions to the nucleotide sequence of SEQ ID No. 7 or the complement thereof, wherein the polynucleotide sequence encodes a polypeptide that has antiproliferative activity when added to NIH-OVCAR3 cells *in vitro*, and one or more cationic compounds selected from the group consisting of cationic lipids, cationic peptides, cationic proteins, cationic polymers, and mixtures thereof. Alternatively, the pharmaceutical composition of the present invention can be a polynucleotide construct comprising a polynucleotide that encodes a polypeptide comprising an amino acid sequence which, except for at least one but not more than 20 amino acid substitutions, deletions, or insertions, is identical to amino acids -23 to 172 or 1 to 172 in SEQ ID No. 8, wherein the polypeptide has antiproliferative activity when added to NIH-OVCAR3 cells *in vitro*, and one or more cationic compounds selected from the group consisting of cationic lipids, cationic peptides, cationic proteins, cationic polymers, and mixtures thereof. Alternatively, the pharmaceutical composition of the present invention can be a polynucleotide construct comprising a polynucleotide that encodes a polypeptide comprising amino acids 86-172 in SEQ ID No. 8, wherein the polypeptide has antiproliferative activity when added to NIH-OVCAR3 cells *in vitro*; and one or more cationic compounds selected from the group

consisting of cationic lipids, cationic peptides, cationic proteins, cationic polymers, and mixtures thereof.

The pharmaceutical composition of the present invention comprises at least one polynucleotide construct comprising at least one polynucleotide encoding an IFN ω , or an active fragment thereof. Preferably, the polynucleotide construct contains a polynucleotide encoding a human IFN ω . More preferably, IFN ω is encoded by nucleotides 1 to 585 in SEQ ID No. 7 (corresponding to amino acids -23 to 172 in SEQ ID No. 8), or by nucleotides 70 to 585 in SEQ ID No. 7 (corresponding to amino acids 1 to 172 in SEQ ID No. 8). Most preferably, the polynucleotide construct is VR4151 in SEQ ID No. 4. The polynucleotide construct may be complexed with one or more cationic compounds selected from the group consisting of cationic lipids, cationic peptides, cationic proteins, cationic polymers, and mixtures thereof. Preferably, the polynucleotide construct is complexed with one or more cationic lipids. More preferably, the polynucleotide construct is complexed with one or more cationic lipids and one or more neutral lipids. Still more preferably, the cationic lipid is (\pm)-N-(2-hydroxyethyl)-N,N-dimethyl-2,3-bis(tetradecyloxy)-1-propaniminium bromide (DMRIE) and the neutral lipid is 1,2-dioleoyl-sn-glycero-3-phosphatidylethanolamine (DOPE) such that the mass ratio of polynucleotide construct to lipid is from about 10:1 and about 0.5:1. More preferably, the mass ratio of polynucleotide construct to lipid is from about 5:1 and about 1:1. Still more preferably, the mass ratio of polynucleotide construct to lipid is about 5:1.

Cytokine-encoding plasmids discussed herein include VR4102 (hIFN α in the VR1012 vector) (SEQ ID No. 1), VR4112 (hIFN α in the VR1055 vector) (SEQ ID No. 2), VR4150 (hIFN ω in the VR1012 vector) (SEQ ID No. 3), VR4151 (hIFN ω in the VR1055 vector) (SEQ ID No. 4), VR4101 (mIFN α in the VR1012 vector) (SEQ ID No. 5), VR4111 (mIFN α in the VR1055 vector) (SEQ ID No. 6), and VR1110 (mIL-2 in the VR1012 vector), VR1103 (hIL-2 in the VR1012 vector) (SEQ ID No. 25), VR4001 (mIL-12 in the VR1033 vector), and VR1700 (mGM-CSF in the VR1012 vector).

Cytokine-encoding cDNAs discussed herein include the cDNA for hIFN ω (SEQ ID No. 7), the cDNA for hIFN α (SEQ ID No. 9), the cDNA for mIFN α (SEQ ID No. 11), the cDNA for hIL-2 (SEQ ID No. 13 and the coding portion of SEQ ID No. 25), the cDNA for mL-2 (for example, as disclosed in Kashima *et al.*, *Nature* 313:402-404 (1985), which is hereby incorporated by reference) the cDNA for mL-12 (for example, as disclosed in Tone *et al.*, *Eur. J. Immunol.* 26:1222-1227(1996), which is hereby incorporated by reference), and the cDNA for mGM-CSF (for example, as disclosed in Gough *et al.*, *EMBO J.* 4:645-653 (1985), which is hereby incorporated by reference). Cytokine polypeptides discussed herein include hIFN ω (SEQ ID No. 8), hIFN α (SEQ ID No. 10), mIFN α (SEQ ID No. 12), and hIL-2 (SEQ ID No. 14 and SEQ ID No. 26).

By "stringent conditions" is intended a hybridization by overnight incubation at 42°C in a solution comprising: 50% formamide, 5x SSC (750 mM NaCl, 15 mM trisodium citrate), 50 mM sodium phosphate (pH 7.6), 5x Denhardt's solution, 10% dextran sulfate, and 20 μ g/ml denatured, sheared salmon sperm DNA, followed by repeatedly washing the filters (at least three times) in 0.1x SSC and 0.1% sodium dodecyl sulfate (w/v) for 20 minutes at about 65°C.

By "active fragment" is intended a fragment of a cytokine that displays the antiproliferative activity of the mature or full length cytokine. For example, a full length hIFN ω is set forth in amino acids -23 to 172 of SEQ ID No. 8. The corresponding mature hIFN ω is set forth in amino acids 1 to 172 of SEQ ID No. 8. Active fragments of hIFN ω include, but are not limited to a polypeptide comprising amino acids 86-172 in SEQ ID No. 8, a polypeptide comprising amino acids 61-172 in SEQ ID No. 8, a polypeptide comprising amino acids 41-172 in SEQ ID No. 8, and a polypeptide comprising amino acids 21-172 in SEQ ID No. 8. A full length hIFN α is set forth in amino acids -23 to 166 of SEQ ID No. 10. The corresponding mature hIFN α is set forth in amino acids 1 to 166 of SEQ ID No. 10. Active fragments of hIFN α include, but are not limited to a polypeptide comprising amino acids 83-166 in SEQ ID

No. 10, a polypeptide comprising amino acids 61-166 in SEQ ID No. 10, a polypeptide comprising amino acids 41-166 in SEQ ID No. 10, and a polypeptide comprising amino acids 21-166 in SEQ ID No. 10. Full length hIL-2 is set forth in amino acids -20 to 133 of SEQ ID No. 14. The corresponding mature hIL-2 is set forth in amino acids 1 to 133 of SEQ ID No. 14. Active fragments of hIL-2 include, but are not limited to a polypeptide comprising amino acids 58 to 105 in SEQ ID No. 14, and a polypeptide comprising amino acids 20 to 126 in SEQ ID No. 14.

Assays of antiproliferative activity *in vitro* are well known to those of ordinary skill in the art. For example, one antiproliferation assay that can be used is to treat cultured cells, such as human ovarian NIH-OVCAR3 cells (ATCC, Rockville, MD), with supernatants from human melanoma UM449 cells transfected with the polynucleotide construct containing a polynucleotide encoding an IFN γ or an active fragment thereof. In this antiproliferation assay, NIH-OVCAR3 cells are cultured and plated in 96 well-tissue culture plates. The plates are incubated for 24 hours at 37°C in a humidified 5% CO₂ atmosphere. Twenty μ l of tissue culture supernatants from transfected UM449 cells are added to duplicate wells. An interferon reference standard (*e.g.*, human leukocyte interferon, Sigma Chemical Co., St. Louis, MO) is included in each assay. The cells are incubated with the test samples or the interferon standard for an additional 72 hours at 37°C. To quantitate the effects on cell proliferation, 50 μ l of XTT/ECR substrate (Cell Proliferation Kit, Boehringer Mannheim, Indianapolis, IN) is added to each well and the plates are incubated for an additional 24 hours at 37°C prior to measurement of the OD₄₉₀. Other cell lines can be used in the antiproliferation assay. For example, any of the cells listed on Table I can be used. Another antiproliferation assay that can be used is provided in Nieroda, *et al* (*Mol. Cell. Differentiation* 4: 335-351 (1996)).

For treatment of cancer, a polynucleotide construct comprising a polynucleotide encoding a cytokine can be delivered locally, systemically or intra-cavity. In the "systemic delivery" embodiment of the invention, one or more polynucleotide construct comprising one or more polynucleotide

encoding one or more cytokine is administered into a tissue such that the polynucleotide is expressed as the cytokine *in vivo* and the cytokine is released into the circulation, and such that a therapeutically effective amount of the cytokine is systemically delivered to the tumor. In this embodiment, the polynucleotide construct can be administered within *ex vivo* cells or associated with *ex vivo* cellular material. Preferably, the cytokine is an IFN ω , IFN α , IFN τ , IFN γ , IFN β , IL-1, IL-2, IL-4, IL-7, IL-12, IL-15, IL-18, GM-CSF, or any combination of these, or any combination of one or more of these and one or more additional cytokines. More preferably, the cytokine is an IFN α , IFN ω , IL-2, or IL-12. Most preferably, the cytokine is an IFN α or IFN ω . Examples of the combination are a polynucleotide encoding an IFN ω and an IFN α ; a polynucleotide encoding an IFN ω and an IL-2; a polynucleotide encoding an IFN α and an IL-2; and a polynucleotide encoding an IFN ω , an IFN α , and an IL-2. More preferably, the polynucleotide construct contains a polynucleotide encoding an IFN ω and/or an IFN α . Even more preferably, the polynucleotide construct contains a polynucleotide encoding a human IFN ω and/or a human IFN α . Even more preferably, the polynucleotide encodes a human IFN ω . Preferably, the polynucleotide construct is administered free from *ex vivo* cells and free from *ex vivo* cellular material.

In this embodiment, administration can be into tissue including but not limited to muscle, skin, brain, lung, liver, spleen, bone marrow, thymus, heart, lymph nodes, blood, bone, cartilage, pancreas, kidney, gall bladder, stomach, intestine, testis, ovary, uterus, rectum, nervous system, eye, gland, or connective tissue. Preferably, the administration is into muscle tissue, *i.e.*, skeletal muscle, smooth muscle, or myocardium, and the polynucleotide construct is naked. Most preferably, the muscle is skeletal muscle. For polynucleotide constructs in which the polynucleotide encoding a cytokine is DNA, the DNA can be operably linked to a cell-specific promoter that directs substantial transcription of the DNA only in predetermined cells.

By "naked" is meant that the polynucleotide construct is free from association with any delivery vehicle known in the art that can act to facilitate

entry into cells, for example, from transfection-facilitating proteins, viral particles, liposomes, cationic lipids, and calcium phosphate precipitating agents.

As used herein, "ex vivo" cells are cells into which the polynucleotide construct is introduced, for example, by transfection, lipofection, electroporation, bombardment, or microinjection. The cells containing the polynucleotide construct are then administered *in vivo* into mammalian tissue. Such *ex vivo* polynucleotide constructs are well-known to those of ordinary skill in the art. For example, see Beldegrun, A., *et al.*, *J. Natl. Cancer Inst.* 85: 207-216 (1993); Ferrantini, M. *et al.*, *Cancer Research* 53: 1107-1112 (1993); Ferrantini, M. *et al.*, *J. Immunology* 153: 4604-4615 (1994); Kaido, T., *et al.*, *Int. J. Cancer* 60: 221-229 (1995); Ogura, H., *et al.*, *Cancer Research* 50: 5102-5106 (1990); Santodonato, L., *et al.*, *Human Gene Therapy* 7:1-10 (1996); Santodonato, L., *et al.*, *Gene Therapy* 4:1246-1255 (1997); and Zhang, J.-F. *et al.*, *Cancer Gene Therapy* 3: 31-38 (1996).

The polynucleotide construct is administered in a "cell-free" fashion when it is administered independently, *i.e.*, free of *ex vivo* cells or *ex vivo* cellular material.

In the "local cytokine delivery" embodiment of the present invention, a polynucleotide construct comprising a polynucleotide encoding IFN ω and/or IFN α is administered *in vivo* into or near a tumor of a mammal, such that the polynucleotide is incorporated into the cells of the tumor. Tumor cells subsequently express the interferon polypeptide in an amount effective to treat cancer.

In this embodiment, a polynucleotide construct comprising a polynucleotide encoding an IFN ω and/or an IFN α can be administered into the tumor. Alternatively, the polynucleotide construct can be administered into non-tumor cells surrounding a tumor, near a tumor, or adjacent to a tumor, such that a therapeutically effective amount of an IFN ω and/or an IFN α is produced *in vivo* near or within the tumor and is delivered to the malignant cells of the tumor. One way to provide local delivery of the polynucleotide construct is by administering intravenously a polynucleotide construct

comprising a tumor-targeted promoter, wherein the polynucleotide is incorporated into the cells of the tumor and the cytokine is expressed in the tumor in an amount effective to treat cancer. Preferably, the polynucleotide construct is administered into the tumor.

5 In the "intra-cavity delivery" embodiment, the present invention provides a method of selectively transfecting malignant cells in a tumor-bearing body cavity of a mammal by introducing a polynucleotide construct into the body cavity, wherein the polynucleotide is incorporated into tumor cells and the tumor cells subsequently express the protein encoded by the polynucleotide in an amount effective to treat cancer. The polynucleotide
10 construct is administered free from *ex vivo* cells and free from *ex vivo* cellular material.

A cavity is a space within the body that can confine a fluid volume for some period of time. The cavity can either be present in a normal animal, or it
15 can be produced as a result of disease, surgery or trauma. Cavities in the normal animal include the peritoneum, the cerebrospinal fluid space, the ventricles of the brain, the plural space around lung, the bronchiolar airways, the nasal sinus, the bladder, the vagina, the ear, the synovium of various joints (knee, hip etc.), the internal network of salivary gland tissue, and the
20 gastrointestinal tract including stomach. Surgical removal of tumor tissue can also produce a space which fits the definition of a cavity. An open wound produced by trauma or surgery and closed by suture can be defined as a cavity, and the area under a blister produced by an infection, abrasion or a burn also fits the definition.

25 There are special bioavailability considerations when a gene delivery system is administered into a cavity. First the fluid volume in the cavity can be substantially comprised of the vehicle in which the delivery system is suspended. Second, the delivery system can have particular access to cells that are either suspended in the cavity, or that are lining the surface of the cavity.
30 Third, in some cases normally differentiated cells that are lining the cavity may be embedded in an extracellular matrix and, may not be accessible to the delivery system. Thus, the delivery system may preferentially transfect cells

that are growing outside the normal extracellular matrix and avoid the cells that are growing within the extracellular matrix, conferring a kind of cell selectivity to the delivery system.

5 With respect to the first point, body fluids such as serum, have been shown to inhibit gene delivery systems. For example, the transfection activities of Lipofectin and LipofectAMINE are inhibited by serum. It is thought that serum factors bind to cationic lipid/DNA complexes and block their uptake into cells. In cavity models the endogenous fluid volume can be removed, the cavity can be washed, and the delivery system can be
10 administered into the cavity in a vehicle that is compatible with optimal gene delivery efficacy. Thus the cavity model allows the investigator to create a fluid environment which allows for optimal gene delivery potency.

With respect to the second point, cells that are either floating in the cavity or are lining the surface of the cavity have preferential access to the
15 delivery system and can be preferentially transfected relative to other cells in the body. Since the delivery system is confined within the cavity, peripheral cells in the body outside of the cavity will not be transfected. Thus, there is tissue targeting to the cells within the cavity. For example, gene delivery systems administered into the peritoneal cavity will have access to metastatic
20 tumor cells derived from colon or ovarian cancers that are floating in the peritoneum or are attached to the surfaces of the peritoneum. Delivery systems administered into the plural space should transfect cancer cells in the plural effusion. Delivery systems administered into the cerebral spinal fluid should have access to metastatic cancer cells present there.

25 With respect to the third point, differentiated cells that are present in normal tissues are often embedded into an extracellular matrix. This matrix can be difficult to penetrate with large particulate delivery systems. Some cells, such as poorly differentiated tumor cells, that are present in cavities can grow outside of the normal extracellular matrix and are therefore more
30 accessible to gene delivery systems. In this way the delivery system can preferentially transfect those cells that are growing outside of the extracellular matrix and not transfect those cells that are growing within the extracellular

matrix. This is another form of in vivo, cell type specific targeting. Examples of normal cells that are not embedded in an extracellular matrix and are therefore more accessible to gene delivery systems are, bronchial airway cells, lung cells in the plural space, and ependimal cells lining the surface of the ventricles of the brain. Normal bladder cells that line the surface of the bladder are embedded in a tight extracellular matrix and are therefore not readily accessible to a gene delivery system delivered into the bladder, but tumor cells which grow up and out of the extracellular matrix into the bladder vesicle are accessible to gene delivery systems administered into the bladder vesicle. Thus normal bladder tissue would be expected to resist transfection whereas, bladder tumor would be expected to be transfectable.

A preferred application of the intra-cavity delivery embodiment is in the treatment of peritoneally disseminated cancers. More specifically, a mammal bearing peritoneal tumor may be injected i.p. with an effective amount of a polynucleotide complexed with a lipid in a physiologically acceptable diluent in a total volume sufficient to access the entire body cavity. The mammal may have tumor ascites in the peritoneal cavity as in an ovarian cancer. In the most preferred application, this methodology may be used in treating ovarian cancer of a human.

Debulking of tumor ascites is commonly performed on human ovarian cancer patients. Debulking involves removal of tumor ascites from the peritoneal cavity. In humans bearing ovarian tumor ascites, the ascites fluid would be debulked by insertion of a catheter i.p. followed by periodic draining of ascites fluid. It is contemplated that the tumor ascites would be debulked before and/or after the i.p. administration of the polynucleotide formulation of the present invention.

Transfection efficacy of the intra-cavity delivery embodiment may be determined by collecting the tumor ascites and serum at various times after the injection and performing diagnostic assays appropriate for the encoded molecule(s). Naturally, other means of determining tumor mass, growth, and viability may also be used to assess the effectiveness of the present invention.

Preferred polynucleotides for the intra-cavity delivery embodiment may encode not only immunogenic molecules such as cytokines (*e.g.*, interleukins 1-18 and $\alpha/\beta/\gamma/\omega$ -interferons, colony stimulating factors, *e.g.*, G-CSF, GM-CSF, M-CSF, and tumor necrosis factors), but also chemokines (*e.g.*, C-X-C and C-C), Class I and II histocompatibility antigens, costimulatory molecules (*e.g.*, B7-1, B7-2, CAMs, and flt3 ligand), growth factors (*e.g.*, epidermal growth factors, fibroblast growth factors, transforming growth factors and growth hormone), and the like. The polynucleotide may also encode bacterial antigens, viral glycoproteins, enzymes (*e.g.*, lysozymes), recombinant antibodies, molecules that interfere with cellular adhesion, adhesion molecules, proliferation and vascular inhibitory factors, ribozymes, and antisense RNAs targeted toward key oncogenic or tumor growth proteins. Moreover, selective delivery of toxic peptides (*e.g.*, ricin, diphtheria toxin, or cobra venom factor) or proteins capable of synthesizing toxic compounds (*e.g.* thymidine kinase and cytosine deaminase) to the malignant cells may have therapeutic benefits. The polynucleotide may also comprise a tumor suppressor gene (*e.g.*, p53). Preferred polynucleotides encode cytokines. Preferred cytokines are IL-2, IFN ω , and IFN α . IL-2 is most preferred.

For treatment of cancer by any of the above disclosed embodiments, any polynucleotide encoding an IFN ω , or an active fragment thereof, can be used. For example, the polynucleotide construct can be a construct comprising a polynucleotide that hybridizes under stringent conditions to the nucleotide sequence of SEQ ID No. 7 or the complement thereof, wherein the polynucleotide sequence encodes a polypeptide that has antiproliferative activity when added to NIH-OVCAR3 cells *in vitro*, and one or more cationic compounds selected from the group consisting of cationic lipids, cationic peptides, cationic proteins, cationic polymers, and mixtures thereof. Alternatively, the construct can be a polynucleotide construct comprising a polynucleotide that encodes a polypeptide comprising an amino acid sequence which, except for at least one but not more than 20 amino acid substitutions, deletions, or insertions, is identical to amino acids -23 to 172 or 1 to 172 in SEQ ID No. 8, wherein the polypeptide has antiproliferative activity when

added to NIH-OVCAR3 cells *in vitro*, and one or more cationic compounds selected from the group consisting of cationic lipids, cationic peptides, cationic proteins, cationic polymers, and mixtures thereof. Alternatively, the construct can be a polynucleotide construct comprising a polynucleotide that encodes a polypeptide comprising amino acids 86-172 in SEQ ID No. 8, wherein the polypeptide has antiproliferative activity when added to NIH-OVCAR3 cells *in vitro*, and one or more cationic compounds selected from the group consisting of cationic lipids, cationic peptides, cationic proteins, cationic polymers, and mixtures thereof. Preferably, IFN α is encoded by nucleotides 1 to 585 in SEQ ID No. 7 (corresponding to amino acids -23 to 172 in SEQ ID No. 8), or by nucleotides 70 to 585 in SEQ ID No. 7 (corresponding to amino acids 1 to 172 in SEQ ID No. 8). More preferably, the polynucleotide construct is VR4151.

For treatment of cancer, any polynucleotide encoding IFN α , or active fragment thereof, can also be used. For example, the polynucleotide construct can be a construct comprising a polynucleotide that hybridizes under stringent conditions to the nucleotide sequence of SEQ ID No. 9 or the complement thereof, wherein the polynucleotide sequence encodes a polypeptide that has antiproliferative activity when added to NIH-OVCAR3 cells *in vitro*, and one or more cationic compounds selected from the group consisting of cationic lipids, cationic peptides, cationic proteins, cationic polymers, and mixtures thereof. Alternatively, the construct can be a polynucleotide construct comprising a polynucleotide that encodes a polypeptide comprising an amino acid sequence which, except for at least one but not more than 20 amino acid substitutions, deletions, or insertions, is identical to amino acids -23 to 166 or 1 to 166 in SEQ ID No. 10, wherein the polypeptide has antiproliferative activity when added to NIH-OVCAR3 cells *in vitro*, and one or more cationic compounds selected from the group consisting of cationic lipids, cationic peptides, cationic proteins, cationic polymers, and mixtures thereof. Alternatively, the construct can be a polynucleotide construct comprising a polynucleotide that encodes a polypeptide comprising amino acids 83-166 in SEQ ID No. 10, wherein the polypeptide has antiproliferative activity when

added to NIH-OVCAR3 cells *in vitro*; and one or more cationic compounds selected from the group consisting of cationic lipids, cationic peptides, cationic proteins, cationic polymers, and mixtures thereof. Preferably, IFN α is encoded by nucleotides 1 to 567 in SEQ ID No. 9 (corresponding to amino acids -23 to 166 in SEQ ID No. 10), or by nucleotides 1 to 567 in SEQ ID No. 9 (corresponding to amino acids 1 to 166 in SEQ ID No. 10). Preferably, the polynucleotide construct is VR4112.

For polynucleotide constructs that do not contain a polynucleotide encoding IFN ω , the polynucleotide construct is preferably a cell-free construct. For polynucleotide constructs that contain a polynucleotide encoding IFN ω , the polynucleotide construct can be administered either within *ex vivo* cells or free of *ex vivo* cells or *ex vivo* cellular material. Preferably, the polynucleotide construct is administered free of *ex vivo* cells or *ex vivo* cellular material.

In the "local delivery" and "intra-cavity delivery" embodiments, the polynucleotide construct is preferably complexed with one or more cationic compounds. More preferably, the polynucleotide construct is complexed with one or more cationic lipids by ionic interaction. Generally, the complex then contacts the cell membrane and is transfected into the cell. This transfection mechanism is referred to as "lipofection," and is a highly efficient transfection procedure (Felgner, *et al.*, *Proc. Natl. Acad. Sci. USA* 84:7413-7417, 1987); and Felgner, *et al.*, *Nature* 337:387-388, 1989). Still more preferably, the polynucleotide construct is complexed with one or more cationic lipids and one or more neutral lipids.

For purposes of the present invention, lipid refers to a synthetic or naturally occurring compound that possesses both a lipophilic region and a polar region, commonly referred to as a head group. Preferred cationic compounds are cationic lipids. Cationic lipids are described in U.S. Pat. Nos. 4,897,355; 4,946,787; 5,049,386; 5,264,618; 5,279,833; 5,334,761; 5,429,127; 5,459,127; 5,589,466; 5,676,954; 5,693,622; 5,580,859; 5,703,055; and 5,578,475; and international publications WO 04/9469, WO 95/14381, 95/14651, 95/17373, 96/18372, 96/26179, 96/40962, 96/40963, 96/41873, and

97/00241, and documents cited therein. As illustrated in the above-cited patents and patent applications, cationic lipids comprise structural features that may be present in a variety of core molecular classes.

Examples of cationic lipids are 5-carboxyspermylglycine dioctadecylamide (DOGS) and dipalmitoyl-phosphatidylethanolamine-5-carboxyspermylamide (DPPES). Cationic cholesterol derivatives are also useful, including {3 β -[N-N',N'-dimethylamino)ethane]-carbomoyl}-cholesterol (DC-Chol). Dimethyldioctadecyl-ammonium bromide (DDAB), N-(3-aminopropyl)-N,N-(bis-(2-tetradecyloxyethyl))-N-methyl-ammonium bromide (PA-DEMO), N-(3-aminopropyl)-N,N-(bis-(2-dodecyloxyethyl))-N-methyl-ammonium bromide (PA-DELO), N,N,N-tris-(2-dodecyloxy)ethyl-N-(3-amino)propyl-ammonium bromide (PA-TELO), and N¹-(3-aminopropyl)((2-dodecyloxy)ethyl)-N²-(2-dodecyloxy)ethyl-1-piperazinaminium bromide (GA-LOE-BP) can also be employed in the present invention.

Non-diether cationic lipids, such as DL-1,2-dioleoyl-3-dimethylaminopropyl- β -hydroxyethylammonium (DORI diester), 1-O-oleyl-2-oleoyl-3-dimethylaminopropyl- β -hydroxyethylammonium (DORI ester/ether), and their salts promote *in vivo* gene delivery. Preferred cationic lipids comprise groups attached via a heteroatom attached to the quaternary ammonium moiety in the head group. A glycol spacer can connect the linker to the hydroxyl group.

Preferred cationic lipids are 3,5-(N,N-dilysyl)-diaminobenzoyl-3-(DL-1,2-dioleoyl-dimethylaminopropyl- β -hydroxyethylamine) (DLYS-DABA-DORI diester), 3,5-(N,N-di-lysyl)amino-benzoylglycyl-3-(DL-1,2-dioleoyl-dimethylaminopropyl- β -hydroxyethylamine) (DLYS-DABA-GLY-DORI diester), and (\pm)-N-(2-hydroxyethyl)-N,N-dimethyl-2,3-bis(tetradecyloxy)-1-propaniminium bromide (DMRIE).

Also preferred are (\pm)-N,N-dimethyl-N-[2-(sperminecarboxamido)ethyl]-2,3-bis(dioleyloxy)-1-propaniminium pentahydrochloride (DOSPA), (\pm)-N-(2-aminoethyl)-N,N-dimethyl-2,3-bis(tetradecyloxy)-1-propaniminium bromide (β -aminoethyl-DMRIE or β AE-DMRIE) (Wheeler, *et al.*, *Biochim. Biophys. Acta* 1280:1-11 (1996)), and (\pm)-

N-(3-aminopropyl)-N,N-dimethyl-2,3-bis(dodecyloxy)-1-propaniminium bromide (GAP-DLRIE) (Wheeler, *et al.*, *Proc. Natl. Acad. Sci. USA* 93:11454-11459 (1996)), which have been developed from DMRIE.

Other examples of DMRIE-derived cationic lipids that are useful for the present invention are (\pm)-N-(3-aminopropyl)-N,N-dimethyl-2,3-(bis-decyloxy)-1-propanaminium bromide (GAP-DDRIE), (\pm)-N-(3-aminopropyl)-N,N-dimethyl-2,3-(bis-tetradecyloxy)-1-propanaminium bromide (GAP-DMRIE), (\pm)-N-((N"-methyl)-N'-ureyl)propyl-N,N-dimethyl-2,3-bis(tetradecyloxy)-1-propanaminium bromide (GMU-DMRIE), (\pm)-N-(2-hydroxyethyl)-N,N-dimethyl-2,3-bis(dodecyloxy)-1-propanaminium bromide (DLRIE), and (\pm)-N-(2-hydroxyethyl)-N,N-dimethyl-2,3-bis-([Z]-9-octadecenyloxy)propyl-1-propaniminium bromide (HP-DORIE).

The lipids of the lipid-containing formulation can comprise a cationic lipid alone, or further comprise a neutral lipid such as cardiolipin, phosphatidylcholine, phosphatidylethanolamine, dioleoylphosphatidylcholine, dioleoylphosphatidyl-ethanolamine, 1,2-dioleoyl-sn-glycero-3-phosphatidylethanolamine (DOPE), sphingomyelin, and mono-, di- or tri-acylglycerol. Other additives, such as cholesterol, fatty acid, ganglioside, glycolipid, neobee, niosome, prostaglandin, sphingolipid, and any other natural or synthetic amphiphiles, can also be used. A preferred molar ratio of cationic lipid to neutral lipid in these lipid-containing formulations is from about 9:1 to about 1:9; an equimolar ratio is particularly preferred. The lipid-containing formulation can further comprise a lyso lipid (*e.g.*, lyso-phosphatidylcholine, lysophosphatidyl-ethanolamine, or a lyso form of a cationic lipid).

More preferably, the cationic lipid is (\pm)-N-(2-hydroxyethyl)-N,N-dimethyl-2,3-bis(tetradecyloxy)-1-propaniminium bromide (DMRIE) and the neutral lipid is 1,2-dioleoyl-sn-glycero-3-phosphatidylethanolamine (DOPE) such that the mass ratio of polynucleotide construct to lipid is from about 10:1 and about 0.5:1. Still more preferably, the mass ratio of polynucleotide construct to lipid is from about 5:1 and about 1:1. Still more preferably, the mass ratio of polynucleotide construct to lipid is about 5:1.

Lipid-containing pharmaceutical composition for use in a complex with the polynucleotide construct of the present invention can also comprise cationic lipid together with an effective amount of a lysophosphatide. The lysophosphatide can have a neutral or a negative head group. Lysophosphatidylcholine and lysophosphatidyl-ethanolamine are preferred, and 1-oleoyl lysophosphatidylcholine is particularly preferred. Lysophosphatide lipids are advantageously present in the lipid-containing formulation in a 1:2 ratio of lysolipid to cationic lipid. Lyso forms of a cationic lipid can also be used to promote polynucleotide delivery. These lyso forms are advantageously present in effective amounts up to about one-third of the total cationic lipid in the lipid-containing formulations.

In a formulation for preparing DNA: lipid complexes, the cationic lipid can be present at a concentration of between about 0.1 mole % and about 100 mole %, preferably about 5 mole % and 100 mole %, and most preferably between about 20 mole % and 100 mole %, relative to other formulation components present in the formulation. The neutral lipid can be present in a concentration of between zero and about 99.9 mole %, preferably zero and about 95 mole %, and most preferably zero and about 80 mole %. In order to produce lipid vesicles having a net positive charge, the quantity of the positively charged component must exceed that of the negatively charged component. The negatively charged lipid can be present at between zero and about 49 mole %, and preferably between zero and about 40 mole %. Cholesterol or a similar sterol can be present at between zero to about 80 mole %, and preferably zero and about 50 mole %.

The polynucleotide to be delivered can be solubilized in a buffer prior to mixing with lipid vesicles. Suitable buffers include, for example, phosphate buffered saline (PBS), normal saline, Tris buffer, and sodium phosphate vehicle (100-150 mM preferred). Insoluble polynucleotides can be solubilized in a weak acid or base, and then diluted to the desired volume with a neutral buffer such as PBS. The pH of the buffer is suitably adjusted, and moreover, a pharmaceutically acceptable additive can be used in the buffer to provide an appropriate osmolarity within the lipid vesicle.

A lipid solution comprising at least one amphipathic lipid can spontaneously assemble to form primary lipid vesicles, heterogeneous in size. Therefore, according to a preferred method, the lipids of the lipid-containing formulation, comprising at least one cationic lipid, are prepared by dissolution in a solvent such as chloroform and the mixture is evaporated to dryness as a film on the inner surface of a glass vessel. On suspension in an aqueous solvent, the amphipathic lipid molecules assemble themselves into primary lipid vesicles. These primary lipid vesicles may be reduced to a selected mean diameter by means of a freeze-thaw procedure. Vesicles of uniform size can be formed prior to drug delivery according to methods for vesicle production known to those in the art; for example, the sonication of a lipid solution as described by Felgner, *et al* (*Proc. Natl. Acad. Sci. USA* 84:7413-7417 (1987)) and U.S. Pat. No. 5,264,618.

The term "mammal" is intended to encompass a singular "mammal" and plural "mammals," and includes, but is not limited to humans; primate mammals such as apes, monkeys, orangutans, and chimpanzees; canine mammals such as dogs and wolves; feline mammals such as cats, lions, and tigers; equine mammals such as horses, donkeys, deer, zebra, and giraffe; and bears. Preferably, the mammal is a human subject.

Tumor cell formation and growth, also known as "transformation," describes the formation and proliferation of cells that have lost their ability to control cellular division; that is the cells are cancerous. "Malignant cells" are defined as cells that have lost the ability to control the cell division cycle, leading to a transformed or cancerous phenotype.

The term "non-tumor tissue" is intended to include, but is not limited to non-tumor bearing tissues such as muscle, skin, brain, lung, liver, spleen, bone marrow, thymus, heart, lymph, blood, bone, cartilage, pancreas, kidney, gall bladder, stomach, intestine, testis, ovary, uterus, rectum, nervous system, eye, gland, or connective tissue. Preferably, the non-tumor tissue is muscle.

Preferably, the polynucleotide construct is delivered to the interstitial space of a tumor or of non-tumor tissues. "Interstitial space" comprises the intercellular, fluid, mucopolysaccharide matrix among the reticular fibers of

organ tissues, elastic fibers in the walls of vessels or chambers, collagen fibers of fibrous tissues, or that same matrix within connective tissue ensheathing muscle cells or in the lacunae of bone. It is similarly the space occupied by the plasma of the circulation and the lymph fluid of the lymphatic channels.

5 The pharmaceutical composition and methods of the present invention can be used to treat a variety of mammalian cancers or tumors. Types of mammalian cancers and tumors that can be treated using the pharmaceutical composition and methods of the present invention include, but are not limited to all solid tumors, cutaneous tumors, melanoma, malignant melanoma, renal
10 cell carcinoma, colorectal carcinoma, colon cancer, hepatic metastases of advanced colorectal carcinoma, lymphomas (including glandular lymphoma), malignant lymphoma, Kaposi's sarcoma, prostate cancer, kidney cancer, ovarian cancer, lung cancer, head and neck cancer, pancreatic cancer, mesenteric cancer, gastric cancer, rectal cancer, stomach cancer, bladder
15 cancer, leukemia (including hairy cell leukemia and chronic myelogenous leukemia), breast cancer, non-melanoma skin cancer (including squamous cell carcinoma and basal cell carcinoma), hemangioma multiple myeloma, and glioma. Preferably, the cancer is melanoma, ovarian cancer, or metastases thereof.

20 By "treatment" is meant reduction in tumor size, a reduction in the rate of metastasis, and/or a slowing of tumor growth, and/or no worsening in disease over a specified period of time.

 A systemic delivery embodiment can be particularly useful for treating nonlocalized tumors (*i.e.*, leukemia and metastases of a variety of tumors), or a
25 disease category that might be responsive to continuous exposure by the systemic route (*i.e.*, myeloma, chronic myelogenous leukemia, lymphoma). A local delivery embodiment can be particularly useful for treating one disease condition that might be responsive to high local concentration (*i.e.*, renal cell carcinoma, melanoma). For tumors involving body cavity of a mammal,
30 "intra-cavity" embodiment is preferred. In particular, the use of this methodology is envisioned in treating cancers involving (i) the peritoneal cavity--pancreatic cancer, gastric cancer, ovarian cancer, mesenteric cancer,

glandular lymphoma and metastatic melanoma; (ii) the thoracic cavity--lung cancer and glandular lymphoma; (iii) the rectal cavity--rectal cancer; (iv) the stomach cavity--stomach cancer; and (v) the urinary bladder vesicle--bladder cancer. When advantageous, systemic, local, and/or intra-cavity delivery can be combined, especially in a mammal having a primary site of tumor and one or more metastases.

An additional embodiment of the present invention is directed to combining any of the methods of the present invention with one or more additional cancer therapies including, but not limited to bone marrow transplant, cord blood cell transplant, surgery, chemotherapy, radiation therapy, and immunotherapy. The polynucleotide construct or pharmaceutical composition of the present invention can be administered prior to the commencement of one or more of the additional cancer therapies, during the practice of one or more of the additional cancer therapies, and after the end of one or more of the additional cancer therapies.

Types of bone marrow transplant include, but are not limited to autologous bone marrow transplant and heterologous (*i.e.*, from a donor) bone marrow transplant.

Types of surgery include, but are not limited to surgery for breast cancer, prostate cancer, colon cancer, brain cancer, and head and neck cancer.

Chemotherapeutic agents include, but are not limited to alkylating agents, including mechlorethamine, cyclophosphamide, ifosfamide, melphalan, chlorambucil, dicarbazine, streptazocine, carmustine, lomustine, semustine, chlorozotocin, busulfan, triethylenemelamine, thiotepa, hexamethylmelamine; antimetabolites, including methotrexate; pyrimidine analogs, including fluorouracil, 5-fluorouracil, floxuridine (5'-fluoro-2'-deoxyuridine), idoxuridine, cytarabine, -phosphonoacetyl-L-aspartate, 5-azacytidine, azaribine, 6-azauridine, pyrazofuran, 3-deazauridine, acivicin; purine analogs, including thioguanine, mercaptopurine, azathioprine, pentostatin, erythrohydroxynonyladenine; vinca alkaloids, including vincristine and vinblastine, epipodophyllotoxins, including etoposide and teniposide; antibiotics, including dactinomycin, daunorubicin, doxorubicin,

bleomycin sulfate, plicamycin, mitomycin; enzymes, including L-asparaginase; platinum coordination complexes, including cisplatin, carboplatin; hydroxyurea, procarbazine, mitotane; and hormones or related agents, including adrenocorticosteroids such as prednisone and prednisolone; aminogluthethimide; progestins such as hydroxyprogesterone caproate, medroxyprogesterone acetate, megestrol acetate, estrogens and androgens such as diethylstilbestrol, fluoxymesterone, ethynyl estradiol, antiestrogens such as tamoxifen, and gonadotropin-releasing hormone analogs such as leuprolide.

The present invention also provides kits for use in treating cancer comprising an administration means and a container means containing one or more cytokine-expressing polynucleotide constructs in a sterile environment. Also provided are kits for use in treating cancer comprising an administration means and a container means containing one or more cytokine-expressing polynucleotide constructs and one or more cationic compounds in a sterile environment. Examples of cationic compounds are described above. The cytokine-expressing polynucleotide constructs and the cationic compounds may be in the same container means or in separate container means. Preferably, the polynucleotide construct is in the amount of 1 ng to 20 mg.

Container means include glass containers, plastic containers, or strips of plastic or paper. In one embodiment, the container means is a syringe and the administration means is a plunger. In another embodiment, the administration means is a catheter.

The cytokine encoded by the polynucleotide construct of the kit of the present invention can be an IFN ω and one or more additional cytokines, including any of the cytokines described herein. Preferably, the cytokine is IFN ω and/or an IFN α . The construct can be in the form of a pharmaceutical composition and can contain a pharmaceutically acceptable carrier. Pharmaceutical compositions are described above. The kit can further comprise a pharmaceutically acceptable carrier in a separate container means.

The kit can further comprise an instruction sheet for administration of the composition into a mammal. The components of the polynucleotide

composition are preferably provided as a liquid solution, such as a suspension, a solution, or an emulsion; or in lyophilized form as a dried powder or a cake. If the polynucleotide construct is provided in lyophilized form, preferably the kit further comprises a container means containing a suitable vehicle, such as sterile pyrogen-free water, for reconstitution of the lyophilized polynucleotide construct, or any buffer described herein, including PBS, normal saline, Tris buffer, and sodium phosphate vehicle.

The term "cytokine" refers to polypeptides, including but not limited to interleukins (*e.g.*, IL-1, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-11, IL-12, IL-13, IL-14, IL-15, IL-16, IL-17, and IL-18), α interferons (*e.g.*, IFN α), β interferons (*e.g.*, IFN β), γ interferons (*e.g.*, IFN γ), ω interferon (IFN ω), τ interferons (IFN τ), colony stimulating factors (CSFs, *e.g.*, CSF-1, CSF-2, and CSF-3), granulocyte-macrophage colony stimulating factor (GM-CSF), epidermal growth factor (EGF), fibroblast growth factors (FGFs, *e.g.*, acidic fibroblast growth factor, basic fibroblast growth factor, FGF-1, FGF-2, FGF-3, FGF-4, and FGF-5), transforming growth factor (TGF, *e.g.*, TGF α and TGF β), platelet-derived growth factor (PDGF), tumor necrosis factors (TNFs, *e.g.*, TNF- α and TNF- β), and insulin-like growth factors (IGFs, *e.g.*, IGF-I and IGF-II).

A "polypeptide" refers to any translation product of a polynucleotide, regardless of the size of the translation product, and regardless of whether the translation product is post-translationally modified (*e.g.*, glycosylated) or not.

The polynucleotide construct of the present invention, whether complexed with cationic vehicle or not, can be administered by any suitable route of administration, including intramuscularly, subcutaneously, intravenously, transdermally, intranasally, by inhalation, or transmucosally (*i.e.*, across a mucous membrane). Similarly, the pharmaceutical composition of the present invention can be administered by any suitable route, including intramuscularly, into or near a tumor, into a cavity (*e.g.*, intraperitoneally), subcutaneously, intravenously, transdermally, intranasally, by inhalation, or transmucosally (*i.e.*, across a mucous membrane).

Any mode of administration can be used so long as the mode results in the expression of one or more cytokines in an amount sufficient to decrease the tumorigenicity of the cancer bearing mammal. This includes needle injection, catheter infusion, biolistic injectors, particle accelerators (*i.e.*, "gene guns"), pneumatic "needleless" injectors (*e.g.*, MedEJet, PedeJet, Bioject), gelfoam sponge depots, other commercially available depot materials, osmotic pumps (*e.g.*, Alza minipumps), oral or suppositorial solid (tablet or pill) pharmaceutical formulations, and decanting or topical applications during surgery. Preferred methods include needle injection and catheter infusion.

A "polynucleotide construct" is a polynucleotide molecule that carries genetic information for encoding one or more molecules, preferably, cytokines. The polynucleotide material delivered to the cells *in vivo* can take any number of forms. It can contain the entire sequence or only a functionally active fragment of a cytokine gene.

The polynucleotide construct comprises at least one polynucleotide (*e.g.*, DNA, RNA, ribozyme, phosphorothioate, or other modified nucleic acid) encoding one or more molecules. Preferred molecules are cytokines. The polynucleotide can be provided in linear, circular (*e.g.* plasmid), or branched form; and double-stranded or single-stranded form. The polynucleotide can involve a conventional phosphodiester bond or a non-conventional bond (*e.g.*, an amide bond as in peptide nucleic acid (PNA)). The choice of polynucleotide encoding a cytokine will depend on the desired kinetics and duration of expression. When long term delivery of the polynucleotide construct is desired, the preferred polynucleotide is DNA. Alternatively, when short term delivery is desired, the preferred polynucleotide is mRNA. RNA will be rapidly translated into polypeptide, but will be degraded by the target cell more quickly than DNA. In general, because of the greater resistance of circular DNA molecules to nucleases, circular DNA molecules will persist longer than linear polynucleotides, and they will be less likely to cause insertional mutation by integrating into the target genome.

In one embodiment, the polynucleotide sequence encoding one or more cytokines is RNA. Most preferably, the RNA is messenger RNA (mRNA). Methods for introducing RNA sequences into mammalian cells is described in U.S. patent No. 5,580,859. A viral alphavector, a non-infectious vector useful for administering RNA, may be used to introduce RNA into mammalian cells. Methods for the *in vivo* introduction of alphaviral vectors to mammalian tissues are described in Altman-Hamamdzie, S., *et al.*, *Gene Therapy* 4: 815-822 (1997). Preferably, the polynucleotide sequence encoding one or more cytokines is DNA. In a DNA construct, a promoter is preferably operably linked to the polynucleotide encoding a cytokine. The promoter may be a cell-specific promoter that directs substantial transcription of the DNA only in predetermined cells. Other transcription control elements, besides a promoter, can be included in the polynucleotide construct to direct cell-specific transcription of the DNA.

An operable linkage is a linkage in which a polynucleotide sequence encoding a cytokine is connected to one or more regulatory sequence in such a way as to place expression of the cytokine sequence under the influence or control of the regulatory sequence(s). Two DNA sequences (such as a coding sequence and a promoter region sequence linked to the 5' end of the coding sequence) are operably linked if induction of promoter function results in the transcription of mRNA encoding the desired polypeptide and if the nature of the linkage between the two DNA sequences does not (1) result in the introduction of a frame-shift mutation, (2) interfere with the ability of the expression regulatory sequences to direct the expression of the polypeptide, antisense RNA, or (3) interfere with the ability of the DNA template to be transcribed. Thus, a promoter region would be operably linked to a DNA sequence if the promoter was capable of affecting transcription of that DNA sequence.

Preferably, the polynucleotide construct is a circular or linearized plasmid containing non-infectious nucleotide sequence. A linearized plasmid is a plasmid that was previously circular but has been linearized, for example, by digestion with a restriction endonuclease. The polynucleotide sequence

encoding a cytokine may comprise a sequence which directs the secretion of the polypeptide.

"Non-infectious" means that the polynucleotide construct does not infect mammalian cells. Thus, the polynucleotide construct can contain functional sequences from non-mammalian (*e.g.*, viral or bacterial) species, but the construct does not contain non-mammalian nucleotide sequences that facilitate infection of the construct into mammalian cells.

"Non-integrating" means that the polynucleotide construct does not integrate into the genome of mammalian cells. The construct can be a non-replicating DNA sequence, or specific replicating sequences genetically engineered to lack the ability to integrate into the genome. The polynucleotide construct does not contain functional sequences that facilitate integration of the cytokine-encoding polynucleotide sequence into the genome of mammalian cells.

The polynucleotide construct is assembled out of components where different selectable genes, origins, promoters, introns, 5' untranslated (UT) sequence, terminators, polyadenylation signals, 3' UT sequence, and leader peptides, etc. are put together to make the desired vector. The precise nature of the regulatory regions needed for gene expression can vary between species or cell types, but shall in general include, as necessary, 5' non-transcribing and 5' non-translating (non-coding) sequences involved with initiation of transcription and translation respectively, such as the TATA box, capping sequence, CAAT sequence, and the like, with those elements necessary for the promoter sequence being provided by the promoters of the invention. Such transcriptional control sequences can also include enhancer sequences or upstream activator sequences, as desired.

The polynucleotide construct can be an expression vector. A typical mammalian expression vector contains the promoter element, which mediates the initiation of transcription of mRNA, the polypeptide coding sequence, and signals required for the termination of transcription and polyadenylation of the transcript. Additional elements include enhancers, Kozak sequences and intervening sequences flanked by donor and acceptor sites for RNA splicing.

Highly efficient transcription can be achieved with the early and late promoters from SV40, the long terminal repeats (LTRS) from retroviruses, e.g., RSV, HTLV, HIV, MPSV and the immediate early promoter of the cytomegalovirus (CMV IEP). However, cellular elements can also be used (e.g., the human actin promoter, metallothionein promoter). In humans, CMV IEP is preferred. Suitable expression vectors for use in practicing the present invention include, for example, vectors such as PSVL and PMSG (Pharmacia, Uppsala, Sweden), pRSVcat (ATCC 37152), pSV2dhfr (ATCC 37146) and pBC12MI (ATCC 67109), VR1012, VR1055, and pcDNA3 (Invitrogen, San Diego, CA). All forms of DNA, whether replicating or non-replicating, which do not become integrated into the genome, and which are expressible, are within the methods contemplated by the invention.

The vector containing the DNA sequence (or the corresponding RNA sequence) which can be used in accordance with the invention can be a eukaryotic expression vector. Techniques for obtaining expression of exogenous DNA or RNA sequences in a host are known. See, for example, Korman, *et al.*, *Proc. Nat. Acad. Sci. (USA)* 84:2150-2154 (1987).

Secretion of a cytokine from a cell can be facilitated by a leader or secretory signal sequence. In a preferred embodiment, either the native leader sequence of a cytokine is used, or a functional derivative of that sequence that retains the ability to direct the secretion of the peptide that is operably linked to it. Alternatively, a heterologous mammalian leader sequence, or a functional derivative thereof, may be used. For example, the wild-type leader sequence may be substituted with the leader sequence of human tissue plasminogen activator or mouse β -glucuronidase.

For the methods of the present invention, a single polynucleotide construct containing more than one polynucleotide sequences encoding one or more molecules, or more than one polynucleotide constructs each containing polynucleotide sequences encoding one or more molecules may be co-injected or sequentially injected. For example, a single polynucleotide construct containing one polynucleotide encoding an interferon and another polynucleotide encoding an additional cytokine or an immunomodulatory

molecule, *i.e.*, MHC class I antigen, tumor antigen, and co-stimulatory molecule, can be injected. Alternatively, two polynucleotide construct can be injected where one encodes a cytokine to enhance anti-tumor efficacy of the other gene product. For example, an IFN γ , IFN α , IL-12 or IL-2-expressing polynucleotide construct can be co-injected with a polynucleotide construct encoding a different cytokine. More specifically, an IL-2 expressing plasmid could be co-injected with a G-CSF or GM-CSF expressing plasmid. Alternatively, one or more plasmids could be administered initially and other plasmid(s) could be administered subsequently at various time intervals. Combination of the present invention with therapeutic agents such as lymphokine-activated killer cells (LAK) and tumor-infiltrating lymphocytes (TIL) is also envisioned.

It will be recognized in the art that some amino acid sequences of the polypeptides described herein can be varied without significant effect on the functional activity of the polypeptides. If such differences in sequence are contemplated, it should be remembered that there will be critical areas on the polypeptide which determine activity. Such variations include deletions, insertions, inversions, repeats, and type substitutions. Guidance concerning which amino acid changes are likely to be phenotypically silent can be found in Bowie, J.U., *et al.*, "Deciphering the Message in Protein Sequences: Tolerance to Amino Acid Substitutions," (*Science* 247:1306-1310 (1990)). Compositions within the scope of the invention can be assayed according to the antiproliferation assay described herein. Amino acids that are critical for cytokine activity can also be determined by structural analysis such as crystallization, nuclear magnetic resonance or photoaffinity labeling (Smith, *et al.*, *J. Mol. Biol.* 224:899-904 (1992) and de Vos, *et al.* *Science* 255:306-312 (1992)).

The present invention further relates to using variants of the cytokine-encoding polynucleotide, which encode portions, analogs or derivatives of the cytokine. Variants may occur naturally, such as a natural allelic variant. By an "allelic variant" is intended one of several alternate forms of a gene occupying a given locus on a chromosome of an organism. *Genes II*, Lewin, B., ed., John

Wiley & Sons, New York (1985). Non-naturally occurring variants may be produced using art-known mutagenesis techniques.

Alterations in the coding regions may produce conservative or non-conservative amino acid substitutions, deletions or additions. Especially preferred among these are silent substitutions, additions and deletions, which do not alter the properties and activities of the cytokine or portions thereof. Also especially preferred in this regard are conservative substitutions. For example, aromatic amino acids that can be conservatively substituted for one another include phenylalanine, tryptophan, and tyrosine. Hydrophobic amino acids that can be conservatively substituted for one another include leucine, isoleucine, and valine. Polar amino acids that can be conservatively substituted for one another include glutamine and asparagine. Basic amino acids that can be conservatively substituted for one another include arginine, lysine, and histidine. Acidic amino acids that can be conservatively substituted for one another include aspartic acid and glutamic acid. Small amino acids that can be conservatively substituted for one another include alanine, serine, threonine, methionine, and glycine.

Substitutions, deletions, or insertions can be made outside of the region encoding the shortest active fragment of the cytokine, without affecting the activity of the cytokine. Further, mutated proteins (or muteins) often retain a biological activity that is similar to that of the naturally occurring protein. For example, Gayle and coworkers (*J. Biol. Chem.* 268: 22105-22111 (1993)) conducted an extensive mutational analysis of the human cytokine IL-1 α . They used random mutagenesis to generate over 3,500 individual IL-1 α mutants with an average of 2.5 amino acid changes per mutein over the entire length of the molecule. Multiple mutations were examined at every possible amino acid and, on average, each mutein's amino acid sequence was 98.4% identical to that of naturally occurring IL-1 α . The investigators observed that most of the molecule could be mutated with little effect on either binding or biological activity, and that 75% of the molecule may not contribute significantly to the biological activity of the molecule.

Similarly, Gronenborn and colleagues (*FEBS Letters* 231: 135-138 (1988)) analyzed the receptor binding activity of six mutant IL-1 α polypeptides. Each mutant contained a single amino acid alteration from the naturally occurring IL-1 α polypeptide and was examined under four sets of experimental conditions. In this study, the investigators found very little difference between the receptor binding activity of the mutants and naturally occurring IL-1 α .

Further, Zurawski and colleagues (*EMBO J.* 12: 5113-5119 (1993)) studied residues 41-142 of mIL-2 by generating 1,090 muteins. The extent of the mutagenesis was such that there was an average of 11 different amino acid substitutions per naturally occurring amino acid residue, with the exception of the extreme N- and C-termini and residues 31-40. The mIL-2 muteins were assayed for specific activity and compared to that of naturally occurring mIL-2. The degree to which the specific activity was antagonized by a previously characterized mIL-2 mutant was also assessed. The investigators observed that in the 149 residue mIL-2 protein, only 23 residues are important for interaction with IL-2R, 18 residues are presumed to be part of the structural core, and three additional residues are important for structure. 98 mIL-2 residues (or 65% of the protein) were assigned as relatively unimportant residues.

The polynucleotide and amino acid sequences encoding an IFN ω include the sequences for the complete IFN ω and the mature IFN ω set forth in U.S. Patent No. 4,917,887; European Patent Publication No. 0 170 204 B1; and Capon, D.J., *et al.*, *Molec. Cell. Biol.* 5: 768-779 (1985); Hauptmann, R. and P. Swetly, *Nucl. Acids Res.* 13: 4739-4749 (1985); Adolf, G.R., *et al.*, *Biochim. Biophys. Acta* 1089: 167-174 (1991); Mege, D., *et al.*, *J. Interf. Res.* 11: 341-350 (1991); Charlier, M., *et al.*, *J. Interf. Res.* 13: 313-322 (1993); Hughes, A.L., *J. Mol. Evol.* 41: 539-548 (1995); and Roberts, R.M., *et al.*, *Prog. Nucl. Acid Res. Molec. Biol.* 56:287-325, edited by W.E. Cohn, Academic Press (1997).

The polynucleotide and amino acid sequences encoding IFN α include the sequences for the complete IFN α and the mature IFN α set forth in U.S. Patent Nos. 4,530,901; 4,695,543; 4,695,623; 4,748,233; 4,892,743; 4,897,471; 4,973,479; 4,975,276; and 5,098,703; and in Pestka, S., *Methods Enzymol.* 119: 3-14 (1986); Hughes, A.L., *J. Mol. Evol.* 41: 539-548 (1995); and Roberts, R.M., *et al.*, *Prog. Nucl. Acid Res. Molec. Biol.* 56:287-325, edited by W.E. Cohn, Academic Press (1997).

The polynucleotide and amino acid sequences encoding an IL-2 include the sequences for the complete human IL-2 and mature IL-2 set forth in Lupker, J. *et al.*, EP 0307285-A3 (1989), U.S. Patent No. 5,641,665, Maeda *et al.*, *Biochem. Biophys. Res. Commun.* 115:1040-1047 (1983), Mita *et al.*, *Biochem. Biophys. Res. Commun.* 117:114-121 (1983), Taniguchi *et al.*, *Nature* 302:305-310 (1983), Devos *et al.*, *Nucleic Acid Res.* 11:4307-4323 (1983), Fujita *et al.*, *Proc. Natl. Acad. Sci. USA* 80:74347-7441 (1983), Clark *et al.*, *Proc. Natl. Acad. Sci. USA* 81:2543-2547 (1984) and Cullen, *DNA* 7:645-650 (1988).

The polynucleotide sequences encoding an IFN ω , an IFN α , and an IL-2 also include sequences that encode the complete polypeptide encoded by the nucleotide sequences set forth in SEQ ID Nos. 7, 9 and 13, respectively, and the mature polypeptides encoded by nucleotide sequences set forth in SEQ ID Nos. 7, 9 and 13, respectively. The polynucleotide sequences encoding IL-2 further includes the sequence that encodes the complete IL-2 polypeptide encoded by the nucleotide sequence set forth in SEQ ID No. 25, shown as SEQ ID No. 26.

Thus, a polynucleotide sequence encoding a polypeptide of the present invention can encode a polypeptide having one to twenty amino acid substitutions, deletions or insertions, either from natural mutations or human manipulation, relative to the full length or mature IFN α , IFN ω , or IL-2. Preferably, no more than one to fifteen substitutions, deletions or insertions are present, relative to the full length or mature IFN α , IFN ω , or IL-2 (excluding the signal sequence). More preferably, no more than one to ten

substitutions, deletions or insertions are present. Still more preferably, no more than one to five substitutions, deletions or insertions are present.

Determining an effective amount of substance to be delivered can depend upon a number of factors including, for example, the chemical structure and biological activity of the substance, the age and weight of the mammal, the precise condition requiring treatment and its severity, and the route of administration. The precise amount, number of doses, and timing of doses will be determined by the attending physician or veterinarian.

If the polynucleotide construct of the present invention is administered as a pharmaceutical composition, the pharmaceutical composition can be formulated according to known methods for preparing pharmaceutical compositions, whereby the substance to be delivered is combined with a pharmaceutically acceptable carrier vehicle. Suitable vehicles and their preparation are described, for example, in Remington's Pharmaceutical Sciences, 16th Edition, A. Osol, Ed., Mack Publishing Co., Easton, PA (1980), and Remington's Pharmaceutical Sciences, 19th Edition, A.R. Gennaro, Ed., Mack Publishing Co., Easton, PA (1995).

The pharmaceutical composition can be in the form of an emulsion, gel, solution, suspension, or other form known in the art. Optionally, it can contain one or more lipids as described above. In addition, the pharmaceutical composition can also contain pharmaceutically acceptable additives including, for example, diluents, binders, stabilizers, and preservatives. Administration of pharmaceutically acceptable salts of the polynucleotides described herein is preferred. Such salts can be prepared from pharmaceutically acceptable non-toxic bases including organic bases and inorganic bases. Salts derived from inorganic bases include sodium, potassium, lithium, ammonium, calcium, magnesium, and the like. Salts derived from pharmaceutically acceptable organic non-toxic bases include salts of primary, secondary, and tertiary amines, basic amino acids, and the like.

For aqueous pharmaceutical compositions used *in vivo*, sterile pyrogen-free water is preferred. Such formulations will contain an effective amount of the substance together with a suitable amount of vehicle in order to

prepare pharmaceutically acceptable compositions suitable for administration to a human or animal.

5 A pharmaceutical composition can be in solution form, or alternatively, in lyophilized form for reconstitution with a suitable vehicle, such as sterile, pyrogen-free water. Both liquid and lyophilized forms will comprise one or more agents, preferably buffers, in amounts necessary to suitably adjust the pH of the injected solution.

10 The container in which the pharmaceutical formulation is packaged prior to use can comprise a hermetically sealed container enclosing an amount of the lyophilized formulation or a solution containing the formulation suitable for a pharmaceutically effective dose thereof, or multiples of an effective dose. The pharmaceutical formulation is packaged in a sterile container, and the hermetically sealed container is designed to preserve sterility of the pharmaceutical formulation until use. Optionally, the container can be
15 associated with administration means and or instruction for use.

Having now generally described the invention, the same will become more readily understood by reference to the following specific examples which are included herein for purposes of illustration only and are not intended to be limiting unless otherwise specified.

20 Described herein are: 1) the *in vitro* characterization of biological activities of and IFNs delivered by plasmid DNA (anti-proliferative activity and anti-viral activity *in vitro*); 2) *in vivo* expression of cytokines following *in vivo* administration of cytokine-expressing pDNA; and 3) the *in vivo* characterization of anti-tumor activity of cytokines in murine models of solid
25 and metastatic tumors following intratumoral, intramuscular or intra-cavity administration of cytokine-encoding pDNA.

The cytokine-encoding polynucleotide constructs have potent anti-proliferative activity *in vitro*. Moreover, the *in vivo* anti-tumor activities of IFN ω , IFN α , IL-2, and IL-12 are herein demonstrated in multiple murine
30 tumor models including nude mice bearing subcutaneous human tumors, or in immunocompetent mice bearing murine solid and metastatic tumors. Intratumoral, intramuscular, or intraperitoneal injection of the cytokine-

encoding plasmids is shown to result in a statistically significant slowing of tumor growth and/or a statistically significant increase in survival. In addition to the potent antitumor effects of the cytokine plasmids delivered via intratumoral or intramuscular injection, this is the first *in vivo* demonstration of anti-tumor activity for human interferon- ω . Moreover, the *in vivo* antitumor activity of IL-2 in the treatment of peritoneally disseminated cancers, such as ovarian metastatic melanoma is also demonstrated.

Example 1

Construction of Expression Vectors

Three basic eukaryotic expression plasmid vectors, termed VR1012, VR1055 and VR1033 were used in the construction of all plasmids used in the following examples. The blank plasmids, VR1012 and VR1055 differ only in transcriptional termination sequences. The backbone of both plasmids is derived from pUC19, with the beta-lactamase (ampicillin resistance) gene replaced by the aminoglycoside acetyltransferase (kanamycin resistance) gene from pET9a (Novagen, Madison, WI). Both plasmids direct eukaryotic gene expression from a cassette containing the human cytomegalovirus immediate early I (CMV IE) gene promoter/enhancer, CMV IE 5' untranslated (UT) sequence, and intron A. Following these regulatory elements is a cloning polylinker for insertion of polypeptide coding sequences. Following the polylinker in VR1012 is the 3' UT sequence from the bovine growth hormone gene for polyadenylation and transcriptional termination. In VR1055, the transcriptional terminator region includes a polyadenylation and termination signals derived from the rabbit b-globin gene. VR1033 is identical to VR1012, except that it contains a cap-independent translational enhancer from the encephalomyocarditis virus within the cloning polylinker sequence. This sequence allows the production of two different polypeptides from a single expressed mRNA.

Plasmid VR4101 (murine interferon α (mIFN α)) was constructed by cloning the murine interferon α cDNA into the vector VR1012 vector. The

cDNA was obtained by amplifying the coding sequence from the plasmid RSV-"1 (Kelly, K.A. and P.M. Pitha, *Nucl. Acids Res.* 13: 805-823 (1985); Kelly, K.A. and P.M. Pitha, *Nucl. Acids Res.* 13: 825-839 (1985)), which was provided by Dr. Paula Pitha-Rowe of Johns Hopkins University. Plasmid VR4111 was constructed by transferring the coding sequences from VR4101 to the VR1055 cloning vector. The oligonucleotide primers used for polymerase chain reaction (PCR) were 5'-AACTGCAGATGGCTAGGCTCTGTGCT-3' (SEQ ID No. 15) and 5'-GAAG-ATCTTCATTTCTTCTC-TCAG-3' (SEQ ID No. 16). Reaction conditions were 30 cycles of 94°C for 1 minute (denaturing), 58°C for 2 minutes (annealing), and 72°C for 1 minute (extension).

Plasmid VR4102 (human interferon α (hIFN α)) was constructed by cloning the human interferon α cDNA into the VR1012 vector. The cDNA was obtained by amplifying the coding sequence from human genomic DNA prepared from a fresh blood sample. Plasmid VR4112 was constructed by transferring the coding sequence sequences from VR4102 to the VR1055 cloning vector. Genomic DNA was isolated using the QIAamp Blood Kit (Qiagen, Inc.). The oligonucleotide primers used for PCR were 5'-AACTGCAGATGGCCTC-GCCCTTTGCT-3' (SEQ ID No. 17) and 5'-CGGGATCCTTATTCCTTC-CTCCTTAATC-3' (SEQ ID No. 18). Reaction conditions were 30 cycles of 94°C for 1 minute (denaturing), 58°C for 2 minutes (annealing), and 72°C for 1 minute (extension).

Plasmid VR4150 (human interferon ω (hIFN ω)) was constructed by cloning the human IFN ω cDNA into the VR1012 cloning vector. The cDNA was obtained by amplifying the coding sequence from human genomic DNA prepared from a fresh blood sample. Plasmid VR4151 (SEQ ID No. 1) was constructed by transferring the coding sequences from VR4150 to the VR1055 cloning vector. The oligonucleotide primers used for PCR were 5'-GCTCTAGATGGCCCTCCTGTTCCCT-3' (SEQ ID No. 19) and 5'-GCGG-ATCCTCAAGATGAGCCCAGGTC-3' (SEQ ID No. 20). Reaction conditions were 30 cycles of 94°C for 1 minute (denaturing), 58°C for 2 minutes (annealing), and 72°C for 1 minute (extension).

Plasmid VR1110 (murine interleukin-2 (mIL-2)) was constructed by cloning modified murine IL-2 cDNA into the VR1012 vector. The 5' UT sequence and the two amino acids of the leader peptide were replaced with the rat insulin II gene 5' UT sequence and coding region of the first six amino acids of the rat preproinsulin leader peptide. The IL-2 cDNA was then cloned into the BamHI site of VR1012.

Plasmid VR1103 (human interleukin-2 (hIL-2)) is identical to VR1110 with the exception that the murine IL-2 cDNA was replaced with the cDNA for human IL-2 (Parker et. al. 1996).

Plasmid VR4001 (murine interleukin-12 (mIL-12)) was constructed by cloning the cDNA's encoding the two murine subunits p35 and p40 into the VR1033 vector. Both cDNA's were obtained by amplifying the coding sequences from plasmids provided by Dr. Thomas Gajewski of The University of Chicago (J. Immun., 154:5637; J. Immun., 156:1095). The oligonucleotides used for PCR of p35 were 5'- CAT GCC ATG GGT CAA TCA CGC TAC CTC CTC TTT TTG G-3' (SEQ ID No. 23) and 5'- GCG GAT CCT CAG GCG GAG CTC AGA TAG CCC-3' (SEQ ID No. 24). The oligonucleotides used for PCR of p40 were 5'- ACG CGT CGA CAT GTG TCC TCA GAA GCT AAC CAT CTC-3' (SEQ ID No. 21) and 5'- GCG GAT CCC TAG GAT CGG ACC CTG CAG GGA ACA C-3' (SEQ ID No. 22). Reaction conditions were 30 cycles of 94°C for 1 minute (denaturing), 58°C for 2 minutes (annealing), and 72°C for 1 minute (extension).

Plasmids VR1223 (luciferase) was constructed by cloning cytoplasmic luciferase gene into the VR1012 vector (Hartikka *et al.*, *Hum. Gene Ther.* 7:1205-1217, 1996). The source of the cytoplasmic luciferase gene found in VR1223 was the plasmid termed pSP-luc+ which was purchased from Promega. An Avr II - Xba I restriction fragment encoding the luciferase cDNA was transferred from pSP-luc+ to VR1012 to make VR1223.

Plasmid VR1412 (β -galactosidase) was constructed by cloning cytoplasmic β -gal gene into the VR1012 vector (Doh *et al.*, *Gene Ther.* 4:648-663).

-50-

Plasmid VR1332 was constructed by inserting a *SalI-BamHI* fragment encoding chloramphenicol acetyltransferase (CAT) from pBS-CAT (Promega) into *SalI/BamHI*-cut VR1012 vector (Hartikka *et al.*, *Hum. Gene Ther.* 7:1205-1217 (1996)).

5

Example 2

Purification of pDNA

10

pDNA was transformed into *Escherichia coli* DH10B-competent cells and grown in Terrific Broth (Sambrook, J. *et al.*, in: *Molecular Cloning: A laboratory manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, p. A.2 (1989)) complemented with 50 µg/ml kanamycin in a 1 Liter shaker flask. Cells were harvested by centrifugation at the end of the exponential growth phase (approximately 16 hr), typically yielding 10 grams of biomass net weight per liter. Covalently closed circular pDNA was isolated by a modified lysis procedure (Horn, N.A. *et al.*, *Human Gene Therapy* 6: 565-573 (1995)) followed by standard double CsCl-ethidium bromide gradient ultracentrifugation with an average yield of approximately 5 mg per liter.

20 Plasmids were ethanol precipitated and resolubilized in saline at 4°C and dialyzed against saline. Endotoxin content was determined by the *Limulus* Amebocyte Lysate assay (Associates of Cape Cod, Inc., Falmouth, MA). All plasmid preparations were free of detectable RNA. Endotoxin levels were less than 0.6 Endotoxin Units/µg of plasmid DNA. The spectrophotometric

25 A260/A280 ratios were between 1.75 and 2.0.

Example 3

In Vitro Evaluation of Biological Activity of IFN ω and IFN α

5 To assure that the interferon plasmid DNA used in the following examples encoded biologically active interferon, cell proliferation and antiviral assays were performed. All culture medium used in this and following examples was obtained from Life Technologies (Gaithersburg, MD), and all serum was obtained from HyClone (Logan, Utah).

10 UM449 cells (American Type Culture Collection, Rockville, MD) were plated at a concentration of 2×10^5 cells per well in a 6 well plate and incubated for 24 hours. Plasmid DNA and the lipid, DMRIE/DOPE (1:1) were each diluted to a concentration of 1 mg in 0.5 ml Optimem medium (Life Technologies, Gaithersburg, MD). The lipid DMRIE/DOPE consists of the
15 cationic lipid (\pm)-N-(2-hydroxyethyl)-N,N-dimethyl-2,3-bis(tetradecyloxy)-1-propanaminium bromide (DMRIE) and the neutral lipid dioleoylphosphatidylethanolamine (DOPE) at a 1:1 mol:mol ratio (Felgner et al., *J. Biol. Chem.* 269:2550-2561, 1994). DMRIE/DOPE has been shown to be effective for both *in vitro* (Felgner et al., *J. Biol. Chem.* 269:2550-2561, 1994) and *in vivo* transfection (Stopeck et al., *J. Clin. Oncol.* 15:341-349, 1997 and Rubin et al., *Gene Ther.* 4:419-425, 1997). The lipid mixture and the DNA mixture were then gently mixed. Medium was removed from the
20 cells which were rinsed gently with PBS, followed by addition of the DNA:lipid mixture (1 ml/well). After incubating the cells for 4-5 h at 37°C, one ml of Optimem with 30% fetal calf serum (FCS) was added to each well. Following an overnight incubation at 37°C, one ml of Optimem with 10% FCS was added to each well. Tissue culture supernatants were collected 48 h after the start of the *in vitro* transfection.

a. Antiproliferative activity

30 To evaluate the antiproliferative and hence, anti-tumor activity of IFN ω and IFN α , supernatants from the above described UM449 cells transfected with the interferon or control plasmid DNA were tested in a cell

proliferation assay of murine or human tumor cell lines (ATCC, Rockville, MD) using the Boehringer Mannheim (Indianapolis, IN) Cell Proliferation Kit II (XTT). Murine or human tumor cells were plated in 96 well plates at the desired concentration (cell concentration varied with each cell line evaluated, for example, at a concentration of 5×10^3 cells/ml for B16F10 cells and 5×10^4 cells/ml for the Cloudman S91 and glioma 261 cells). The plates were incubated at 37°C for 24 hours followed by addition of tissue culture supernatants from UM449 cells *in vitro* transfected with either interferon plasmid DNA or control plasmid DNA. As a positive control for mIFN α plasmid DNA, mIFN α protein (ICN Pharmaceuticals Inc., Costa Mesa, CA) was serially diluted and added to the wells. For the hIFN plasmid DNA, an interferon reference standard (human leukocyte interferon, Sigma Chemical Co., St. Louis, MO) was included in each assay. Following a 24-72 hour incubation, at 37°C, 50 μ l of XTT/ECR substrate was added to each well. Plates were incubated for 6-24 hours at 37°C and the optical density (OD) at 490 nm was determined. Increasing amounts of interferon result in inhibition of cell proliferation and a reduction in the OD₄₉₀. The percent reduction in cell proliferation due to addition of the supernatants was determined by the formula:

$$1 - \frac{\text{OD}_{490} \text{ of cells incubated with interferon plasmid DNA supernatants}}{\text{OD}_{490} \text{ of cells incubated with control plasmid DNA supernatants}} \times 100$$

As shown in Table 1, both human interferons displayed the characteristic potent anti-proliferative activity against a wide variety of human tumor cell lines, with the most sensitive line being the NIH-OVCAR3 ovarian line and the least sensitive being the SK-OV-3 ovarian line. Also, the supernatants from the mIFN α pDNA (VR4111)-transfected UM449 cells inhibited the proliferation of murine B16F10 melanoma (generous gift from Dr. Suzuki at the University of Texas, Galveston, Texas), murine Cloudman melanoma S91 (American Type Culture Collection, Rockville, MD), and

murine glioma 261 cell lines (Division of Cancer Treatment Tumor Repository, National Cancer Institute, Frederick Cancer Research and Development Center, Frederick, MD) by 40, 42 and 17%, respectively.

5

Table 1. IFN ω (VR4150) and IFN α (VR4102) *in vitro* biological assay: anti-proliferation activity against human tumor cell lines

Cell line (tissue type)	% reduction in cell proliferation (compared to control plasmid DNA supernatants)	
	Interferon ω	Interferon α
NIH-OVCAR3 (ovarian)	60	43
SCC-4 (squamous)	36	36
ACHN (renal)	41	35
A431 (epidermoid)	24	19
SCC-15 (squamous)	29	29
U87MG (glioblastoma)	36	30
A375 (melanoma)	24	21
PC3 (prostate)	20	22
UMUC3 (bladder)	14	6
A549 (lung)	17	15
MCF7 (breast)	18	18
SK-OV-3 (ovarian)	<10	<10

10

b. Antiviral Activity

An antiviral assay was performed to evaluate the ability of the supernatants from the interferon plasmid DNA-transfected cells to protect murine L929 cells or human A549 cells from infection by murine encephalomyocarditis (EMC) virus (Assay performed at IIT Institute, Chicago, IL). *In vitro* transfections were performed as described above and supernatants were collected from cells transfected with either VR4151 (hIFNT), VR4112 (hIFN α), VR4111 (mIFN α) or VR1055 (control). Antiviral activity of the supernatants was performed by IIT Research Institute (Chicago, IL). The antiviral assay evaluated the degree of

15

20

protection of either human A549 or murine L929 cells from infection with murine encephalomyocarditis (EMC) virus. Briefly, 2.5×10^4 L929 cells were plated into 96-well plates and incubated for 24 h. Tissue culture supernatants were serially diluted and added to the L929 cells which were incubated for another 24 h. Supernatants were then removed from the wells, the cells were washed and murine EMC virus was added to each well at a multiplicity of infection of 0.04. Assay plates were incubated further for 24 h followed by removal of supernatants, washing of wells, fixation with 5% formalin and staining with 1% crystal violet. Samples with interferon activity protected the cells from virus infection, resulting in darkly stained cell monolayers.

Supernatants from UM449 cells transfected with VR4151, VR4112, or VR4111 had antiviral activity of 30,000, 3,000 or 30 Units/ml, respectively, on human A549 cells. When evaluated for antiviral activity on the murine L929 cell line, supernatants from UM449 cells transfected with VR4151, VR4112, or VR4111 had antiviral activity of 300, 1000 and 30,000 Units/ml, respectively (Table 2) showing species specificity of the hIFNs for human cells and mIFNs for mouse cells.

Table 2. Antiviral Activity of interferon Plasmid DNA

Plasmid	Interferon (Units/ml)	
	Human cell line	Murine cell line
VR4151 (hIFN ω)	30,000	300
VR4112 (hIFN α)	3,000	1,000
VR4111 (mIFN α)	30	30,000

Example 4

Systemic Interferon Therapy; Intramuscular Administration of Cytokine-Expressing Plasmids

Cell lines and tumor models

Murine B16F10 cells were grown in RPMI-1640 (GibcoBRL) and 5% fetal bovine serum (FBS). Murine Cloudman S91 cells were grown in Ham's F-10 medium with 25 mM Hepes, 0.1 mM, non-essential amino acids, 1 mM sodium pyruvate, 0.05 mM β -mercaptoethanol, 2.5% FBS and 12.5% horse serum. Human melanoma UM449 cells were grown in RPMI 1640 with 10% FBS.

Murine glioma 261 tumor fragments and M5076 reticulum cell sarcoma cells were obtained from the Division of Cancer Treatment Tumor Repository (National Cancer Institute, Frederick Cancer Research and Development Center, Frederick, MD). The glioma 261 tumor fragments (2mm³) were initially implanted into the inguinal region of C57BL/6 mice using a 13 g trocar (Popper Sons, Inc., New Hyde Park, NY). Tumors which grew in the mice were used to establish a tumorigenic cell line. Minced tumor fragments were placed in Iscove's tissue culture medium with 10% FBS. Glioma 261 tumor cells began to attach to the flasks after several days, and the cells were propagated using standard tissue culture techniques. The M5076 cells were grown as ascites in C57BL/6 mice and frozen in liquid nitrogen. Human A431 cells were obtained from the American Type Culture Collection and were grown in DMEM and 10% FBS.

C57BL/6, DBA/2, nude (*nu/nu*) and beige-nude (*bg/nu/xid*) female mice between the ages of 6-8 weeks were obtained from Harlan Sprague Dawley (San Diego, CA). All animal experiments in this and the following examples were conducted in accordance with Vical's Institutional Animal Care and Use Committee as well as the standards set forth in the National Research Council guidelines concerning animal care and use.

To establish subcutaneous B16F10 melanoma tumors, C57BL/6, nude or nude-beige mice were injected subcutaneously on the flank with 10⁴

B16F10 cells. The Cloudman melanoma model was established by subcutaneous injection of 10^5 Cloudman S91 cells on the flank of DBA/2 mice and the glioma 261 model was established by subcutaneous injection of 5×10^4 glioma 261 cells on the flank of C57BL/6 mice. To establish human epidermoid carcinomas, nude mice were injected subcutaneously on the flank with 5×10^3 A431 cells.

To establish intradermal M5076 tumors and liver metastases thereof, C57BL/6 mice were injected intradermally with 10^5 M5076 reticulum cell sarcoma cells. In this model, primary intradermal tumors spontaneously metastasize to the liver. On day 29 after tumor cell injection, mice were sacrificed, livers removed and fixed in 10% buffered formalin (Fisher Scientific, Pittsburgh, PA), and the liver nodules were counted.

To establish lung metastases of B16F10 melanoma, C57BL/6 mice were injected intravenously with 2×10^4 B16F10 cells. On day 25 after tumor cell injection, the mice were sacrificed, lungs removed and fixed in 10% buffered formalin, and the lung nodules were counted.

To monitor the primary tumor growth, tumor sizes were determined 2-3 times per week by measuring with calipers ($l \times w \times h$), and tumor volumes were determined using the formula: tumor volume (mm^3) = $0.52 (l \times w \times h)$.

Tumor volume was analyzed using the Mann-Whitney U non-parametric Statistical Test to identify groups having significantly different mean weights. Mouse survival was analyzed using a Kaplan-Meier survival plot followed by a Logrank (Mantel-Cox) test to identify significant differences in survival between groups. Differences were considered statistically significant when the p value was less than or equal to 0.05.

Intramuscular injections

Fifty to 100 μg of plasmid DNA in 50 μl of saline was injected into the rectus femoris muscle of each hind leg for a total DNA dose of 100 to 200 mg. The muscle injections were performed using a 300 μl sterile tuberculin syringe fitted with a 28G $\frac{1}{2}$ needle (Becton Dickinson) and a plastic collar cut

from a 200 µl micropipette tip. The collar length was adjusted to limit the needle from penetrating further than 2 mm into the rectus femoris muscle.

Serum levels of interferon following intramuscular injection of interferon plasmid DNA

Serum samples from C57BL/6 mice injected intramuscularly with either VR4111 (mIFN α plasmid) or VR1055 (control plasmid DNA) were analyzed in a mIFN α ELISA (n=10). For the ELISA, 96-well plates (Immulon 4HBX high binding plates from Dynex Technologies, Chantilly, VA) were coated with rat anti-mouse IFN α monoclonal antibody (mAb) from Caltag Laboratories (Burlingame, CA) at a concentration of 5 µg/ml in 100 mM sodium carbonate buffer, pH 9.5 (50 µl per well). Plates were incubated with the coating mAb for 16 hours at 4°C. The plates were then washed 3 times with a wash buffer (phosphate-buffered saline (PBS)), pH 7.2 and 0.05% Tween-20 (Sigma, St. Louis, MO)). The plates were blocked in PBS containing 3% bovine serum albumin (BSA, Sigma) and 0.05% Tween-20 (400 µl per well) and incubated for 24 hours at 4°C followed by washing three times with wash buffer.

Serum samples (10 µl) from mice injected intramuscularly with VR4111 (mIFN α) were mixed with 40 µl of assay buffer (PBS, 1% BSA, 0.05% Tween-20) and the mixture was added to each assay well. The positive control was mIFN α polypeptide (Biosource International, Camarillo, CA), which was serially diluted in assay buffer and 50 µl was added to the positive control wells. The negative control was serum from mice injected intramuscularly with VR1055. After adding the test samples and controls, the plates were incubated for 16 hours at 4°C. The plates were then washed 6 times with wash buffer, followed by addition of a sheep anti-mouse IFN α polyclonal antibody (pAb) (Biosource International, Camarillo, CA). The pAb was added at a 1:500 dilution in assay buffer (50 µl per well) and the plates were incubated for 5 hours at room temperature.

Following incubation with the pAb, the plates were washed six times with wash buffer followed by addition of anti-sheep IgG conjugated with peroxidase (Sigma) at a 1:5000 dilution in assay buffer (50 μ l per well) and incubated for 1 hour at room temperature. The plates were washed 6 times with wash buffer and 200 μ l of 3,3',5,5'-tetramethylbenzidine liquid substrate (TMB) (Sigma Chemical Co.) was added per well. Plates were incubated at room temperature for 30 minutes, followed by determination of the optical density of the wells at 650 nm. A standard curve was generated by plotting the ng/ml of mIFN α polypeptide versus the optical density at 650 nm. The concentration of mIFN α in the test serum samples was determined from the linear portion of the mIFN α standard curve. The sensitivity of the mIFN α ELISA was 50 ng/ml.

C57BL/6 mice injected intramuscularly with VR4111 had detectable serum levels of mIFN α after 5 intramuscular injections of 100 μ g VR4111 (twice a week for two weeks, followed by one injection the next week). The average serum level of mIFN α after 5 intramuscular injections of VR4111 was 1465 ng/ml (average of 16 mice). At the time of this study, no commercial mIFN α ELISA kit had been developed. Since sensitivity of the in-house mIFN α ELISA is 50 ng/ml, lower serum levels of mIFN α could exist in the mice at earlier timepoints, but we were unable to detect this in our ELISA.

To determine the serum levels of hIFN ω , C57BL/6 or nude mice received a single intramuscular injection of 100 μ g of VR4151 (hIFN ω plasmid DNA) or VR1055 (control plasmid DNA) (50 μ g per leg bilaterally) in the rectus femoris. Serum samples were collected daily over a two week period and analyzed in the hIFN ω ELISA kit (Alexis, San Diego, CA) which was sensitive to 2 pg/ml. Serum samples were collected from 4-5 mice per day. In the C57BL/6 mice, measurable serum levels of hIFN ω were detected as early as one day after injection (69 pg/ml) (Figure 2A). In these mice, peak serum levels were found six days after injection (254 pg/ml) and expression continued out to day 14 (50 pg/ml), the final timepoint of the study.

In nude mice, serum levels of IFN α were found as early as one day after injection (133 pg/ml). Peak serum levels were found on day 7 (648 pg/ml) and expression continued out to day 14 (134 pg/ml), the final time point of the study (FIG. 2B). Thus, interferon could be detected in the serum after a single intramuscular injection of an interferon-encoding plasmid DNA.

Systemic interferon treatment inhibits primary tumor growth

As shown in FIGs. 3-5 and FIG. 7A, mice bearing different tumors were found to significantly benefit from intramuscular injection of different cytokines. To test the efficacy of IFN α plasmid, C57BL/6 mice bearing subcutaneous B16F10 melanoma, subcutaneous glioma 261, or intradermal M5076 tumors, or DBA/2 mice bearing subcutaneous Cloudman melanoma were injected with 100 μ g either of VR4111 (mIFN α) or VR1055 (control), twice per week for three weeks, beginning on day 4 after tumor cell injection (n = 8-10 mice per group). In all three subcutaneous tumor models, the mice treated intramuscularly with VR4111 had a significant reduction in tumor volume (p<0.05) (FIGs. 3A, 3C, and 3E), and a significant enhancement of survival (p<0.02) compared to the mice that received the control plasmid (FIGs. 3B, 3D, and 3F). In the intradermal tumor model, mice treated with intramuscular VR4111 had a significant reduction in primary tumor volume (p<0.001) compared to the mice that received the control plasmid (FIG. 7A).

To compare the efficacy of IL-2, IL-12 to IFN α plasmids, C57BL/6 mice bearing subcutaneous B16F10 melanoma were injected with 100 μ g of VR4111 (mIFN α), VR4001 (mIL-12), VR1110 (mIL-2), or VR1012 (control) (n = 15-16 mice per group) twice per week for three weeks. Mice receiving intramuscular injections of VR4111 had a significant reduction in tumor growth (p < 0.0002) (FIG. 4A) by day 17 as well as a significant increase in survival (p = 0.00001) (FIG. 4B). By day 28 of the study, 100% of the VR4111-treated mice were still alive, compared to only 20% of the VR1012-treated mice. Mice treated with VR1110 had a modest reduction in tumor

-60-

growth by day 17 ($p < 0.02$) (FIG. 4A) but did not have an increase in survival compared to the VR1012-treated mice (FIG. 4B). Mice treated with VR4001 also had a modest reduction in tumor growth by day 17 ($p < 0.03$) (FIG. 4A) as well as a significant increase in survival ($p = 0.02$) (FIG. 4B). By day 28, 55% of the mice treated with VR4001 were alive, compared to 20% of the VR1012-treated mice.

To test the efficacy of IFN ω , mice bearing human A431 tumors between 30-80 mm³ were injected intramuscularly with 200 μ g of either VR4151 (hIFN ω) or VR1055 (control) twice per week for three weeks ($n=15$). Mice bearing subcutaneous A431 tumors and injected intramuscularly with VR4151 had a significant reduction in tumor volume ($p < 0.05$) (FIG. 5A) and a significant increase in survival ($p < 0.05$), compared to the mice that received the control plasmid (FIG. 5B).

Systemic mIFN α plasmid DNA treatment inhibits the growth of tumor metastases

As shown in FIG. 6 and FIG. 7, mice bearing different tumor metastases were found to significantly benefit from intramuscular injection of IFN α . C57BL/6 mice bearing lung metastases of B16F10 melanoma were injected intramuscularly with 100 μ g of either VR4111 (mIFN α) or VR1055 (control) twice per week for three weeks, beginning on day 4 after tumor cell injection ($n=10$). On day 25 after tumor cell injection, the mice were sacrificed, lungs removed and fixed in 10% buffered formalin (Fisher Scientific, Pittsburgh, PA) followed by counting of lung nodules.

While 70% of the control plasmid-treated mice had lung nodules that were too numerous to count, 80% of the mice treated with mIFN α plasmid DNA had 10 or fewer nodules (FIG. 6). TNTC denotes lungs with nodules that were too numerous to count.

In the liver metastases model, C57BL/6 mice bearing intradermal M5076 murine reticulum sarcoma were injected intramuscularly with 100 μ g of either VR4111 or VR1055 twice per week for three weeks, beginning on

day 4 after tumor cell injection (n=10-13 mice per group). On day 29 after tumor cell injection, the mice were sacrificed, livers removed and fixed in 10% buffered formalin (Fisher Scientific, Pittsburgh, PA) followed by counting of liver nodules.

5 While the control plasmid-treated mice had a mean of 190 hepatic tumor nodules or had nodules that were too numerous to count, mIFN α plasmid DNA-treated mice had a mean of 35 hepatic tumor nodules (FIG. 7B).

10 These results demonstrate that intramuscular injection of mIFN α plasmid DNA can effectively inhibit the growth of both primary and metastatic lesions. Thus, for patients with metastatic disease, intramuscular administration of therapeutic plasmid DNAs would be advantageous for the treatment of undiagnosed or inaccessible metastatic lesions.

Regimen optimization of mIFN α therapy in the B16F10 melanoma model

15 A regimen optimization study was conducted to evaluate the antitumor efficacy of fewer injections and/or a lower dose of VR4111(mIFN α) in the subcutaneous B16F10 melanoma model. C57BL/6 mice were injected with either 100 or 50 μ g of VR4111 or VR1055 over a 6 week period (n=10). Mice
20 received intramuscular injections either twice per week, once per week or once every other week. All intramuscular injections began four days after the initial subcutaneous B16F10 tumor cell injection. Mice which received intramuscular injections of 100 μ g of VR4111 at any of the time courses had a significant reduction in tumor volume ($p \leq 0.005$) and a significant increase in
25 survival ($p = 0.007$) (FIGs. 8A and 8B). The mice which received the 100 μ g dose of VR4111 once every other week for 6 weeks had a total of only three intramuscular injections with significant antitumor efficacy. In contrast, mice receiving the 50 μ g dose of VR4111 revealed a dose response based on the frequency of injection. While mice injected with 50 μ g of VR4111 once or
30 twice per week had a significant reduction in tumor volume ($p \leq 0.03$), and a significant increase in survival ($p = 0.002$), mice injected only once every other

week did not have a significant antitumor response for either tumor growth or survival (FIGs. 8C and 8D).

Mechanism of mIFN α Antitumor Effect

5

To investigate the role of natural killer (NK) and T cells in mediating the antitumor effect of systemically delivered mIFN α , VR4111 or VR1055 was administered intramuscularly to nude mice (which are T cell deficient) and to beige-nude mice (which are NK and T cell deficient) bearing subcutaneous B16F10 melanoma tumors. Beginning on day 4 after injection of 10⁴ B16F10 cells, 50 μ g of plasmid DNA in 50 μ l of saline was injected into the rectus femoris muscle of each hind leg for a total DNA dose of 100 μ g twice per week for three weeks (n=15 mice per group).

10

15

There was neither a significant reduction in tumor volume nor enhancement of survival in the nude mice (FIGs. 9A and 9B) or beige-nude mice (FIGs. 9C and 9D). These results suggest that T cells may be required for the mIFN α antitumor response. NK cells appeared not to be required for the antitumor effect since nude mice (NK⁺) did not have a greater antitumor response compared to the beige-nude mice (NK⁻).

20

25

30

To further explore the role of T cells in the antitumor effect of mIFN α DNA therapy, C57BL/6 mice bearing subcutaneous B16F10 tumors were injected with depleting doses of monoclonal antibodies (mAbs) specific for either CD4⁺ or CD8⁺ T cells. For depletion of T cell subsets, anti-CD4 (clone GK1.5, rat IgG) and anti-CD8 (clone 2.43, rat IgG) hybridomas (American Type Culture Collection, Rockville, MD) were used to generate the corresponding mAb. The anti-CD8 hybridoma was grown as ascites in nude mice and the mAbs were purified from ascites using ion exchange chromatography (Harlan Bioproducts for Science, San Diego, CA). The anti-CD4 hybridoma was grown *in vitro* with Dulbecco's Modified Eagle Medium, 10% fetal bovine serum and low IgG. The anti-CD4 mAb was purified from tissue culture supernatant by ammonium sulfate precipitation to 30%. The

protein pellet was resolubilized and extensively dialyzed in Dulbecco's $\text{Ca}^{2+}/\text{Mg}^{2+}$ -free phosphate-buffered saline (Zymed Laboratories Inc., San Francisco, CA).

Beginning on day 4 after subcutaneous injection with 10^4 B16F10 cells, mice were injected intramuscularly with 100 μg of either VR4111 or VR1055 twice per week for three weeks. For depletion of CD4^+ and CD8^+ T cells, mice were injected intraperitoneally with 500 μg of either the anti-CD4 mAb (clone GK1.5, rat IgG) or anti-CD8 mAb (clone 2,43, rat IgG) one day prior to each intramuscular DNA injection ($n = 10$ mice per group). Control tumor-bearing mice were injected intraperitoneally with 500 μg of normal rat IgG (Simga Chemical Co., St. Louis, MO) ($n=10$). To assure complete depletion, sentinel mice were injected according to the same regimen, and once per week, spleens were collected, dissociated and assessed for the presence of CD4^+ and CD8^+ T cells. Spleen cells were stained with FITC-conjugated anti-CD4 and PE-conjugated anti-CD8 mAbs (Pharmingen, San Diego, CA) and analyzed by flow cytometry (Cytometry Research Services, San Diego, CA). The depletion of CD4^+ and CD8^+ T cells was consistently greater than 98%, as determined by cytometry.

The $\text{mIFN}\alpha$ DNA therapy significantly reduced tumor growth ($p \leq 0.002$) and enhanced survival ($p \leq 0.008$) of both normal mice and mice depleted of CD4^+ T cells, compared to mice injected with control plasmid and treated with normal IgG (FIGs. 10A and 10B). These results suggest that CD4^+ T cells are not required for the $\text{mIFN}\alpha$ antitumor effect. In contrast, mice depleted of CD8^+ T cells and injected with $\text{mIFN}\alpha$ DNA displayed tumor volumes and survival profiles that were not significantly different from mice treated with the control plasmid (FIGs. 10A and 10B). This result suggests that CD8^+ cells are involved in the $\text{mIFN}\alpha$ antitumor response.

Example 5

Local Interferon Therapy: Intratumoral Administration of interferon Plasmids

5 The anti-tumor activity of IFN ω and IFN α was evaluated *in vivo* in nude mice bearing subcutaneous human ovarian (NIH-OVCAR3) or human melanoma (A375) (nude/human/xenograft model), or in C57BL/6 mice bearing murine melanoma (B16F10) tumors following intratumoral
10 administration of DNA complexed with a cationic lipid.

Cell Lines and Tumor Models

15 Athymic nude (*nu/nu*) and C57BL/6 mice between the ages of 6-10 weeks were obtained from Harlan Sprague Dawley (San Diego, CA).

Human A375 melanoma cells and human NIH-OVCAR3 ovarian carcinoma cells were obtained from the American Type Culture Collection (Rockville, MD), and grown in Dulbecco's modified Eagle's medium (GibcoBRL, Gaithersburg, MD) supplemented with 10% FBS. B16F10 cells
20 were a generous gift from Dr. Suzuki at the University of Texas (Galveston, Texas). Cells were grown in RPMI-1640 (GibcoBRL) and 5% fetal bovine serum (FBS).

To establish subcutaneous A375 melanoma tumors and subcutaneous NIH-OVCAR3 ovarian tumors, athymic nude/nude mice (10 mice/group)
25 were injected subcutaneously with 5×10^6 A375 cells and 5×10^7 NIH-OVCAR3 cells respectively. To establish subcutaneous murine B16F10 melanoma tumors, C57BL/6 mice were injected subcutaneously with 10^4 B16F10 cells.

Mice were monitored for tumor growth and survival. Tumor sizes
30 were determined 3 times per week by measuring with calipers (l x w x h) and tumor volumes were determined using the formula: tumor volume (mm³) = $0.52 (l \times w \times h)$. Statistical analysis was done as described in Example 4.

Local Interferon Plasmid DNA Inhibits Tumor Growth

Established subcutaneous A375 human melanoma and NIH-OVCAR3 human ovarian carcinoma tumors in nude mice were transfected *in vivo* by intratumoral administration of pDNA/DMRIE/DOPE (DNA: lipid) complexes (n=10). When tumors became palpable (80-300 mm³, at day 27 post tumor cell implant for the A375 cells, and at day 41 post tumor cell implant for the NIH-OVCAR3 cells), mice were injected intratumorally with 100 µg of VR4112 (hIFN α), VR4151 (hIFN γ), VR1055 (control) or VR1012 (control) complexed with DMRIE/DOPE (1:1 DNA:DMRIE mass ratio). Tumor bearing animals were treated intratumorally with DNA: lipid for 6 consecutive days followed by 5 treatments every other day for a total of 11 treatments (A375 melanoma model), or every other day for a total of 11 treatments (NIH-OVCAR3 ovarian cancer model).

As shown in FIG. 11A, in the A375 model of melanoma, the direct intratumoral injection of VR4151: lipid complex for 6 consecutive days, followed by 5 additional injections every other day (100 µg plasmid DNA/injection, total of 11 injections), resulted in a statistically significant slowing of tumor growth, as compared to the control ($p < 0.03$, days 40-44).

As shown in FIG. 2B, in the NIH-OVCAR3 model of ovarian cancer, the direct intratumoral injection of VR4151: lipid complex every other day for a total of 11 injections (100 µg plasmid DNA/injection), resulted in sustained and statistically significant reduction in tumor growth as compared to the control plasmid control ($p < 0.001 - 0.05$, Days 45-65). A similar treatment regimen with VR4112: lipid was found to have a moderate effect on tumor growth which only reached statistical significance vs. the control plasmid at two time-points during the study ($p < 0.05$, days 53 and 57).

As shown in FIG. 12A, in the B16F10 melanoma model, direct intratumoral injection of mIFN α resulted in decrease in tumor volume. Once palpable tumors were established (80-300mm³) on day 12 after tumor cell injection in C57BL/6 mice bearing subcutaneous B16F10 melanoma tumors, mice were injected intratumorally with 100 µg of either VR4101 (mIFN α) or

-66-

VR1012 (control) complexed with the cationic lipid, DMRIE/DOPE at a DNA/DMRIE mass ratio of 1:1 in a 100 μ l volume. Mice received six consecutive intratumoral injections of either VR4101 or VR1012 (n=10 mice per group). As shown in FIG. 12A, although not statistically significant, intratumoral injection of VR4101 resulted in a 54% reduction in tumor volume by day 19 of the study. As shown in FIG. 12B, a significant increase in survival ($p = 0.02$) was found for the VR4101-treated mice, compared to the mice that received VR1012.

Example 6

Local Cytokine Therapy: Intraperitoneal Administration of Cytokine-expressing Plasmids

The goal of this example is to show that the present invention provides an effective method of treating malignant tumors of murine ovarian carcinoma via intraperitoneal (i.p.) injection of cytokine-expressing plasmid DNA. Since late-stage ovarian carcinoma is usually limited to the peritoneal cavity, it was envisioned that continuous secretion of a cytokine in this cavity would produce beneficial anti-tumor immune response. In particular, the present example clearly shows that the ovarian cancer therapy by intraperitoneal injection of a cytokine-expressing plasmid DNA:lipid complexes (1) results in sustained levels of the cytokine in the ascites, avoiding the need for frequent injections of the protein (in contrast to intraperitoneal injection of recombinant cytokine wherein the cytokine level declines shortly after injection), (2) targets tumor ascites, rather than peritoneal tissues (suggesting that systemic cytokine side effects should be reduced using this method), (3) inhibits tumor growth and enhances survival, and (4) can be combined with debulking of tumor ascites to enhance the antitumor effect.

Cell Lines and Tumor Models

As a model for human ovarian cancer, the murine ovarian teratocarcinoma (MOT) model in C3H/HeN mice was used. MOT exhibits many of the characteristics of late-stage human ovarian cancer including peritoneal spread, production of tumor ascites and tumor cell blockage of lymphatics (Ozols et al., 1979 and Fekete et al., 1952).

Murine ovarian teratocarcinoma (MOT) cells were obtained from Dr. Robert Knapp and Dr. Robert C. Bast at the Dana-Farber Cancer Center (Boston, MA). The MOT cells (10^5) were grown by serial intraperitoneal (i.p) transplantation in C3H/HeN mice and a stock of the cells was frozen in liquid nitrogen.

CTLL-2 cells were obtained from the American Type Culture Collection (ATCC, Rockville, MD) and were grown in RPMI 1640 with glutamine, 1% sodium pyruvate, 1% penicillin-streptomycin (Life Technologies, Gaithersburg, MD), 10% fetal bovine serum (HyClone, Logan, Utah) and 10 U/ml murine IL-2 (Boehringer Mannheim, Indianapolis, IN).

C3H/HeN and nude (*nu/nu*) female mice between the ages of 6-10 weeks were obtained from Harlan Sprague Dawley (San Diego, CA). All animal experiments were conducted in accordance with Vical's Institutional Animal Care and Use Committee as well as the standards set forth in the National Research Council guidelines concerning animal care and use.

To establish i.p. MOT tumors, C3H/HeN mice were injected i.p. with 10^5 MOT cells in 100 μ l of medium. In the MOT tumor model, tumor growth is typically monitored by weighing the mice which reflects the increase in volume of tumor ascites (Berek et al., *Cancer Res.*, 44:1871-1875, 1984). The nude mouse study were performed in the same manner as the studies in C3H/HeN mice with injection i.p. of 10^5 MOT cells and monitoring the weight of the mice. Statistical analysis on mouse weight and survival was done as described in Example 4.

Preparation of plasmid DNA:lipid complexes and intraperitoneal injection

To yield a pDNA:DMRIE mass ratio of 1:1, 100 µg of VR1110 (mIL-2) was diluted in 500 µl 0.9% saline (Radix Labs, Eau Claire, WI),
5 DMRIE/DOPE lipid (100 µg DMRIE) was diluted in 500 µl of 0.9% saline in a separate vial, and the pDNA and cationic lipid were combined and vortexed for 5 seconds. To yield a pDNA:DMRIE mass ratio of 5:1, 500 µg of mIL-2 pDNA was diluted in 500 µl 0.9% saline (Radix Labs, Eau Claire, WI),
10 DMRIE/DOPE lipid (100 µg DMRIE) was diluted in 500 µl of 0.9% saline in a separate vial, and the pDNA and cationic lipid were combined and vortexed for 5 seconds.

The 1 ml pDNA:DMRIE/DOPE (DNA: lipid) complex was injected i.p. into mice bearing i.p. MOT tumors on various days after tumor cell implant. Control MOT tumor-bearing mice received i.p. injections of VR1012
15 (control): lipid at the same ratio (1:1 pDNA:DMRIE, 100 µg pDNA) and were injected i.p. on the same days as the cytokine-expressing or reporter gene treatment groups.

Intraperitoneal injection of plasmid DNA:lipid results in targeted expression in tumor ascites

The pDNA:lipid therapy was evaluated for the ability to target malignant cells within a cavity. C3H/HeN mice were injected i.p. with 10⁵ MOT cells followed by i.p. injection of 100 µg of VR1223 (luciferase): lipid
25 (1:1 pDNA:DMRIE mass ratio) on days 5 and 6 after tumor cell implant. Control MOT tumor-bearing mice were injected i.p. with 100 µg of either VR1012: lipid or VR1223 without lipid on days 5 and 6 after MOT tumor cell injection. An additional group of control mice did not receive MOT tumor cells and were injected i.p. with VR1223: lipid on the same days as the other
30 treatment groups (n=3). Three days later the mice were euthanized and tumor ascites and tissues (liver, kidney, spleen, diaphragm, intestine and ovary) were collected. Luciferase was extracted from the tissues by freeze-thawing and

grinding of the samples in cell lysis reagent (Promega, Madison, WI) as previously described (Hartikka et al., *Hum. Gene Ther.*, 7:1205-1217, 1996). The tumor ascites was diluted 1:5 in cell lysis reagent followed by three cycles of freeze-thaw and collection of supernatant from the cell lysate. Samples were read in a microplate luminometer (Dynatech, Chantilly, VA) following addition of luciferase substrate (Promega, Madison, WI). The relative light units (RLU) of the samples were determined from a standard curve using purified firefly luciferase (Analytical Luminescence Laboratory, Sparks, MD). The protein concentration of each sample was determined using the BCA protein assay kit (Pierce Chemical Company, Rockford, IL). Luciferase levels were expressed as RLU per mg of protein. There may be a decrease in IL-2 pDNA expression from day 1 to day 3 post DNA injection.

On day three, tumor ascites had 900,000 RLU of luciferase/mg, while diaphragm and ovary tissue had only 327 and 16 RLU/mg (FIG. 13). Kidney, liver, spleen and intestinal tissue had no detectable luciferase activity. These results suggest that i.p. injection of pDNA: lipid complexes appears to target the tumor ascites in the peritoneal cavity with limited or negligible transfection of surrounding tissues. Luciferase detected in the diaphragm and ovary tissue was only found in MOT tumor-bearing mice injected with VR1223: lipid. When naive non-tumor bearing mice were injected with the same DNA: lipid complex, no luciferase activity was found in any of the tissues (data not shown). These results suggest that the low levels of luciferase in the diaphragm and ovary in MOT tumor-bearing mice may reflect metastases of tumor cells to these tissues. Tumor-bearing mice injected with luciferase pDNA without cationic lipid had no luciferase activity in either tumor ascites or surrounding tissues, indicating that lipid is required for optimal *in vivo* transfection of ovarian tumor ascites. Mice injected with VR1012: lipid had no detectable luciferase activity in either tumor ascites or tissues.

A follow-up study investigated the specific cell type in the ovarian tumor ascites that was transfected after i.p. injection of a reporter gene pDNA:DMRIE/DOPE complex. On days 5 and 6 after tumor cell implant (10^5

-70-

cells), C3H/HeN mice were injected i.p. with 100 μ g of VR1412 (β -galactosidase (β -gal)): lipid, VR1012: lipid (1:1 DNA:DMRIE mass ratio), or with VR1412 without cationic lipid (n=3 mice per group). One day later the mice were sacrificed and the tumor ascites was collected. The ascites was spun at 2500 rpm for 2 minutes to pellet the cells, and the supernatant was removed. The tumor cells were fixed in 10% buffered formalin (Fisher Scientific, Pittsburgh, PA), placed in a cryomold containing OCT embedding medium (VWR, S. Plainfield, NJ), frozen in isopentane and then stored at -70°C. The embedded and frozen samples were then cryostat sectioned (5 μ m), further fixed (0.5% glutaraldehyde in PBS), washed (PBS), stained with X-gal reagent (1 mg/ml X-gal diluted in PBS containing 5 mM potassium ferricyanide, 5 mM potassium ferrocyanide, and 1 mM magnesium chloride), washed again (PBS), and counterstained with hematoxylin and eosin. (The samples were cryostat sectioned and stained by Pathology Associates (Frederick, MD).)

The ascites from mice treated with either VR1012 or VR1412 without lipid had no β -gal activity in the samples. In contrast, the tumor ascites from mice injected with VR1412: lipid had β -gal staining primarily in the tumor cells (data not shown). In a few slides, several macrophages and lymphocytes were also positive for β -gal while neutrophils were negative for β -gal.

Intraperitoneal injection of IL-2 pDNA:lipid results in sustained expression of IL-2 in tumor ascites

A time-course study was done to determine the length of time that IL-2 could be detected after multiple i.p. injections of IL-2 pDNA: lipid or a single i.p. injection of either IL-2 protein or IL-2 pDNA: lipid in mice bearing i.p. ovarian tumor ascites. Beginning on day 5 after tumor cell injection, mice were injected multiple times with VR1110 (mIL-2): lipid or a single time with VR1110: lipid or IL-2 protein. Mice were sacrificed at various times and ascites and serum were analyzed for IL-2 levels. Ascites was collected from the sacrificed mice, the samples were spun at 14,000 rpm for 2 minutes and the

supernatant was harvested. Blood was collected from the mice on the same day as the ascites collection and the serum was separated from blood cells by allowing the blood to clot in serum separator tubes (Microtainer, Becton Dickinson, Franklin Lakes, NJ) followed by centrifugation at 14,000 rpm for 10 minutes and collection of the serum supernatant. IL-2 concentration (pg/ml) in the ascites and serum samples was determined using a murine IL-2 ELISA (R & D Systems, Minneapolis, MN). Since the volume of tumor ascites increases over time, the volume of ascites was also determined for each mouse. The total concentration of IL-2 in ascites was determined using the formula: IL-2 pg/ml x ml of ascites = pg IL-2/ total ascites. Serum IL-2 concentrations were reported as pg/ml.

IL-2 in serum and ascites after multiple injections of IL-2 pDNA: lipid.

Beginning on day 5 after tumor cell injection, mice were injected with VR1110: lipid for either 2, 4 or 6 consecutive days or with control VR1012: lipid for 6 consecutive days (100 µl of plasmid DNA complexed with 100 µl DMRIE/DOPE at 1:1 mass ratio in a total volume of 1 ml). An additional group of mice received VR1110 that was not complexed with DMRIE/DOPE. Every two days for up to 17 days after the DNA:lipid injections, 3-4 mice were sacrificed per treatment group and ascites and serum were collected and analyzed using mIL-2 ELISA assay as described above.

Two injections of VR1110: lipid (100 µl DNA per day, 1:1 DNA:cationic lipid mass ratio) into mice bearing i.p. MOT ovarian tumors yielded high levels of IL-2 protein in the tumor ascites. One day after VR1110: lipid i.p. injection, 28,000 pg/ml of IL-2 was measured in the tumor ascites (Table 3). IL-2 expression in the ascites continued for over two weeks after DNA:lipid injection with 750 pg/ml detected 17 days after the last pDNA:lipid injection. Mice injected with either four or six consecutive injections of VR1110: lipid also had high levels of IL-2 in the tumor ascites, however, due to the very high expression levels after more frequent VR1110: lipid injections, IL-2-mediated side effects were noted in the mice which did

-72-

not survive beyond day 9 or 13 after DNA injection. In contrast, two consecutive injections of VR1110 plus DMRIE/DOPE did not cause observable IL-2 side effects yet IL-2 expression levels remained high for over 2 weeks. The IL-2 expressed after i.p. DNA:lipid injection of MOT-bearing mice appeared to remain localized in the peritoneal cavity as the IL-2 serum levels after VR1110: lipid i.p. injection were always less than 10% of the levels in the tumor ascites. Injection of VR1110 without lipid yielded very low levels of IL-2 in ascites and serum (0-32 pg/ml). Injection of the control vector, VR1012, resulted in only background levels of IL-2.

-73-

Table 3
mIL-2 concentration in ascites (pg/ml)

Days post tumor cell injection

	Treatment	<u>7</u>	<u>9</u>	<u>11</u>	<u>13</u>	<u>15</u>	<u>17</u>	<u>19</u>	<u>21</u>	<u>23</u>
5										
10	VR1110 (mIL-2) without DM/DP, 6 injs.	28	12	0	1	0	1	0		
	VR1012 (control) + DM/DP, 6 injs.	0	0	8	0	0	1	43		
15	VR1110 (mIL-2) + DM/DP, 6 injs.			7196	7716	7824	3895	3200		
	VR1110 (mIL-2) + DM/DP, 4 injs.		3524	4187	5968	3050	3984	2392	441	
20	VR1110 (mIL-2) + DM/DP, 2 injs.	28882	4750	11725	8047	1246	1445	1407	774	753

mIL-2 concentration in serum (pg/ml)

Days post tumor cell injection

	Treatment	<u>7</u>	<u>9</u>	<u>11</u>	<u>13</u>	<u>15</u>	<u>17</u>	<u>19</u>	<u>21</u>	<u>23</u>
30										
35	VR1110 (mIL-2) without DM/DP, 6 injs.	0	4	0	32	0	0	0		
	VR1012 (control) + DM/DP, 6 injs.	0	0	0	0	0	0	62		
40	VR1110 (mIL-2) + DM/DP, 6 injs.			66	85	34	5	4		
	VR1110 (mIL-2) + DM/DP, 4 injs.		114	81	40	10	3	2	0	
45	VR1110 (mIL-2) + DM/DP, 2 injs.	625	418	181	76	6	0	1	0	0

IL-2 in serum and ascites after single injection of pDNA:lipid injection compared to protein injection.

Five days after i.p. injection of 10^5 MOT tumor cells, C3H/HeN mice were injected with 100 μ g of either VR1110: lipid or VR1012: lipid (1:1 DNA:DMRIE mass ratio) or with 100 μ g of VR1110 without lipid. For the IL-2 protein-treated group, mice were injected with 1 μ g recombinant murine IL-2 protein (R & D Systems, Minneapolis, MN). The pDNA: lipid, pDNA alone, or recombinant protein was injected i.p. in a total volume of 1 ml saline per mouse. Five mice from each group were sacrificed beginning at 4 hours and continuing on days 1, 2, 3, 6 and 10 post DNA or protein injection. Ascites and serum were collected and analyzed using mIL-2 ELISA assay as described above.

Mice injected i.p. with IL-2 protein had peak levels of IL-2 in ascites (10 ng) at 4 hours after injection of IL-2 and a 1000-fold reduction in IL-2 one day later (0.009 ng) (FIG. 14A). In contrast, mice injected with IL-2 pDNA: lipid had peak IL-2 levels in ascites 2 days after injection (64 ng) and only a 2.6 fold reduction in IL-2 by 10 days after injection (25 ng) (FIG. 14A). These results indicate that mice receiving IL-2 pDNA: lipid have more sustained levels of IL-2 in the ascites compared to mice receiving IL-2 protein. Tumor-bearing mice injected i.p. with either VR1012: lipid or VR1110 without cationic lipid had no IL-2 in the tumor ascites. In a related study, MOT tumor-bearing mice injected i.p. with 10-fold less VR1110: lipid (10 μ g DNA) still had detectable IL-2 in the tumor ascites 11 days after the VR1110: lipid injection (data not shown).

Serum levels of IL-2 after either i.p. protein or DNA injection reflected a similar pattern as that found in tumor ascites; however, the serum IL-2 levels were markedly reduced compared to the ascites IL-2 levels. Four hours after protein injection, IL-2 in the serum was 2.4 ng/ml and negligible by one day after protein injection. Serum levels of IL-2 one day after VR1110: lipid injection was 1 ng/ml and undetectable by 6 days after the injection (FIG.

14B). These results suggest that the majority of the IL-2 after either IL-2 protein or pDNA: lipid delivery remains in the peritoneal cavity.

5 ***Intraperitoneal injection of IL-2 plasmid DNA:lipid inhibits tumor growth and enhances survival***

Six consecutive-day treatments. The plasmid DNA:lipid therapy was evaluated for the ability to reduce tumor growth and to increase survival of mice with MOT tumors. On day 5 after MOT tumor cell injection, mice were
10 injected i.p. with 100 µg of either VR1110 or VR1012, both complexed with DMRIE/DOPE. The plasmid DNA was complexed at either a 5:1 or 1:1 DNA:DMRIE mass ratio. An additional treatment group received VR1110 that was not complexed with lipid. A total volume of 1 ml DNA:lipid or DNA alone in physiological saline was injected i.p. The DNA treatments occurred
15 over 6 consecutive days, beginning on day 5 (days 5-10). MOT tumor growth was measured by weighing the mice. All treatment groups consisted of 10 mice per group.

The high IL-2 expression level in ascites was accompanied by significant antitumor effects. Mice treated with VR1110: lipid on days 5-10
20 after tumor cell injection had a significant reduction in MOT tumor growth compared to the mice treated with the VR1012: lipid ($p=0.01$) (FIG. 15A). A significant enhancement in survival was also found for mice injected i.p. with VR1110: lipid ($p=0.05$) (FIG. 15B).

Complexing the pDNA with a cationic lipid seemed necessary for the
25 antitumor effect as treatment of tumor-bearing mice with VR1110 without lipid was not effective (FIGs. 15E and 15F). Furthermore, a 1:1 DNA:cationic lipid mass ratio was found to be more effective at reducing tumor burden and increasing survival than a 5:1 DNA:cationic lipid mass ratio (FIGs. 15C and 15D).

30 *Three alternative-day treatments.* C3H/HeN mice bearing i.p. MOT tumor ascites were injected i.p. with VR1110: lipid or with VR1012: lipid (100 µg DNA) on days 5, 8 and 11 after tumor cell implant. By day 14 post

tumor cell injection, mice treated with VR1110: lipid had a significant reduction in mean weight ($p=0.001$) compared to the mice treated with the control pDNA: lipid (FIG. 16A). In addition, a significant increase in survival ($p=0.008$) was found for the VR1110: lipid-treated mice compared to the mice
5 treated with the VR1012: lipid (FIG. 16B). By day 26 post tumor cell injection, none of the mice treated with the VR1012 were still alive, while 50% of the mice treated with VR1110: lipid remained alive. By day 55 post tumor cell injection, 20% of the mice treated with VR1110: lipid appeared to be tumor-free.

10 Whether the IL-2 pDNA: lipid antitumor effect required T cells was investigated by implanting nude mice with i.p. MOT tumors followed by same VR1110: lipid regimen used in the C3H/HeN tumor-bearing mice (DNA treatment on days 5, 8 and 11 after tumor cell implant). No significant antitumor effect was found for the nude mice treated with VR1110: lipid
15 suggesting that T cells may be required for the antitumor effect (data not shown).

IL-2 plasmid DNA:lipid antitumor effect enhanced by debulking of tumor ascites

20 Debulking of tumor ascites is commonly performed on human ovarian cancer patients. A similar procedure was performed in the mice bearing MOT tumors and treated with VR1110: lipid. Mice bearing MOT tumors and injected i.p. with 100 μ l DNA:lipid on days 5-10 as described above (1:1
25 DNA:lipid mass ratio), were also debulked of tumor ascites on day 14 after tumor cell injection (4 days after the last DNA:lipid injection). Mice were debulked of 5 ml of tumor ascites by insertion of a 22 G needle attached to a 5 ml syringe and removal of 5 ml of fluid. The mice were anesthetized with methoxyflurane during the debulking procedure. All treatment groups in this
30 experiment consisted of 8-10 mice per group.

Debulking of ovarian tumor ascites in mice previously treated with IL-2 plasmid DNA:lipid further enhanced the efficacy of the treatment resulting

-77-

in a significant reduction in tumor growth ($p=0.01$) and an increase in survival. Forty four percent of the IL-2 plasmid DNA:lipid-treated and debulked mice were alive on day 57 vs. 17% of the plasmid control-treated and debulked mice (FIG. 17). These results show that plasmid-mediated gene therapy in combination with conventional procedures such as debulking of tumor ascites may hold promise for future treatment of human ovarian cancer.

Dose-response of IL-2 pDNA: lipid

A dose-response study was initiated to determine the minimum dose of VR1110: lipid that could still result in a significant antitumor effect. C3H/HeN mice were injected with 25, 50 or 100 μ g of IL-2 pDNA: lipid on days 5, 8 and 11 after MOT tumor cell injection. A control group of MOT tumor-bearing mice were injected with 100 μ g of VR1012 complexed with lipid. By day 15 post tumor cell injection, mice treated with either the 50 or 100 μ g dose of VR1110 complexed with lipid had a significant inhibition of tumor growth ($p=0.002$) compared to the mice treated with the VR1012: lipid (FIG. 18A). A significant increase in survival ($p=0.01$) was also found for the mice treated with either the 50 or 100 μ g dose of VR1110: lipid (FIG. 18B). On day 25, none of the mice treated with the VR1012: lipid survived, while the mice injected with 50 or 100 μ g of VR1110: lipid had 27 and 33% survival, respectively. By day 37, mice treated with 50 or 100 μ g of VR1110: lipid had 20 and 27% survival, respectively. Tumor-bearing mice treated with 25 μ g of VR1110 complexed with lipid were not significantly different from the control mice for either tumor volume or survival.

Cytokine profile of ovarian tumor ascites

Since i.p. injection of IL-2 pDNA: lipid into mice bearing i.p. MOT tumors resulted in high levels of IL-2 expression in the ascites, it was of interest to determine whether the IL-2 therapy initiated a cytokine cascade in

the tumor ascites. C3H/HeN mice were injected i.p. with 10^5 MOT cells. On days 5, 8 and 11 after tumor cell implant, the mice were injected i.p. with either VR1110: lipid or VR1012: lipid (1:1 pDNA:DMRIE mass ratio) or received no treatment after the MOT tumor cell injection. Two days after each injection of pDNA: lipid (days 7, 10 and 13 after tumor cell implant), 5 mice per group were sacrificed and the tumor ascites was collected. The total volume of ascites was determined per mouse. The ascites samples were spun at 14,000 rpm for 2 minutes followed by collection of the supernatants. The ascites supernatants were assayed for the concentration of the cytokines: IL-2, IL-4, IL-6, IL-10, IL-12, granulocyte-macrophage colony stimulating factor (GM-CSF), interferon gamma ($\text{IFN}\gamma$) and tumor necrosis factor alpha ($\text{TNF}\alpha$) using ELISA (R & D Systems, Minneapolis, MN). The concentration of transforming growth factor beta ($\text{TGF}\beta$) in the ascites was assayed using the $\text{TGF}\beta_2$ Emax Immunoassay System (Promega, Madison, WI). The amount of cytokine in the tumor ascites was calculated using the formula: cytokine concentration in pg/ml x ml of total ascites = pg of cytokine/total ascites.

As expected, tumor-bearing mice injected i.p. with VR1110: lipid had a marked increase in IL-2 levels with negligible levels in untreated tumor-bearing mice or mice injected with the VR1012: lipid (FIG. 19A). The levels of $\text{IFN}\gamma$ and GM-CSF were also markedly elevated in the mice treated with VR1110: lipid (FIGs. 19B and 19C). The $\text{IFN}\gamma$ and GM-CSF levels in these mice increased on days 10 and 13 after tumor cell injection but not on the day 7 timepoint suggesting that this could be due to IL-2 secretion. Some non-specific increase in $\text{IFN}\gamma$ was also found in the ascites of tumor-bearing mice after injection of the VR1012: lipid; however, by day 13, the levels of $\text{IFN}\gamma$ in the tumor-bearing mice treated with VR1110: lipid were 6-fold higher than in the mice treated with VR1012: lipid suggesting that expression of IL-2 upregulates $\text{IFN}\gamma$ production.

Levels of IL-6, $\text{TNF}\alpha$ and IL-10 were increased in both the IL-2 pDNA: lipid group as well as the control pDNA: lipid group suggesting that pDNA: lipid complexes may non-specifically stimulate production of these

particular cytokines in the tumor ascites (FIGs. 19D, 19E, and 19F). No differences were found for IL-4, IL-12 or TGF β in the ascites from any of the groups and levels of these cytokines were low (0-300 pg/ml for IL-4 and IL-12 and 0-2000 pg/ml for TGF β , data not shown). For all of the cytokines evaluated, mice treated with control pDNA without lipid, IL-2 pDNA without lipid or with lipid alone had similar cytokine levels as the untreated mice.

Intraperitoneal injection of IFN α pDNA:lipid enhances survival

C3H/HeN mice were injected i.p. with 10^5 MOT cells to establish ovarian i.p. tumors. The mice then received i.p. injections of 100 μ g of VR4111 (mIFN α) or VR1012 (control) complexed with DMRIE:DOPE cationic lipid at a 1:1 DNA:DMRIE mass ratio in a total volume of 1 ml saline. The mice received the i.p. injections of pDNA: lipid on days 5, 8 and 11 after tumor cell injection. The mice were weighed 3-6 times per week. Fifteen mice were included in each treatment group.

Mice bearing i.p. MOT tumors and treated with i.p. VR4111: lipid had a significant increase in survival ($p < 0.006$) compared to the mice receiving the control plasmid (FIG. 20B). No significant reduction in tumor volume was found for the VR4111: lipid-treated mice (FIG. 20A).

Example 7

Selective Transfection of Malignant Cells in Murine Intraperitoneal Melanoma Tumor Model

The anti-tumor effect of DNA formulations with or without lipids in mouse i.p. melanoma model have been evaluated in the present example.

Cell line and tumor model

B16F10 mouse melanoma cells were grown *in vitro* in DMEM and 10% FCS. Two hundred thousand B16F10 mouse melanoma cells were
5 implanted i.p. in C57BL/6 mice in 1-3 ml saline using a 28 G 1/2 needle and without puncturing internal organs.

Preparation of plasmid DNA:lipid complexes

10 The plasmid DNA-cationic lipid formulations were prepared just prior to use. Equal volumes of DNA and DMRIE : DOPE (1:1) were mixed by swirling to achieve a target concentration of 0.5 mg DNA/ml, 100 µg DMRIE/ml and 0.12 mg DOPE/ml as previously described (Parker et al, 1996; Saffran et al, 1998). The other cationic lipids were mixed similarly so that the
15 mass ratio of DNA to cationic lipid was 5:1 and the cationic lipid to DOPE molar ratio was 1:1. The formulated material was then vortexed at high speed for 30 seconds and kept at room temperature until dose administration.

CAT assay

20

Tumor or other tissues were collected, immediately frozen in liquid nitrogen and ground into a powder using a reversible drill as described (Manthorpe, M., Hartikka, J., Vahlsing, H.L. and Sawdey, M. Quantification of plasmid DNA transfection *in vivo*. In Gene Quantification. F. Ferre, ed.
25 Birkhauser, Boston, Ma., 1998 in press.). The dry frozen powder was thawed and extracted in lysis buffer and high speed supernates assayed for CAT activity using a two-phase partition assay as described (Sankaran, L., Analytical Biochemistry 200: 180-186 (1992)).

Cationic lipids enhances tumor transfection

C57BL/6 mice were injected i.p. with 200,000 B16F10 murine melanoma cells, and seven days later, injected i.p. with CAT pDNA (VR1332):DMRIE/DOPE. Two days later, tumor tissues were collected, extracted, and assayed for CAT activity.

As shown in Table 4, DNA alone transfected tumor, but DMRIE:DOPE increased transfection by 78 fold (from 2,326 to 182,052).

Table 4

pgs CAT per gm of tumor tissue collected
(n = 4 to 6 mice as indicated); saline values subtracted

Mouse #	DNA only	DNA:DMRIE	DMRIE, no DNA
1	0	82,600	0
2	307	137,551	0
3	706	220,600	0
4	791	287,458	0
5	5,892		0
6	6,258		
Average	2,326	182,052	0
Std Error	1,306	52,140	0
Fold higher	1	78	0

10 *Selective transfection of tumor cells*

Normal animal. Normal, non-tumor bearing BALB/c mice were injected i.p. with 1mg/2ml VR1332, with or without a cationic lipid, and with or without the neutral lipid, DOPE. Two days later, selected i.p. tissues (liver, lung, kidney, spleen, mesentery) were collected, extracted, and assayed for CAT activity.

As shown in Table 5, normal intraperitoneal organs were not at all transfected or transfected very little with a variety of cationic lipids. There was a low level transfection of mesentery tissues, and a moderate transfection of 1 of 5 spleens (the one transfected spleen may have been punctured by the injection needle).

Table 5

Average pgs CAT per gm of tumor tissue (avg., n = 5 mice)

Saline background values have been subtracted

ORGANS	DNA only	+ DMRIE	+ β AE-DMRIE	+ GAP-DLRIE
Liver	0	0	0	0
Lung	0	0	0	0
Kidney	0	0	0	0
Spleen	0	2,563(0)*	0	0
Mesentery	370	1,549	1,866	1,127

ORGANS	+ DOSPA	+ DMRIE, no DOPE	+ β AE-DMRIE, no DOPE
Liver	0	0	0
Lung	0	0	0
Kidney	0	0	0
Spleen	0	0	0
Mesentery	600	1,091	3,604

*one mouse with a CAT value; the rest = 0

Tumor-bearing animal compared to normal animal. C57BL/6 mice were injected i.p. with 200,000 B16F10 murine melanoma cells, and seven days later, injected i.p. with VR1332 with or without cationic lipid/DOPE. Two days later, selected i.p. tissues (liver, lung, kidney, spleen, mesentery) were collected, extracted, and assayed for CAT activity.

As shown in Table 6, tumor tissues were transfected at high levels with pDNA complexed with cationic lipid/DOPE. Normal i.p. organs were not transfected well compared to i.p. tumor tissues.

Table 6

Average pgs CAT per gm of tumor tissue (n = 5 mice); saline values subtracted

ORGANS	DNA only	DNA + DMRIE:DOPE	DNA + DMRIE:DOPE normal mice no tumors
Tumor	0	200,195	n/a
Liver	0	0	0
Lung	0	0	0
Kidney	0	0	0
Spleen	493	0	0
Mesentery	6,795	nd	2,562
Ovary	nd	nd	0

5

Dose Response

C57BL/6 mice were injected i.p. with 200,000 B16F10 murine melanoma cells, and seven days later, injected i.p. with 0.15 or 1.5 mgs of VR1332 with or without cationic lipid/DOPE in 3 ml saline. Two days later, tumor tissues were collected, extracted, and assayed for CAT activity.

10

As shown in Table 7, a higher dose of DNA transfected tumor tissues better than a lower dose. Also, DMRIE transfected better than the two other cationic lipids tested.

Table 7

Average pgs CAT per gm of tumor tissue (n = 5 mice); saline values subtracted

DNA DOSE	DNA only	+ DMRIE	+ GAP-DMRIE	+ PA-DELO
1.5 mgs	16,592	1,270,466	524,006	581,616
0.15 mgs	1,937	131,089	47,866	99,797

5 ***Testing a variety of cationic lipids***

C57BL/6 mice were injected i.p. with 200,000 B16F10 murine melanoma cells, and seven days later, injected i.p. with 1 mg of VR1332 with or without cationic lipid/DOPE in 3 ml saline. Two days later, tumor tissues
10 were collected, extracted, and assayed for CAT activity.

As shown in Table 8, all cationic lipids tested increased transfection level, and DMRIE was one of three preferred cationic lipids.

Table 8

Average pgs CAT per gm of tumor tissue (n = 5 mice);
saline values have been subtracted

CL:DOPE	Run 1	Run 2	Average
None	6,247	not done	6,247
GAP-DDRIE	20,816	7,027	13,922
GMU-DMRIE	9,355	52,888	31,122
HP-DORIE	36,615	26,795	31,705
DOSPA	33,494	57,026	45,260
PA-TELO	63,326	32,681	48,004
GA-LOE-BP	59,729	63,528	61,629
GAP-DMRIE	80,760	63,002	71,881
PA-DELO	73,692	82,265	77,979
GAP-DLRIE	77,553	107,629	92,591
DMRIE	77,128	122,225	99,677
DLRIE	122,999	128,225	125,612
PA-DEMO	155,369	154,884	155,127

In sum, the success of the intra-cavity delivery embodiment of the present invention is also exemplified in the murine melanoma tumor model. Following i.p. injection of a polynucleotide, transfection occurs predominantly in tumor tissues, and normal intraperitoneal organs, such as liver, lung, and kidney are poorly transfected, if at all, with the polynucleotide formulation.

It clear that the invention may be practiced otherwist than as particular described in the foregoing description and examples.

Numerous modifications and variations of the present invention are possible in light of the above teachings and, therefore, are withing the scope of the appended claims.

The entire disclosure of all publications (including patents, patent application, journal articles, laboratory manuals, books, or other documents) cited herein are hereby incorporated by reference.

What Is Claimed Is:

1. A pharmaceutical composition comprising about 1 ng to 20 mg of a non-infectious, non-integrating polynucleotide construct comprising a
5 polynucleotide selected from the group consisting of:

(a) a polynucleotide that hybridizes under stringent conditions to the nucleotide sequence of SEQ ID No. 7 or the complement thereof,

wherein said polynucleotide sequence encodes a polypeptide that has antiproliferative activity when added to NIH-OVCAR3 cells *in vitro*;

10 (b) a polynucleotide that encodes a polypeptide comprising an amino acid sequence which, except for at least one but not more than 20 amino acid substitutions, deletions, or insertions, is identical to amino acids -23 to 172 or 1 to 172 in SEQ ID No. 8, wherein said polypeptide has antiproliferative activity when added to NIH-OVCAR3 cells *in vitro*; and

15 (c) a polynucleotide that encodes a polypeptide comprising amino acids 86 to 172 in SEQ ID No. 8, wherein said polypeptide has antiproliferative activity when added to NIH-OVCAR3 cells *in vitro*; and

one or more cationic compounds selected from the group consisting of cationic lipids, cationic peptides, cationic proteins, cationic polymers, and
20 mixtures thereof.

2. The pharmaceutical composition of claim 1, wherein said polynucleotide is (a).

25 3. The pharmaceutical composition of claim 1, wherein said polynucleotide is (b).

4. The pharmaceutical composition of claim 1, wherein said polynucleotide is (c).

30

-88-

5 5. The pharmaceutical composition of claim 2, wherein said polynucleotide encodes a polypeptide comprising amino acids -23 to 172 of SEQ ID No. 8.

10 6. The pharmaceutical composition of claim 2, wherein said polynucleotide encodes a polypeptide comprising amino acids 1 to 172 of SEQ ID No. 8.

15 7. The pharmaceutical composition of claim 3, wherein said polynucleotide encodes a polypeptide comprising an amino acid sequence which, except for at least one but not more than 15 amino acid substitutions, deletions, or insertions, is identical to amino acids -23 to 172 or 1 to 172 in SEQ ID No. 8.

20 8. The pharmaceutical composition of claim 7, wherein said polynucleotide encodes a polypeptide comprising an amino acid sequence which, except for at least one but not more than 10 amino acid substitutions, deletions, or insertions, is identical to amino acids -23 to 172 or 1 to 172 in SEQ ID No. 8.

25 9. The pharmaceutical composition of claim 8, wherein said polynucleotide encodes a polypeptide comprising an amino acid sequence which, except for at least one but not more than 5 amino acid substitutions, deletions, or insertions, is identical to amino acids -23 to 172 or 1 to 172 in SEQ ID No. 8.

 10. The pharmaceutical composition of claim 1, wherein said polynucleotide further encodes one or more additional molecules.

30 11. The pharmaceutical composition of claim 1, wherein said polynucleotide further encodes one or more additional cytokines.

12. The pharmaceutical composition of claim 4, wherein said polypeptide further comprises amino acids 61 to 85 of SEQ ID No. 8.

13. The pharmaceutical composition of claim 12, wherein said
5 polypeptide further comprises amino acids 41 to 60 of SEQ ID No. 8.

14. The pharmaceutical composition of claim 13, wherein said polypeptide further comprises amino acids 21 to 40 of SEQ ID No. 8.

10 15. The pharmaceutical composition of claim 1, wherein said cationic compound is one or more cationic lipids.

16. The pharmaceutical composition of claim 15, wherein said cationic lipids are selected from the group consisting of (±)-N-(2-
15 hydroxyethyl)-N,N-dimethyl-2,3-bis(tetradecyloxy)-1-propaniminium bromide, (±)-N-(2-aminoethyl)-N,N-dimethyl-2,3-bis(tetradecyloxy)-1-propaniminium bromide, (±)-N-(3-aminopropyl)-N,N-dimethyl-2,3-bis(dodecyloxy)-1-propaniminium bromide, 5-carboxyspermylglycine dioctadecylamide, dipalmitoyl-phosphatidylethanolamine 5-carboxy-
20 spennylamide, {3β-[N,N',N'-dimethylamino]ethane}-carbomoyl}-cholesterol, dimethyldioctdecyl-ammonium bromide, (±)-N,N-dimethyl-N-[2-(sperminecarboxamido)ethyl]-2,3-bis(dioleoyloxy)-1-propaniminium pentahydrochloride, DL-1,2-dioleoyl-3-dimethylaminopropyl-β-hydroxyethylammonium, 1-O-oleyl-2-oleoyl-3-dimethylaminopropyl-β-hydroxyethylammonium, cationic lipids are 3,5-(NN-dilysyl)-diaminobenzoyl-
25 3-(DL-1,2-dioleoyl-dimethylaminopropyl-β-hydroxyethylamine) and 3,5-(N,N-di-lysyl)diamino-benzoylglycyl-3-(DL-1,2-dioleoyl-dimethylaminopropyl-β-hydroxyethylamine) and mixtures thereof.

30 17. The pharmaceutical composition of claim 15, wherein said cationic lipids comprise (±)-N-(2-hydroxyethyl)-N,N-dimethyl-2,3-bis(tetradecyloxy)-1-propaniminium bromide.

18. The pharmaceutical composition of claim 15, further comprising one or more neutral lipids.

5 19. The pharmaceutical composition of claim 18, wherein said neutral lipids comprise 1,2-dioleoyl-glycero-3-phosphoethanolamine.

20. The pharmaceutical composition of claim 18, wherein said polynucleotide construct is complexed with (\pm)-N-(2-hydroxyethyl)-N,N-
10 dimethyl-2,3-bis(tetradecyloxy)-1-propaniminium bromide and 1,2-dioleoyl-glycero-3-phosphoethanolamine.

21. The pharmaceutical composition of claim 18, wherein the mass ratio of said polynucleotide construct to lipid is from about 5:1 to about 1:1.

15

22. The pharmaceutical composition of claim 1, wherein the polynucleotide encoding the polypeptide is DNA operably linked to a promoter.

20

23. The pharmaceutical composition of claim 1, wherein the polynucleotide encoding the polypeptide is RNA.

24. The pharmaceutical composition of claim 1, wherein said polynucleotide construct is VR4151 complexed with (\pm)-N-(2-hydroxyethyl)-
25 N,N-dimethyl-2,3-bis(tetradecyloxy)-1-propaniminium bromide and 1,2-dioleoyl-glycero-3-phosphoethanolamine such that the mass ratio of VR4151 (SEQ ID No. 4) to lipid is from about 5:1 to about 1:1.

25. The pharmaceutical composition of claim 1, wherein said
30 polynucleotide comprises a region regulating gene expression.

26. The pharmaceutical composition of claim 25, wherein said region regulating gene expression is cell specific or tissue specific.

27. The pharmaceutical composition of claim 26, wherein said
5 region is tumor cell or tumor tissue specific.

28. A pharmaceutical composition obtained by complexing about 1 ng to 20 mg of a non-infectious, non-integrating polynucleotide construct comprising a polynucleotide selected from the group consisting of:

10 (a) a polynucleotide that hybridizes under stringent conditions to the nucleotide sequence of SEQ ID No. 7 or the complement thereof,

wherein said polynucleotide sequence encodes a polypeptide that has antiproliferative activity when added to NIH-OVCAR3 cells *in vitro*;

(b) a polynucleotide that encodes a polypeptide comprising an
15 amino acid sequence which, except for at least one but not more than 20 amino acid substitutions, deletions, or insertions, is identical to amino acids -23 to 172 or 1 to 172 in SEQ ID No. 8, wherein said polypeptide has antiproliferative activity when added to NIH-OVCAR3 cells *in vitro*; and

(c) a polynucleotide that encodes a polypeptide comprising amino
20 acids 86-172 of SEQ ID No. 8, and wherein said polypeptide has antiproliferative activity when added to NIH-OVCAR3 cells *in vitro*;

with one or more cationic compounds selected from the group consisting of cationic lipids, cationic peptides, cationic proteins, cationic
25 polymers, and mixtures thereof.

29. The pharmaceutical composition of claim 28, wherein said polynucleotide is (a).

30. The pharmaceutical composition of claim 28, wherein said
30 polynucleotide is (b).

-92-

31. The pharmaceutical composition of claim 28, wherein said polynucleotide is (c).

32. The pharmaceutical composition of claim 29, wherein said
5 polynucleotide encodes a polypeptide comprising amino acids -23 to 172 of
SEQ ID No. 8.

33. The pharmaceutical composition of claim 29, wherein said
polynucleotide encodes a polypeptide comprising amino acids 1 to 172 of
10 SEQ ID No. 8.

34. The pharmaceutical composition of claim 30, wherein said
polynucleotide encodes a polypeptide comprising an amino acid sequence
which, except for at least one but not more than 15 amino acid substitutions,
15 deletions, or insertions, is identical to amino acids -23 to 172 or 1 to 172 in
SEQ ID No. 8.

35. The pharmaceutical composition of claim 34, wherein said
polynucleotide encodes a polypeptide comprising an amino acid sequence
20 which, except for at least one but not more than 10 amino acid substitutions,
deletions, or insertions, is identical to amino acids -23 to 172 or 1 to 172 in
SEQ ID No. 8.

36. The pharmaceutical composition of claim 35, wherein said
25 polynucleotide encodes a polypeptide comprising an amino acid sequence
which, except for at least one but not more than 5 amino acid substitutions,
deletions, or insertions, is identical to amino acids -23 to 172 or 1 to 172 in
SEQ ID No. 8.

30 37. The pharmaceutical composition of claim 28, wherein said
polynucleotide further encodes one or more additional molecules.

38. The pharmaceutical composition of claim 28, wherein said polynucleotide further encodes one or more additional cytokines.

39. The pharmaceutical composition of claim 31, wherein said polypeptide further comprises amino acids 61 to 85 of SEQ ID No. 8.

40. The pharmaceutical composition of claim 39, wherein said polypeptide further comprises amino acids 41 to 60 of SEQ ID No. 8.

41. The pharmaceutical composition of claim 40, wherein said polypeptide further comprises amino acids 21 to 40 of SEQ ID No. 8.

42. The pharmaceutical composition of claim 23, wherein said cationic compound is one or more cationic lipids.

43. The pharmaceutical composition of claim 42, wherein said cationic lipids are selected from the group consisting of (\pm)-N-(2-hydroxyethyl)-N,N-dimethyl-2,3-bis(tetradecyloxy)-1-propaniminium bromide, (\pm)-N-(2-aminoethyl)-N,N-dimethyl-2,3-bis(tetradecyloxy)-1-propaniminium bromide, (\pm)-N-(3-aminopropyl)-N,N-dimethyl-2,3-bis(dodecyloxy)-1-propaniminium bromide, 5-carboxyspermylglycine dioctadecylamide, dipalmitoyl-phosphatidylethanolamine 5-carboxy-spermylamide, {3 β -[N,N',N'-dimethylamino]ethane]-carbomoyl}-cholesterol, dimethyldioctadecyl-ammonium bromide, (\pm)-N,N-dimethyl-N-[2-(sperminecarboxamido)ethyl]-2,3-bis(dioleoyloxy)-1-propaniminium pentahydrochloride, DL-1,2-dioleoyl-3-dimethylaminopropyl- β -hydroxyethylammonium, 1-O-oleyl-2-oleoyl-3-dimethylaminopropyl- β -hydroxyethylammonium, cationic lipids are 3,5-(NN-dilysyl)-diaminobenzoyl-3-(DL-1,2-dioleoyl-dimethylaminopropyl- β -hydroxyethylamine) and 3,5-(N,N-di-lysyl)diamino-benzoylglycyl-3-(DL-1,2-dioleoyl-dimethylaminopropyl- β -hydroxyethylamine) and mixtures thereof.

44. The pharmaceutical composition of claim 42, wherein said cationic lipids comprise (\pm) -N-(2-hydroxyethyl)-N,N-dimethyl-2,3-bis(tetradecyloxy)-1-propaniminium bromide.

5 45. The pharmaceutical composition of claim 42, further comprising one or more neutral lipids.

46. The pharmaceutical composition of claim 45, wherein said neutral lipids comprise 1,2-dioleoyl-glycero-3-phosphoethanolamine.

10

47. The pharmaceutical composition of claim 45, wherein said polynucleotide construct is complexed with (\pm) -N-(2-hydroxyethyl)-N,N-dimethyl-2,3-bis(tetradecyloxy)-1-propaniminium bromide and 1,2-dioleoyl-glycero-3-phosphoethanolamine.

15

48. The pharmaceutical composition of claim 45, wherein the mass ratio of said polynucleotide construct to lipid is from about 5:1 to about 1:1.

49. The pharmaceutical composition of claim 28, wherein the polynucleotide encoding the polypeptide is DNA operably linked to a promoter.

20

50. The pharmaceutical composition of claim 28, wherein the polynucleotide encoding the polypeptide is RNA.

25

51. The pharmaceutical composition of claim 28, wherein said polynucleotide construct is VR4151 complexed with (\pm) -N-(2-hydroxyethyl)-N,N-dimethyl-2,3-bis(tetradecyloxy)-1-propaniminium bromide and 1,2-dioleoyl-glycero-3-phosphoethanolamine such that the mass ratio of VR4151 (SEQ ID No. 4) to lipid is from about 5:1 to about 1:1.

30

52. The pharmaceutical composition of claim 28, wherein said polynucleotide comprises a region regulating gene expression.

53. The pharmaceutical composition of claim 52, wherein said
5 region regulating gene expression is cell specific or tissue specific.

54. The pharmaceutical composition of claim 53, wherein said region is tumor cell or tumor tissue specific.

10 55. A method of treating cancer or metastasis thereof in a mammal, comprising:
administering into a tissue of said mammal a non-infectious, non-integrating polynucleotide construct comprising a polynucleotide encoding a cytokine, or an active fragment thereof,
15 such that said polynucleotide is expressed as said cytokine *in vivo*, and
such that said cytokine is present in the blood stream of said mammal
in an amount effective to treat said cancer.

56. The method of claim 55, wherein said polynucleotide construct
20 is administered free from ex vivo cells or ex vivo cellular material.

57. The method of claim 55, wherein said cancer is selected from the group consisting of renal cell carcinoma, colorectal carcinoma, lymphoma, Kaposi's sarcoma, melanoma, prostate cancer, ovarian cancer, lung cancer,
25 liver cancer, head and neck cancer, bladder cancer, uterine cancer, bone cancer, leukemia, breast cancer, non-melanoma skin cancer, glioma, solid cutaneous tumor, metastases of any of thereof, and combinations of any of thereof.

30 58. The method of claim 57, wherein said cancer is one or more metastases of one or more of the cancers of claim 57.

59. The method of claim 57, wherein said cancer is lung or liver metastasis.

60. The method of claim 55, wherein said construct is administered
5 into a tissue selected from the group consisting of muscle, skin, brain, lung, liver, spleen, bone marrow, thymus, heart, lymph, blood, bone, cartilage, pancreas, kidney, gall bladder, stomach, intestine, testis, ovary, uterus, rectum, nervous system, eye, gland, and connective tissue.

10 61. The method of claim 60, wherein said tissue is muscle.

62. The method of claim 61, wherein said tissue is skeletal muscle, smooth muscle, or myocardium.

15 63. The method of claim 55, wherein said construct is administered intravenously.

64. The method of claim 55, wherein said construct is administered intramuscularly.

20 65. The method of claim 55, wherein said polynucleotide construct comprises a polynucleotide encoding a cytokine selected from the group consisting of IFN ω , IFN α , IFN τ , IFN γ , IFN β , IL-1, IL-2, IL-4, IL-7, IL-12, IL-15, IL-18, GM-CSF, an active fragment of any of thereof, and a
25 combination of any of thereof.

66. The method of claim 65, wherein said cytokine is an interferon ω .

30 67. The method of claim 66, wherein said interferon ω is a polypeptide comprising amino acids -23 to 172 in SEQ ID No. 8.

-97-

68. The method of claim 66, wherein said interferon ω is a polypeptide comprising amino acids 1 to 172 in SEQ ID No. 8.

69. The method of claim 66, wherein said polynucleotide construct
5 is VR4151 (SEQ ID No. 4).

70. The method of claim 65, wherein said cytokine is an interferon α .

71. The method of claim 70, wherein said interferon α is a
10 polypeptide comprising amino acids -23 to 166 of SEQ ID No. 10.

72. The method of claim 70, wherein said interferon α is a polypeptide comprising amino acids 1 to 166 of SEQ ID No. 10.

73. The method of claim 70, wherein said polynucleotide construct
15 is VR4112 (SEQ ID No. 2).

74. The method of claim 65, wherein said cytokine is an interleukin-2.
20

75. The method of claim 74, wherein interleukin-2 is a polypeptide comprising amino acids 58 to 105 of SEQ ID No. 14.

76. The method of claim 74, wherein interleukin-2 is a polypeptide
25 comprising amino acids 20 to 126 of SEQ ID No. 14.

77. The method of claim 74, wherein said polynucleotide construct is VR1103 (SEQ ID No. 25).

78. The method of claim 55, wherein a polynucleotide construct
30 comprising a DNA molecule encoding an interferon ω is administered intramuscularly to treat melanoma or metastasis thereof.

79. The method of claim 55, wherein a polynucleotide construct comprising a DNA molecule encoding an interferon ω is administered intramuscularly to treat metastasis of melanoma.

5

80. The method of claim 79, wherein the metastasis of melanoma is lung metastasis.

81. The method of claim 55, wherein a polynucleotide construct comprising a DNA molecule encoding an interferon ω is administered to treat glioma.

82. The method of claim 55, wherein a polynucleotide construct comprising a DNA molecule encoding an interferon α is administered intramuscularly to treat melanoma.

83. The method of claim 55, wherein a polynucleotide construct comprising a DNA molecule encoding an interferon α is administered intramuscularly to treat metastasis of melanoma.

20

84. The method of claim 83, wherein the metastasis of melanoma is lung metastasis.

85. The method of claim 55, wherein a polynucleotide construct comprising a DNA molecule encoding an interferon α is administered to treat glioma.

25

86. The method of claim 55 wherein said mammal is a human.

30

87. The method of claim 55 wherein said construct is free from association with transfection-facilitating proteins, viral particles, liposomes, cationic lipids, and calcium phosphate precipitating agents.

88. The method of claim 55, wherein said construct is administered as a complex of said construct and one or more cationic compounds selected from the group consisting of cationic lipids, cationic peptides, cationic proteins, cationic polymers, and mixtures thereof.

89. The method of claim 88, wherein said one or more cationic compounds are one or more cationic lipids.

90. The method of claim 89, wherein said cationic lipids are selected from the group consisting of (\pm)-N-(2-hydroxyethyl)-N,N-dimethyl-2,3-bis(tetradecyloxy)-1-propaniminium bromide, (\pm)-N-(2-aminoethyl)-N,N-dimethyl-2,3-bis(tetradecyloxy)-1-propaniminium bromide, (\pm)-N-(3-aminopropyl)-N,N-dimethyl-2,3-bis(dodecyloxy)-1-propaniminium bromide, 5-carboxyspermylglycine dioctadecylamide, dipalmitoyl-phosphatidylethanolamine 5-carboxy-spermylamide, $\{3\beta$ -[N,N'-dimethylamino)ethane]-carbomoyl}-cholesterol, dimethyldioctdecylammonium bromide, (\pm)-N,N-dimethyl-N-[2-(sperminecarboxamido)ethyl]-2,3-bis(dioleyloxy)-1-propaniminium pentahydrochloride, DL-1,2-dioleoyl-3-dimethylaminopropyl- β -hydroxyethylammonium, 1-O-oleyl-2-oleoyl-3-dimethylaminopropyl- β -hydroxyethylammonium, cationic lipids are 3,5-(NN-dilysyl)-diaminobenzoyl-3-(DL-1,2-dioleoyl-dimethylaminopropyl- β -hydroxyethylamine) and 3,5-(N,N-di-lysyl)diamino-benzoylglycyl-3-(DL-1,2-dioleoyl-dimethylaminopropyl- β -hydroxyethylamine) and mixtures thereof.

91. The method of claim 89, wherein said cationic lipids comprise (\pm)-N-(2-hydroxyethyl)-N,N-dimethyl-2,3-bis(tetradecyloxy)-1-propaniminium bromide.

92. The method of claim 89, wherein said complex further comprising one or more neutral lipids.

-100-

93. The method of claim 89, wherein said neutral lipids comprise 1,2-dioleoyl-glycero-3-phosphoethanolamine.

94. The method of claim 94, wherein said polynucleotide construct
5 is complexed with (\pm)-N-(2-hydroxyethyl)-N,N-dimethyl-2,3-bis(tetradecyloxy)-1-propaniminium bromide and 1,2-dioleoyl-glycero-3-phosphoethanolamine.

95. The method of claim 94, wherein the ratio of said
10 polynucleotide construct to lipid is about 5:1 to about 1:1.

96. The method of claim 55, wherein said polynucleotide construct is VR4151 (SEQ ID No. 4) complexed with (\pm)-N-(2-hydroxyethyl)-N,N-dimethyl-2,3-bis(tetradecyloxy)-1-propaniminium bromide and 1,2-dioleoyl-glycero-3-phosphoethanolamine such that the ratio of VR4151 to lipid is about
15 5:1 to 1:1.

97. The method of claim 55, wherein said polynucleotide construct is VR4112 (SEQ ID No. 4) complexed with (\pm)-N-(2-hydroxyethyl)-N,N-dimethyl-2,3-bis(tetradecyloxy)-1-propaniminium bromide and 1,2-dioleoyl-glycero-3-phosphoethanolamine such that the mass ratio of VR4112 (SEQ ID
20 No. 4) to lipid is from about 5:1 to about 1:1.

98. The method of claim 55, wherein said polynucleotide construct
25 is VR1103 (SEQ ID No. 25) complexed with (\pm)-N-(2-hydroxyethyl)-N,N-dimethyl-2,3-bis(tetradecyloxy)-1-propaniminium bromide and 1,2-dioleoyl-glycero-3-phosphoethanolamine such that the mass ratio of VR1103 (SEQ ID No. 25) to lipid is from about 5:1 to about 1:1.

30 99. The method of claim 55, wherein said polynucleotide is DNA operably linked to a promoter.

-101-

100. The method of claim 55, wherein said polynucleotide is RNA.

101. The method of claim 55, wherein said polynucleotide construct further comprises a polynucleotide encoding one or more additional molecule.

5

102. The method of claim 55, wherein said polynucleotide construct further comprises a polynucleotide encoding one or more additional cytokines.

10

103. The method of claim 55, wherein said polynucleotide comprises a region regulating gene expression.

104. The method of claim 103, wherein said region regulating gene expression is cell specific or tissue specific.

15

105. The method of claim 104, wherein said region is tumor cell or tumor tissue specific.

106. A method of treating cancer, or metastasis thereof, in a mammal, comprising:

20

the method of claim 55 in combination with one or more additional cancer treatment methods selected from the group consisting of surgery, radiation therapy, chemotherapy, immunotherapy, and gene therapy.

25

107. The method of claim 106, wherein said polynucleotide construct is administered prior to the commencement of said one or more additional cancer treatment methods.

30

108. The method of claim 106, wherein said polynucleotide construct is administered during the practice of said at one or more additional cancer treatment methods.

-102-

109. The method of claim 106, wherein said polynucleotide construct is administered after the end of said one or more additional cancer treatment methods.

5 110. A method of treating cancer in a mammal, comprising:
administering into a tissue of said mammal a non-infectious, non-integrating polynucleotide construct comprising a polynucleotide encoding a cytokine, or an active fragment thereof, selected from the group consisting of an interferon ω , an interferon α , a combination of an interferon ω and an
10 interferon α , and an active fragment of any of thereof, such that said cytokine is delivered to a tumor in a therapeutically effective amount.

111. The method of claim 110, wherein said polynucleotide construct is administered free from *ex vivo* cells or *ex vivo* cellular material.

15

112. The method of claim 110, wherein said cancer is selected from the group consisting of solid tumor, cutaneous tumor, renal cell carcinoma, colorectal carcinoma, lymphoma, Kaposi's sarcoma, melanoma, prostate cancer, ovarian cancer, lung cancer, head and neck cancer, bladder cancer,
20 uterine cancer, bone cancer, leukemia, breast cancer, non-melanoma skin cancer, glioma, metastases thereof, and combinations thereof.

113. The method of claim 110, wherein said polynucleotide construct is administered directly into a tissue selected from the group
25 consisting of muscle, skin, breast, prostate, blood, bone marrow, lung, heart, liver, kidney, small intestine, colon, and brain.

114. The method of claim 110, wherein said construct is administered intravenously.

30

115. The method of claim 114, wherein said construct is targeted to a tumor.

116. The method of claim 114, wherein said construct is targeted to a tumor by a promoter.

5 117. The method of claim 110, wherein said construct is administered directly into a tumor.

118. The method of claim 117, wherein said tumor is a tumor of tissue selected from the group consisting of skin, breast, prostate, blood, bone
10 marrow, lung, heart, liver, kidney, small intestine, colon, brain, and ovary.

119. The method of claim 110, wherein said construct is administered into cells surrounding a tumor.

15 120. The method of claim 110, wherein said construct is injected into a cavity within which a tumor is found.

121. The method of claim 110, wherein said construct is injected intraperitoneally.
20

122. The method of claim 110, wherein said construct is administered into one or both lungs of said mammal.

123. The method of claim 110, wherein said cytokine is an
25 interferon ω .

124. The method of claim 123, wherein interferon ω is a polypeptide comprising amino acids -23 to 172 in SEQ ID No. 8.

30 125. The method of claim 123, wherein interferon ω is a polypeptide comprising amino acids 1 to 172 in SEQ ID No. 8.

126. The method of claim 123, wherein said polynucleotide construct is VR4151 (SEQ ID No. 44).

127. The method of claim 110, wherein said cytokine is an
5 interferon α .

128. The method of claim 127, wherein interferon α is a polypeptide comprising amino acids -23 to 166 of SEQ ID No. 10.

10 129. The method of claim 127, wherein interferon α is a polypeptide comprising amino acids 1 to 166 of SEQ ID No. 10.

130. The method of claim 127, wherein said polynucleotide construct is VR4112 (SEQ ID No. 2).

15

131. The method of claim 110, wherein a polynucleotide construct comprising a DNA molecule encoding an interferon ω is administered directly into a tumor to treat melanoma.

20 132. The method of claim 110, wherein said mammal is a human.

133. The method of claim 110, wherein said construct is free from association with transfection-facilitating proteins, viral particles, liposomes,
25 cationic lipids, and calcium phosphate precipitating agents.

134. The method of claim 110, wherein said construct is administered as a complex of said construct and one or more cationic compounds selected from the group consisting of cationic lipids, cationic
30 peptides, cationic proteins, cationic polymers, and mixtures thereof.

135. The method of claim 134, wherein said one or more cationic compounds are one or more cationic lipids.

136. The method of claim 135, wherein said cationic lipids are
5 selected from the group consisting of (±)-N-(2-hydroxyethyl)-N,N-dimethyl-2,3-bis(tetradecyloxy)-1-propaniminium bromide, (±)-N-(2-aminoethyl)-N,N-dimethyl-2,3-bis(tetradecyloxy)-1-propaniminium bromide, (±)-N-(3-aminopropyl)-N,N-dimethyl-2,3-bis(dodecyloxy)-1-propaniminium bromide, 5-carboxyspermylglycine dioctadecylamide, dipalmitoyl-
10 phophatidylethanolamine 5-carboxy-spermylamide, {3β-[N-N',N'-dimethylamino)ethane]-carbomoyl}-cholesterol, dimethyldioctdecylammonium bromide, (±)-N,N-dimethyl-N-[2-(sperminecarboxamido)ethyl]-2,3-bis(dioleyloxy)-1-propaniminium pentahydrochloride, DL-1,2-dioleoyl-3-dimethylaminopropyl-β-hydroxyethylammonium, 1-O-oleyl-2-oleoyl-3-
15 dimethylaminopropyl-β-hydroxyethylammonium, cationic lipids are 3,5-(NN-dilysyl)-diaminobenzoyl-3-(DL-1,2-dioleoyl-dimethylaminopropyl-β-hydroxyethylamine) and 3,5-(N,N-di-lysyl)diamino-benzoylglycyl-3-(DL-1,2-dioleoyl-dimethylaminopropyl-β-hydroxyethylamine) and mixtures thereof.

20 137. The method of claim 135, wherein said cationic lipids comprise (±)-N-(2-hydroxyethyl)-N,N-dimethyl-2,3-bis(tetradecyloxy)-1-propaniminium bromide.

138. The method of claim 135, wherein said complex further
25 comprising one or more neutral lipids.

139. The method of claim 135, wherein said neutral lipids comprise 1,2-dioleoyl-glycero-3-phosphoethanolamine.

30 140. The method of claim 138, wherein said polynucleotide construct is complexed with (±)-N-(2-hydroxyethyl)-N,N-dimethyl-2,3-

-106-

bis(tetradecyloxy)-1-propaniminium bromide and 1,2-dioleoyl-glycero-3-phosphoethanolamine.

141. The method of claim 140, wherein the ratio of said
5 polynucleotide construct to lipid is about 5:1 to about 1:1.

142. The method of claim 110, wherein said polynucleotide
construct is VR4151 (SEQ ID No. 4) complexed with (\pm)-N-(2-hydroxyethyl)-
N,N-dimethyl-2,3-bis(tetradecyloxy)-1-propaniminium bromide and 1,2-
10 dioleoyl-glycero-3-phosphoethanolamine such that the ratio of VR4151 to
lipid is about 5:1 to 1:1.

143. The method of claim 110, wherein said polynucleotide
construct is VR4112 (SEQ ID No. 4) complexed with (\pm)-N-(2-hydroxyethyl)-
15 N,N-dimethyl-2,3-bis(tetradecyloxy)-1-propaniminium bromide and 1,2-
dioleoyl-glycero-3-phosphoethanolamine such that the mass ratio of VR4112
(SEQ ID No. 4) to lipid is from about 5:1 to about 1:1.

144. The method of claim 110, wherein said polynucleotide is DNA
20 operably linked to a promoter.

145. The method of claim 110, wherein said polynucleotide is RNA.

146. The method of claim 110, wherein said polynucleotide
25 construct further comprises a polynucleotide encoding one or more additional
molecule.

147. The method of claim 110, wherein said polynucleotide
construct further comprises a polynucleotide encoding one or more additional
30 cytokines.

-107-

148. The method of claim 110, wherein said polynucleotide comprises a region regulating gene expression.

149. The method of claim 148, wherein said region regulating gene expression is cell specific or tissue specific.

150. The method of claim 149, wherein said region is tumor cell or tumor tissue specific.

151. A method of treating cancer, or metastasis thereof, in a mammal, comprising
the method of claim 110 in combination with one or more additional cancer treatment methods selected from the group consisting of surgery, radiation therapy, chemotherapy, immunotherapy, and gene therapy.

152. The method of claim 151, wherein said polynucleotide construct is administered prior to the commencement of said one or more additional cancer treatment methods.

153. The method of claim 151, wherein said polynucleotide construct is administered during the practice of said at one or more additional cancer treatment methods.

154. The method of claim 151, wherein said polynucleotide construct is administered after the end of said one or more additional cancer treatment methods.

155. A method of treating cancer in a mammal, comprising:
administering into a body cavity of said mammal a non-infectious, non-integrating polynucleotide construct comprising a polynucleotide encoding a cytokine, or an active fragment thereof, such that said cytokine is delivered to a tumor in a therapeutically effective amount.

156. The method of 155, wherein said cytokine is selected from the group consisting of IFN ω , IFN α , IFN τ , IFN γ , IFN β , IL-1, IL-2, IL-4, IL-7, IL-12, IL-15, IL-18, GM-CSF, an active fragment of any of thereof, and a combination of any of thereof.

157. The method of claim 156, wherein said cytokine is an interferon ω .

158. The method of claim 157, wherein said interferon ω is a polypeptide comprising amino acids -23 to 172 in SEQ ID No. 8.

159. The method of claim 157, wherein said interferon ω is a polypeptide comprising amino acids 1 to 172 in SEQ ID No. 8.

160. The method of claim 157, wherein said polynucleotide construct is VR4151 (SEQ ID No. 4).

161. The method of claim 156, wherein said cytokine is an interferon α .

162. The method of claim 161, wherein said interferon α is a polypeptide comprising amino acids -23 to 166 of SEQ ID No. 10.

163. The method of claim 161, wherein said interferon α is a polypeptide comprising amino acids 1 to 166 of SEQ ID No. 10.

164. The method of claim 162, wherein said polynucleotide construct is VR4112 (SEQ ID No. 2).

165. The method of claim 156, wherein said cytokine is an interleukin-2.

166. The method of claim 165, wherein interleukin-2 is a polypeptide comprising amino acids 58 to 105 of SEQ ID No. 14.

167. The method of claim 165, wherein interleukin-2 is a polypeptide comprising amino acids 20 to 126 of SEQ ID No. 14.

168. The method of claim 165, wherein said polynucleotide construct is VR1103 (SEQ ID No. 25).

169. The method of claim 155, wherein said polynucleotide construct is administered free from *ex vivo* cells or *ex vivo* cellular material.

170. The method of 155, wherein said tumor disseminates in a body cavity.

171. The method of 170, wherein the body cavity is selected from the group consisting of peritoneal cavity, thoracic cavity, plural cavity, rectal cavity, stomach cavity, and urinary bladder vesicle.

172. The method of 171, wherein the body cavity is peritoneal cavity.

173. The method of claim 155, wherein said cancer is selected from the group consisting of renal cell carcinoma, colorectal carcinoma, lymphoma, melanoma, prostate cancer, ovarian cancer, lung cancer, stomach cancer, glandular lymphoma, bladder cancer, uterine cancer, mesenteric cancer, pancreatic cancer, gastric cancer, metastases thereof, and combinations thereof.

-110-

174. The method of claim 155, wherein said construct is targeted to a tumor.

175. The method of claim 155, wherein said construct is injected
5 intraperitoneally.

176. The method of claim 155, wherein said mammal is a human.

10 177. The method of claim 155, wherein said construct is free from association with transfection-facilitating proteins, viral particles, liposomes, cationic lipids, and calcium phosphate precipitating agents.

178. The method of claim 155, wherein said construct is
15 administered as a complex of said construct and one or more cationic compounds selected from the group consisting of cationic lipids, cationic peptides, cationic proteins, cationic polymers, and mixtures thereof.

179. The method of claim 178, wherein said one or more cationic
20 compounds are one or more cationic lipids.

180. The method of claim 179, wherein said cationic lipids are selected from the group consisting of (\pm)-N-(2-hydroxyethyl)-N,N-dimethyl-2,3-bis(tetradecyloxy)-1-propaniminium bromide, (\pm)-N-(2-aminoethyl)-N,N-dimethyl-2,3-bis(tetradecyloxy)-1-propaniminium bromide, (\pm)-N-(3-aminopropyl)-N,N-dimethyl-2,3-bis(dodecyloxy)-1-propaniminium bromide, 5-carboxyspermylglycine dioctadecylamide, dipalmitoyl-phosphatidylethanolamine 5-carboxy-spermylamide, {3 β -[N-N',N'-dimethylamino)ethane]-carbomoyl}-cholesterol, dimethyldioctdecyl-ammonium bromide, (\pm)-N,N-dimethyl-N-[2-(sperminecarboxamido)ethyl]-2,3-bis(dioleyloxy)-1-propaniminium pentahydrochloride, DL-1,2-dioleoyl-3-dimethylaminopropyl- β -hydroxyethylammonium, 1-O-oleyl-2-oleoyl-3-

25
30

dimethylaminopropyl- β -hydroxyethylammonium, cationic lipids are 3,5-(NN-dilysyl)-diaminobenzoyl-3-(DL-1,2-dioleoyl-dimethylaminopropyl- β -hydroxyethylamine) and 3,5-(N,N-di-lysyl)diamino-benzoylglycyl-3-(DL-1,2-dioleoyl-dimethylaminopropyl- β -hydroxyethylamine) and mixtures thereof.

5

181. The method of claim 179, wherein said cationic lipids comprise (\pm)-N-(2-hydroxyethyl)-N,N-dimethyl-2,3-bis(tetradecyloxy)-1-propaniminium bromide.

10

182. The method of claim 179, wherein said complex further comprising one or more neutral lipids.

183. The method of claim 179, wherein said neutral lipids comprise 1,2-dioleoyl-glycero-3-phosphoethanolamine.

15

184. The method of claim 182, wherein said polynucleotide construct is complexed with (\pm)-N-(2-hydroxyethyl)-N,N-dimethyl-2,3-bis(tetradecyloxy)-1-propaniminium bromide and 1,2-dioleoyl-glycero-3-phosphoethanolamine.

20

185. The method of claim 184, wherein the ratio of said polynucleotide construct to lipid is about 5:1 to about 1:1.

25

186. The method of claim 155, wherein said polynucleotide construct is VR4151 (SEQ ID No. 4) complexed with (\pm)-N-(2-hydroxyethyl)-N,N-dimethyl-2,3-bis(tetradecyloxy)-1-propaniminium bromide and 1,2-dioleoyl-glycero-3-phosphoethanolamine such that the mass ratio of VR4151 (SEQ ID No. 4) to lipid is from about 5:1 to about 1:1.

30

187. The method of claim 155, wherein said polynucleotide construct is VR4112 (SEQ ID No. 4) complexed with (\pm)-N-(2-hydroxyethyl)-N,N-dimethyl-2,3-bis(tetradecyloxy)-1-propaniminium bromide and 1,2-

dioleoyl-glycero-3-phosphoethanolamine such that the mass ratio of VR4112 (SEQ ID No. 4) to lipid is from about 5:1 to about 1:1.

188. The method of claim 155, wherein said polynucleotide
5 construct is VR1103 (SEQ ID No. 25) complexed with (\pm)-N-(2-hydroxyethyl)-N,N-dimethyl-2,3-bis(tetradecyloxy)-1-propaniminium bromide and 1,2-dioleoyl-glycero-3-phosphoethanolamine such that the mass ratio of VR1103 (SEQ ID No. 25) to lipid is from about 5:1 to about 1:1.

10 189. The method of claim 156, wherein a polynucleotide construct comprising a DNA molecule encoding interferon ω is administered to treat ovarian carcinoma.

15 190. The method of claim 156, wherein a polynucleotide construct comprising a DNA molecule encoding interferon ω is administered to treat metastasis of ovarian carcinoma.

20 191. The method of claim 156, wherein a polynucleotide construct comprising a DNA molecule encoding interleukin-2 is administered to treat ovarian carcinoma.

25 192. The method of claim 156, wherein a polynucleotide construct comprising a DNA molecule encoding interleukin-2 is administered to treat metastasis of ovarian carcinoma.

193. The method of claim 156, wherein a polynucleotide construct comprising a DNA molecule encoding an interleukin-2 is administered to treat melanoma.

30 194. The method of claim 156, wherein a polynucleotide construct comprising a DNA molecule encoding an interleukin-2 is administered to treat metastasis of melanoma.

195. The method of claim 155, wherein said polynucleotide is DNA operably linked to a promoter.

5 196. The method of claim 155, wherein said polynucleotide is RNA.

197. The method of claim 155, wherein said polynucleotide construct further comprises a polynucleotide encoding one or more additional molecule.

10

198. The method of claim 155, wherein said polynucleotide construct further comprises a polynucleotide encoding one or more additional cytokines.

15 199. The method of claim 155, wherein said polynucleotide comprises a region regulating gene expression.

200. The method of claim 199, wherein said region regulating gene expression is cell specific or tissue specific.

20

201. The method of claim 200, wherein said region is tumor cell or tumor tissue specific.

202. A method of treating cancer, or metastasis thereof, in a mammal, comprising:

25

the method of claim 155 in combination with one or more additional cancer treatment methods selected from the group consisting of surgery, radiation therapy, chemotherapy, immunotherapy, and gene therapy.

30

203. The method of claim 202, wherein said polynucleotide construct is administered prior to the commencement of said one or more additional cancer treatment methods.

204. The method of claim 202, wherein said polynucleotide construct is administered during the practice of said at one or more additional cancer treatment methods.

5

205. The method of claim 202, wherein said polynucleotide construct is administered after the end of said one or more additional cancer treatment methods.

10

206. A method of selectively transfecting malignant cells in a body cavity of a mammal, comprising:

administering into a body cavity of said mammal a non-infectious, non-integrating polynucleotide construct comprising a polynucleotide encoding a molecule, or an active fragment thereof, such that said molecule is delivered substantially to and expressed in malignant cells within said body cavity.

15

207. The method of claim 206, wherein said molecule is a peptide, polypeptide or protein.

20

208. The method of claim 206, wherein said molecule is cytoplasmic, nuclear, secreted, or membrane bound.

209. The method of claim 206, wherein said molecule induces wherein said molecule induces rejection, regression or prevents progression of said malignant cells in the mammal.

25

210. The method of claim 206, wherein said molecule is immunogenic.

30

-115-

211. The method of claim 206, wherein said polynucleotide encodes at least one member selected from the group consisting of cytokine, lymphokine Class I or II histocompatibility antigens, costimulatory molecules, bacterial antigen, viral glycoprotein, lysozyme, recombinant
5 antibody, adhesion molecule, toxic peptide, and tumor suppressor.

212. The method of 206, wherein said molecule is a cytokine.

213. The method of 212, wherein said cytokine is selected from the
10 group consisting IFN ω , IFN α , IFN τ , IFN γ , IFN β , IL-1, IL-2, IL-4, IL-7, IL-12, IL-15, IL-18, GM-CSF, an active fragment of any of thereof, and a combination of any of thereof.

214. The method of claim 213, wherein said cytokine is an
15 interferon ω .

215. The method of claim 214, wherein said interferon ω is a polypeptide comprising amino acids -23 to 172 in SEQ ID No. 8.

20 216. The method of claim 214, wherein said interferon ω is a polypeptide comprising amino acids 1 to 172 in SEQ ID No. 8.

217. The method of claim 214, wherein said polynucleotide construct is VR4151 (SEQ ID No. 4).

25

218. The method of claim 213, wherein said cytokine is an interferon α .

219. The method of claim 218, wherein interferon α is a polypeptide
30 comprising amino acids -23 to 166 of SEQ ID No. 10.

-116-

220. The method of claim 218, wherein interferon α is a polypeptide comprising amino acids 1 to 166 of SEQ ID No. 10.

221. The method of claim 218, wherein said polynucleotide
5 construct is VR4112 (SEQ ID No. 2).

222. The method of claim 213, wherein said cytokine is an interleukin-2.

10 223. The method of claim 222, wherein interleukin-2 is a polypeptide comprising amino acids 58 to 105 of SEQ ID No. 14.

224. The method of claim 222, wherein interleukin-2 is a polypeptide comprising amino acids 20 to 126 of SEQ ID No. 14.

15 225. The method of claim 222, wherein said polynucleotide construct is VR1103 (SEQ ID No. 25).

226. The method of claim 206, wherein said malignant cells form a
20 cancer in the mammal.

227. The method of claim 226, wherein said molecule is encoded in an amount sufficient to suppress the dissemination of said malignant cells within said body cavity.

25 228. The method of claim 206, wherein the body cavity is selected from the group consisting of peritoneal cavity, thoracic cavity, plural cavity, rectal cavity, stomach cavity, and urinary bladder vesicle.

30 229. The method of claim 228, wherein the body cavity is peritoneal cavity.

230. The method of claim 227, wherein said cancer is selected from the group consisting of renal cell carcinoma, colorectal carcinoma, lymphoma, melanoma, prostate cancer, ovarian cancer, lung cancer, stomach cancer, glandular lymphoma, bladder cancer, uterine cancer, mesenteric
5 cancer, pancreatic cancer, gastric cancer, metastases thereof, and combinations thereof.

231. The method of claim 226, wherein said cancer is pancreatic, gastric, colon, ovarian, or mesenteric cancer.

10

232. The method of claim 206, wherein said administering is by injection or infusion.

233. The method of claim 206, wherein said polynucleotide
15 construct is administered free from *ex vivo* cells or *ex vivo* cellular material.

234. The method of claim 206, wherein said construct is targeted to a tumor.

20 235. The method of claim 206, wherein said construct is injected intraperitoneally.

236. The method of claim 206, wherein said mammal is a human.

25

237. The method of claim 206, wherein said construct is free from association with transfection-facilitating proteins, viral particles, liposomes, cationic lipids, and calcium phosphate precipitating agents.

30 238. The method of claim 206, wherein said construct is administered as a complex of said construct and one or more cationic

-118-

compounds selected from the group consisting of cationic lipids, cationic peptides, cationic proteins, cationic polymers, and mixtures thereof.

239. The method of claim 238, wherein said one or more cationic
5 compounds are one or more cationic lipids.

240. The method of claim 239, wherein said cationic lipids are
selected from the group consisting of (\pm)-N-(2-hydroxyethyl)-N,N-dimethyl-
2,3-bis(tetradecyloxy)-1-propaniminium bromide, (\pm)-N-(2-aminoethyl)-N,N-
10 dimethyl-2,3-bis(tetradecyloxy)-1-propaniminium bromide, (\pm)-N-(3-
aminopropyl)-N,N-dimethyl-2,3-bis(dodecyloxy)-1-propaniminium bromide,
5-carboxyspermylglycine dioctadecylamide, dipalmitoyl-
phosphatidylethanolamine 5-carboxy-spermylamide, {3 β -[N-N',N'-
dimethylamino)ethane]-carbomoyl}-cholesterol, dimethyldioctdecyl-
15 ammonium bromide, (\pm)-N,N-dimethyl-N-[2-(sperminecarboxamido)ethyl]-
2,3-bis(dioleoyloxy)-1-propaniminium pentahydrochloride, DL-1,2-dioleoyl-3-
dimethylaminopropyl- β -hydroxyethylammonium, 1-O-oleyl-2-oleoyl-3-
dimethylaminopropyl- β -hydroxyethylammonium, cationic lipids are 3,5-(NN-
dilysyl)-diaminobenzoyl-3-(DL-1,2-dioleoyl-dimethylaminopropyl- β -
20 hydroxyethylamine) and 3,5-(N,N-di-lysyl)diamino-benzoylglycyl-3-(DL-1,2-
dioleoyl-dimethylaminopropyl- β -hydroxyethylamine) and mixtures thereof.

241. The method of claim 239, wherein said cationic lipids comprise
(\pm)-N-(2-hydroxyethyl)-N,N-dimethyl-2,3-bis(tetradecyloxy)-1-
25 propaniminium bromide.

242. The method of claim 239, wherein said complex further
comprising one or more neutral lipids.

30 243. The method of claim 239, wherein said neutral lipids comprise
1,2-dioleoyl-glycero-3-phosphoethanolamine.

244. The method of claim 242, wherein said polynucleotide construct is complexed with (\pm)-N-(2-hydroxyethyl)-N,N-dimethyl-2,3-bis(tetradecyloxy)-1-propaniminium bromide and 1,2-dioleoyl-glycero-3-phosphoethanolamine.

5

245. The method of claim 244, wherein the ratio of said polynucleotide construct to lipid is about 5:1 to about 1:1.

246. The method of claim 206, wherein said polynucleotide
10 construct is VR4151 (SEQ ID No. 4) complexed with (\pm)-N-(2-hydroxyethyl)-N,N-dimethyl-2,3-bis(tetradecyloxy)-1-propaniminium bromide and 1,2-dioleoyl-glycero-3-phosphoethanolamine such that the mass ratio of VR4151 (SEQ ID No. 4) to lipid is from about 5:1 to about 1:1.

15 247. The method of claim 206, wherein said polynucleotide construct is VR4112 (SEQ ID No. 4) complexed with (\pm)-N-(2-hydroxyethyl)-N,N-dimethyl-2,3-bis(tetradecyloxy)-1-propaniminium bromide and 1,2-dioleoyl-glycero-3-phosphoethanolamine such that the mass ratio of VR4112 (SEQ ID No. 4) to lipid is from about 5:1 to about 1:1.

20

248. The method of claim 206, wherein said polynucleotide
construct is VR1103 (SEQ ID No. 25) complexed with (\pm)-N-(2-
hydroxyethyl)-N,N-dimethyl-2,3-bis(tetradecyloxy)-1-propaniminium
bromide and 1,2-dioleoyl-glycero-3-phosphoethanolamine such that the mass
25 ratio of VR1103 (SEQ ID No. 25) to lipid is from about 5:1 to about 1:1.

249. The method of claim 213, wherein a polynucleotide construct comprising a DNA molecule encoding interferon ω is administered to treat ovarian carcinoma.

30

250. The method of claim 213, wherein a polynucleotide construct comprising a DNA molecule encoding interferon ω is administered to treat metastasis of ovarian carcinoma.

5 251. The method of claim 213, wherein a polynucleotide construct comprising a DNA molecule encoding interleukin-2 is administered to treat ovarian carcinoma.

252. The method of claim 213, wherein a polynucleotide construct
10 comprising a DNA molecule encoding interleukin-2 is administered to treat metastasis of ovarian carcinoma.

253. The method of claim 213, wherein a polynucleotide construct comprising a DNA molecule encoding an interleukin-2 is administered to treat
15 melanoma.

254. The method of claim 213, wherein a polynucleotide construct comprising a DNA molecule encoding an interleukin-2 is administered to treat metastasis of melanoma.
20

255. The method of any one of claim 206, wherein said polynucleotide is DNA operably linked to a promoter.

256. The method of claim 206, wherein said polynucleotide is RNA.
25

257. The method of claim 206, wherein said polynucleotide construct further comprises a polynucleotide encoding one or more additional molecule.

30 258. The method of claim 206, wherein said polynucleotide construct further comprises a polynucleotide encoding one or more additional cytokines.

-121-

259. The method of claim 206, wherein said polynucleotide comprises a region regulating gene expression.

5 260. The method of claim 259, wherein said region regulating gene expression is cell specific or tissue specific.

261. The method of claim 260, wherein said region is tumor cell or tumor tissue specific.

10

262. A method of treating cancer, or metastasis thereof, in a mammal, comprising:

the method of claim 227 in combination with one or more additional cancer treatment methods selected from the group consisting of surgery,
15 radiation therapy, chemotherapy, immunotherapy, and gene therapy.

263. The method of claim 262, wherein said polynucleotide construct is administered prior to the commencement of said one or more additional cancer treatment methods.

20

264. The method of claim 262, wherein said polynucleotide construct is administered during the practice of said at one or more additional cancer treatment methods.

25

265. The method of claim 262, wherein said polynucleotide construct is administered after the end of said one or more additional cancer treatment methods.

1/27




FIG.1

2/27




FIG. 2A




FIG. 2B

3/27




FIG.3A




FIG.3B

4/27




FIG.3C




FIG.3D

5/27




FIG.3E




FIG.3F

6/27




FIG.4A




FIG.4B

7/27




FIG.5A




FIG.5B

8/27




FIG.6

9/27




FIG.7A




FIG.7B

10/27




FIG.8A




FIG.8B

11/27




FIG.8C




FIG.8D

12/27




FIG.9A




FIG.9B




FIG.9C




FIG.9D

14/27




FIG. 10A




FIG. 10B

15/27




FIG.11A




FIG.11B

16/27




FIG.12A




FIG.12B

17/27




FIG. 13A




FIG. 13B

18/27




FIG.14A




FIG.14B




FIG. 15B




FIG. 15A

20/27




FIG. 15D




FIG. 15C

21/27




FIG. 15F




FIG. 15E




FIG.16A




FIG.16B

23/27




FIG.17A




FIG.17B




FIG.18A




FIG.18B

25/27






FIG.19C

FIG.19B

FIG.19A




FIG.19F




FIG.19E




FIG.19D

27/27




FIG.20A




FIG.20B

SEQUENCE LISTING

<110> VICAL

Horton, Holly

Parker, Suezanne

Manthorpe, Marston

Felgner, Philip

<120> Treatment of Cancer Using Cytokine-Expressing
Polynucleotides and Compositions Therefor

<130> 1530.006PC03

<140>

<141>

<150> US 60/067,087

<151> 1997-11-20

<150> US 60/079,914

<151> 1998-03-30

<150> US 60/100,820

<151> 1998-09-15

<160> 26

<170> PatentIn Ver. 2.0

<210> 1

<211> 5428

<212> DNA

<213> Homo sapiens

<400> 1

tcgcgcgttt cgtgatgac ggtgaaaacc tctgacacat gcagctcccg gagacgggtca 60
cagcttgtct gtaagcggat gccgggagca gacaagcccg tcagggcgcg tcagcgggtg 120
ttggcgggtg tcggggctgg cttaactatg cggcatcaga gcagattgta ctgagagtgc 180
accatatgcg gtgtgaaata ccgcacagat gcgtaaggag aaaataccgc atcagattgg 240
ctattggcca ttgcatacgt tgtatccata tcataatag tacatttata ttggctcatg 300
tccaacatta ccgccatggt gacattgatt attgactagt tattaatagt aatcaattac 360

ggggtcatta gttcatagcc catatatgga gttccgcgtt acataactta cggtaaattgg 420
cccgcctggc tgaccgcccac acgacccccg cccattgacg tcaataatga cgtatgttcc 480
catagtaacg ccaataggga ctttccattg acgtcaatgg gtggagtatt tacggtaaac 540
tgcccacttg gcagtacatc aagtgtatca tatgccaaagt acgcccccta ttgacgtcaa 600
tgacggtaaa tggcccgcct ggcattatgc ccagtacatg accttatggg actttcctac 660
ttggcagtac atctacgtat tagtcacgc tattaccatg gtgatgcggg tttggcagta 720
catcaatggg cgtggatagc ggtttgactc acggggattt ccaagtctcc accccattga 780
cgtcaatggg agtttgtttt ggcacaaaaa tcaacgggac tttccaaaat gtcgtaacaa 840
ctccgccccca ttgacgcaaa tgggcggtag gcgtgtacgg tgggaggtct atataagcag 900
agctcgttta gtgaaccgtc agatcgctg gagacgccat ccacgtgtt ttgacctcca 960
tagaagacac cgggaccgat ccagcctcg cggccgggaa cgggtgcattg gaacgcggat 1020
tccccgtgcc aagagtgaac taagtaccgc ctatagactc tataggcaca cccctttggc 1080
tcttatgcat gctatactgt ttttggcttg gggcctatac acccccgctt ccttatgcta 1140
taggtgatgg tatagcttag cctatagggtg tgggttattg accattattg accactcccc 1200
tattggtgac gatactttcc attactaatc cataacatgg ctctttgcca caactatctc 1260
tattggetat atgccaatac tctgtccttc agagactgac acggactctg tatttttaca 1320
ggatggggtc ccatttatta ttacaaatt cacatataca acaacgccgt cccccgtgcc 1380
cgcagttttt attaaacata gcgtgggac tccacgcgaa tctcgggtac gtgttccgga 1440
catgggctct tctccggtag cggcggagct tccacatccg agccctggtc ccatgcctcc 1500
agcggctcat ggtcgtcgg cagctccttg ctctaacag tggaggccag acttaggcac 1560
agcacaatgc ccaccaccac cagtgtgcc cacaaggccg tggcggtagg gtatgtgtct 1620
gaaaatgagc gtggagattg ggctcgcacg gctgacgcag atggaagact taaggcagcg 1680
gcagaagaag atgcaggcag ctgagttgtt gtattctgat aagagtcaga ggtaactccc 1740
gttgcggtgc tgtaacggg ggagggcagt gtagtctgag cagtactcgt tgctgcccg 1800
cgcgccacca gacataatag ctgacagact aacagactgt tcctttccat gggctctttc 1860
tgcatatggc ctgcctctt gctttactga tggctcgtt ggtgctcagc tgcaagtcaa 1920

gctgctctct gggctgtgat ctccctgaga cccacagcct ggataacagg aggaccttga 1980
tgctcctggc acaaatgagc agaatctctc ctctcctctg tctgatggac agacatgact 2040
ttggatttcc ccaggaggag tttgatggca accagttcca gaaggctcca gccatctctg 2100
tcctccatga gctgatccag cagatcttca acctctttac cacaaaagat tcctctgctg 2160
cttgggatga ggacctcta gacaaattct gcaccgaact ctaccagcag ctgaatgact 2220
tggaagcctg tgtgatgcag gaggagaggg tgggagaaac tccctgatg aatgcggact 2280
ccatcttggc tgtgaagaaa tacttccgaa gaatcactct ctatctgaca gagaagaaat 2340
acagcccttg tgccctggag gttgtcagag cagaaatcat gagatccctc tctttatcaa 2400
caaacttgca agaaagatta aggaggaagg aataaggatc cagatctgct gtgccttcta 2460
gttgccagcc atctgttgtt tgccctccc ccgtgccttc cttgacctg gaaggtgcca 2520
ctcccactgt ccttctctaa taaaatgagg aaattgcac gcattgtctg agtaggtgtc 2580
attctattct ggggggtggg gtggggcagc acagcaaggg ggaggattgg gaagacaata 2640
gcaggcatgc tggggatgcg gtgggctcta tgggtacca ggtgctgaag aattgaccog 2700
gttctctctg ggccagaaag aagcaggcac atcccttct ctgtgacaca ccctgtccac 2760
gcccctgggt cttagttcca gcccactca taggacactc atagctcagg agggctccgc 2820
cttcaatccc acccgctaaa gtacttgag cggtctctcc ctccctcacc agcccaccaa 2880
accaaacta gcctccaaga gtgggaagaa attaaagcaa gataggctat taagtgcaga 2940
gggagagaaa atgcctcca catgtgagga agtaatgaga gaaatcatag aatttcttcc 3000
gcttctctgc tcactgactc gctgcgctcg gtcgttcggc tgcggcgagc ggtatcagct 3060
cactcaaagg cggtaatagc gttatccaca gaatcagggg ataacgcagg aaagaacatg 3120
tgagcaaaag gccagcaaaa ggccaggaac cgtaaaaagg ccgcgttgct ggcgtttttc 3180
cataggctcc gccccctga cgagcatcac aaaaatcgac gctcaagtca gaggtggcga 3240
aaccgcagag gactataaag ataccaggcg ttccccctg gaagctccct cgtgcgctct 3300
cctgttccga ccctgccgct tacgggatac ctgtccgctt ttctcccttc ggaagcgtg 3360
gcgctttctc aatgctcacg ctgtaggtat ctgagttcgg tgtaggtcgt tcgctccaag 3420
ctgggctgtg tgcacgaacc cccggttcag cccgaccgct gcgccttacc cggttaactat 3480

cgtcttgagt ccaacccggt aagacacgac ttatcgccac tggcagcagc cactggtaac 3540
aggattagca gagcgaggta tgtaggcggt gctacagagt tcttgaagtg gtggcctaac 3600
tacggetaca ctagaaggac agtatttggt atctgcgctc tgctgaagcc agttaccttc 3660
ggaaaaagag ttggtagctc ttgatccggc aaacaaacca ccgctggtag cgggtggtttt 3720
tttgtttgca agcagcagat tacgcgcaga aaaaaaggat ctcaagaaga tcctttgatc 3780
ttttctacgg ggtctgacgc tcagtgaac gaaaactcac gttaagggat tttggtcatg 3840
agattatcaa aaaggatctt cacctagatc cttttaaatt aaaaatgaag ttttaaatca 3900
atctaaagta tatatgagta aacttggtct gacagttacc aatgcttaat cagtgaggca 3960
cctatctcag cgatctgtct atttcgttca tccatagttg cctgactccg gggggggggg 4020
gcgctgaggt ctgcctcgtg aagaagggtg tgctgactca taccaggeet gaatcgcccc 4080
atcatccagc cagaaagtga gggagccacg gttgatgaga gctttgttgt aggtggacca 4140
gttggtgatt ttgaactttt gctttgccac ggaacggtct gcgttgctcg gaagatgcgt 4200
gatctgatcc ttcaactcag caaaagtctg atttattcaa caaagccgcc gtcccgtaa 4260
gtcagcgtaa tgctctgcca gtgttacaac caattaacca attctgatta gaaaaactca 4320
tcgagcatca aatgaaactg caatttattc atatcaggat tatcaatacc atatttttga 4380
aaaagccgtt tctgtaatga aggagaaaac tcaccgaggc agttccatag gatggcaaga 4440
tcctggtatc ggtctgcgat tccgactcgt ccaacatcaa tacaacctat taatttccc 4500
tcgtcaaaaa taaggttatc aagtgagaaa tcaccatgag tgacgactga atccggtgag 4560
aatggcaaaa gcttatgcat ttctttccag acttggtcaa caggccagcc attacgctcg 4620
tcacaaaaat cactcgcac aaccaaaccg ttattcattc gtgattgcgc ctgagcgaga 4680
cgaaatacgc gatcgctgtt aaaaggacaa ttacaaacag gaatcgaatg caaccggcgc 4740
aggaacactg ccagcgcac aacaatattt tcacctgaat caggatattc ttctaatacc 4800
tggaatgctg ttttccggg gatcgagtg gtgagtaacc atgcatcatc aggagtacgg 4860
ataaaatgct tgatggctcg aagaggcata aattccgtca gccagtttag tctgaccatc 4920
tcattctgtaa catcattggc aacgtacct ttgcatgtt tcagaaacaa ctctggcgca 4980
tcgggcttcc catacaatcg atagattgtc gcacctgatt gcccgacatt atcgcgagcc 5040

catttatacc catataaatc agcatccatg ttggaattta atcgcggcct cgagcaagac 5100
gtttcccggt gaatatggct cataacaccc ctgtattac tgtttatgta agcagacagt 5160
tttattgttc atgatgatat atttttatct tgtgcaatgt aacatcagag attttgagac 5220
acaacgtggc tttccccccc ccccatattat tgaagcattt atcaggggta ttgtctcatg 5280
agcggataca tatttgaatg tatitagaaa aataaataaa taggggttcc gcgcacattt 5340
ccccgaaaag tgccacctga cgtctaagaa accattatta tcatgacatt aacctataaa 5400
aataggcgta tcacgaggcc ctttcgtc 5428

<210> 2

<211> 5259

<212> DNA

<213> Homo sapiens

<400> 2

tcgcgcgttt cggatgatgac ggtgaaaacc totgacacat gcagctcccg gagacgggtca 60
cagcttgtct gtaagcggat gccgggagca gacaagcccg tcagggcgcg tcagcgggtg 120
ttggcgggtg tcggggctgg cttaactatg cggcatcaga gcagattgta ctgagagtgc 180
accatatgag gtgtgaaata ccgcacagat gcgtaaggag aaaataccgc atcagattgg 240
ctattggcca ttgcatacgt tgtatccata tcataatatg tacatttata ttggctcatg 300
tccaacatta ccgcatggtt gacattgatt attgactagt tattaatagt aatcaattac 360
ggggtcatta gttcatagcc catatatgga gttccgcgtt acataactta cggtaaatgg 420
cccgcctggc tgaccgcccc acgacccccg cccattgacg tcaataatga cgtatgttcc 480
catagtaacg ccaataggga ctttccattg acgtcaatgg gtggagtatt tacggtaaac 540
tgcccacttg gcagtacatc aagtgtatca tatgccaagt acgcccccta ttgacgtcaa 600
tgacggtaaa tggccgcgct ggcattatgc ccagtacatg acctatggg actttectac 660
ttggcagtag atctacgtat tagtcatcgc tattaccatg gtgatgcggt tttggcagta 720
catcaatggg cgtggatagc ggtttgactc acggggattt ccaagtctcc accccattga 780
cgtcaatggg agtttgtttt ggcacaaaaa tcaacgggac tttccaaaat gtcgtaacaa 840
ctccgcccc a ttgacgcaaa tgggcggtag gcgtgtacgg tgggaggtct atataagcag 900

agctcgttta gtgaaccgtc agatcgcttg gagacgccat ccacgtgtt ttgacctcca 960
tagaagacac cgggaccgat ccagcctccg cggccgggaa cgggtgcattg gaacgcggat 1020
tccccgtgcc aagagtgcg taagtaccgc ctatagactc tataggcaca cccctttggc 1080
tcttatgcat gctatactgt ttttggttg gggcctatac acccccgctt ccttatgcta 1140
taggtgatgg tatagcttag cctataggtg tgggttattg accattattg accactcccc 1200
tattggtgac gatactttcc attactaatc cataacatgg ctctttgcc caactatctc 1260
tattggctat atgccaatac tctgtccttc agagactgac acggactctg tatttttaca 1320
ggatggggtc ccatttatta ttacaaatt cacatataca acaacgcctt cccccgtgcc 1380
cgcagttttt attaaacata gcgtgggac tccacgcgaa tctcgggtac gtgttccgga 1440
catgggctct tctccggtag cggcggagct tccacatccg agcctgggtc ccatgcctcc 1500
agcggctcat ggtcgtcgg cagctccttg ctctaacag tggaggccag acttaggcac 1560
agcacaatgc ccaccaccac cagtgtgccg cacaaggccg tggcggtagg gtatgtgtct 1620
gaaaatgagc gtggagattg ggctcgacg gctgacgcag atggaagact taaggcagcg 1680
gcagaagaag atgcaggcag ctgagttgtt gtattctgat aagagtcaga ggtaactccc 1740
gttgcggtgc tgttaacggt ggagggcagt gtagtctgag cagtactcgt tgctgccgcg 1800
cgcgccacca gacataatag ctgacagact aacagactgt tcctttccat gggctctttc 1860
tgcagatggc ctgcctttt gctttactga tggctctggt ggtgctcagc tgcaagtcaa 1920
gctgctctct gggctgtgat ctccctgaga cccacagcct ggataacagg aggacctga 1980
tgctcctggc acaaatgagc agaactcttc ctctcctctg tctgatggac agacatgact 2040
ttggatttcc ccaggaggag tttgatggca accagttcca gaaggctcca gccatctctg 2100
tcctccatga gctgatccag cagatcttca acctctttac cacaaaagat tcatctgctg 2160
cttgggatga ggacctcta gacaaattct gcaccgaact ctaccagcag ctgaatgact 2220
tggaagcctg tgtgatgcag gaggagagg tgggagaaac tcccctgatg aatgcggact 2280
ccatcttggc tgtgaagaaa tacttccgaa gaatcactct ctatctgaca gagaagaaat 2340
acagcccttg tgctgggag gttgtcagag cagaaatcat gagatccctc tctttatcaa 2400
caaaacttga agaaagatta aggaggaagg aataaggatc cagatctact tctggcta 2460

aaaagatcag agctctagag atctgtgtgt tgggtttttg tgtggtaccc aggtgctgaa 2520
gaattgaccc gggttcctcct gggccagaaa gaagcaggca catccccttc tctgtgacac 2580
accctgtcca cgccctcgtt tcttagttcc agccccactc ataggacact catagctcag 2640
gagggctccg ccttcaatcc caccgctaa agtacttga gcggtctctc cctccctcat 2700
cagcccacca aaccaaact agcctccaag agtggaaga aattaaagca agataggcta 2760
ttaagtgcag agggagagaa aatgcctcca acatgtgagg aagtaatgag agaaatcata 2820
gaatttcttc cgcttcctcg ctactgact cgctgcgctc ggtcggttcg ctgcccgcag 2880
cggatcagc tcaactcaaag gcggaatac ggttatccac agaatacagg gataacgcag 2940
gaaagaacat gtgagcaaaa ggccagcaaa aggccaggaa ccgtaaaaag gccgcgttgc 3000
tggcggtttt ccataggctc cgccccctg acgagcatca caaaaatcga cgctcaagtc 3060
agaggtggcg aaacccgaca ggactataaa gataccaggc gtttccccct ggaagctccc 3120
tcgtgcgctc tctgtttcg accctgccgc ttaccggata cctgtccgcc tttctccctt 3180
cgggaagcgt ggcgctttct caatgctcac gctgtaggta tctcagttcg gtgtaggteg 3240
ttcgctcaa gctgggctgt gtgcacgaac cccccgttca gcccgaccgc tgcgccttat 3300
ccggttaacta tcgtcttgag tccaaccg taagacacga cttatcgcca ctggcagcag 3360
ccactggtaa caggattagc agagcgaggt atgtaggcgg tgctacagag ttcttgaagt 3420
ggtggcctaa ctacggctac actagaagga cagtatttgg tatctgcgct ctgctgaagc 3480
cagttacctt cggaaaaaga gttggtagct cttgatccgg caaacaacc accgctggtg 3540
gcggtggttt ttttgtttgc aagcagcaga ttacgcgcag aaaaaaagga tctcaagaag 3600
atcctttgat cttttctacg ggggtctgacg ctcaagtggaa cgaaaactca cgttaaggga 3660
ttttggtcat gagattatca aaaaggatct tcacctagat ctttttaaataaaaaatgaa 3720
gttttaaatc aatctaaagt atatatgagt aaacttggtc tgacagttac caatgcttaa 3780
tcagtgaggc acctatctca gcgatctgtc tatttcgttc atccatagtt gctgactcc 3840
gggggggggg ggcgctgagg tctgcctcgt gaagaagggtg ttgctgactc ataccaggcc 3900
tgaatcgccc catcatccag ccagaaagtg agggagccac ggttgatgag agctttgttg 3960
taggtggacc agttggtgat tttgaacttt tgctttgccg cggaacggtc tgcgttgctg 4020

ggaagatgcg tgatctgac cttcaactca gcaaaagttc gatttattca acaaagccgc 4080
cgtcccgtca agtcagcgta atgctctgcc agtggtacaa ccaattaacc aattctgatt 4140
agaaaaactc atcgagcatc aaatgaaact gcaatttatt catatcagga ttatcaatac 4200
catatttttg aaaaagccgt ttctgtaatg aaggagaaaa ctcaccgagg cagttccata 4260
ggatggcaag atcctggtat cggctctgca ttecgactcg tccaacatca atacaaccta 4320
ttaatttccc ctctgcaaaa ataaggttat caagtgagaa atcaccatga gtgacgactg 4380
aatccggtga gaatggcaaa agcttatgca tttctttcca gacttggtca acaggccagc 4440
cattacgctc gtcacaaaa tcaactcgat caaccaaacc gttattcatt cgtgattgcg 4500
cctgagcgag acgaaatacg cgatcgctgt taaaaggaca attacaaaca ggaatcgaat 4560
gcaaccggcg caggaacact gccagcgcat caacaatatt ttcacctgaa tcaggatatt 4620
cttctaatac ctggaatgct gttttccgg ggatcgagc ggtgagtaac catgcatcat 4680
caggagtacg gataaaatgc ttgatggtcg gaagaggcat aaattccgtc agccagttta 4740
gtctgaccat ctcatctgta acatcattgg caacgctacc ttgcatgt ttcagaaaca 4800
actctggcgc atcgggcttc ccatacaatc gatagattgt cgcacctgat tgcccacat 4860
tatcgcgagc ccatttatac ccataataat cagcatccat gttggaattt aatcgcggcc 4920
tcgagcaaga cgtttccgt tgaatatggc tcataacacc ccttgatta ctgtttatgt 4980
aagcagacag ttttattggt catgatgata tatttttata ttgtgcaatg taacatcaga 5040
gattttgaga cacaacgtgg ctttcccccc cccccatta ttgaagcatt tatcagggtt 5100
attgtctcat gagcggatgc atatttgaat gtatttagaa aaataaacia ataggggttc 5160
cgcgacatt tccccgaaaa gtgccacctg acgtctaaga aaccattatt atcatgacat 5220
taacctataa aaataggcgt atcacgagge cctttcgtc 5259

<210> 3
<211> 5480
<212> DNA
<213> Homo sapiens

<400> 3
tcgcgcgttt cggatgatgac ggtgaaaacc tctgacacat gcagctcccg gagacgggtca 60
cagcttgtct gtaagcggat gccgggagca gacaagcccg tcagggcgcg tcagcgggtg 120

ttggcgggtg tcggggctgg cttactatg cggcatcaga gcagattgta ctgagagtgc 180
accatatgcg gtgtgaaata ccgcacagat gcgtaaggag aaaataccgc atcagattgg 240
ctattggcca ttgcatacgt tgtatecata tcataatatg tacatttata ttggcctcatg 300
tccaacatta ccgcatgtt gacattgatt attgactagt tattaatagt aatcaattac 360
ggggtcatta gttcatagcc catatatgga gttccgcgtt acataactta cggtaaatgg 420
ccgcctggc tgaccgcca acgaccccg ccattgacg tcaataatga cgtatgtcc 480
catagtaacg ccaatagga ctttccattg acgtcaatgg gtggagtatt tacggtaaac 540
tgcccacttg gcagtacac aagtgtatca tatgccaagt acgccccta ttgacgcaa 600
tgacggtaaa tggcccgct ggcattatgc ccagtacatg accttatggg actttctac 660
ttggcagtac atctacgtat tagtcacgc tattaccatg gtgatgcgtt ttggcagta 720
catcaatggg cgtggatagc gggttgactc acggggattt ccaagtctcc acccattga 780
cgtcaatggg agtttgttt ggcacaaaa tcaacgggac ttccaaaat gtcgtaacaa 840
ctccgcccc ttgacgcaa tgggcggtag gcgtgtacgg tgggaggtct atataagcag 900
agctcgttta gtgaaccgtc agatcgctg gagacgcat ccacgtgtt ttgacctca 960
tagaagacac cgggaccgat ccagcctccg cggccgggaa cggtgcatg gaacgcggat 1020
tcccgtgcc aagagtgcg taagtaccgc ctatagactc tataggcaca ccccttggc 1080
tcttatgcat gctatactgt ttttgcttg gggcctatac accccgctt ccttatgcta 1140
taggtgatgg tatagcttag cctatagtg tgggttattg accattattg accactccc 1200
tattggtgac gatactttcc attactaatc cataacatgg ctctttgcca caactatctc 1260
tattggctat atgccaatac tctgtcctc agagactgac acggactctg tatttttaca 1320
ggatggggtc ccatttatta ttacaaatt cacatataca acaacgctt ccccggtgc 1380
cgcagtttt attaaacata gcgtgggac tccacgcga tctcggtac gtgttcgga 1440
catgggctct tctccgtag cggcgagct tccacatcg agcctggtc ccatgcctcc 1500
agcggctcat ggtcgctcg cagtccttg ctctaacag tggaggccag acttaggcac 1560
agcacaatgc ccaccaccac cagtgtgcg cacaaggccg tggcggtagg gtatgtctc 1620
gaaaatgagc gtggagattg ggctcgacg gctgacgcag atggaagact taaggcagc 1680

gcagaagaag atgcaggcag ctgagttggt gtattctgat aagagtcaga ggtaactccc 1740
gttgcggtgc tgttaacggt ggagggcagt gtagtctgag cagtactpgt tgctgccgcg 1800
cgcgccacca gacataatag ctgacagact aacagactgt tcctttccat gggctctttc 1860
tgcagtcacc gtcgtcgaca cgtgtgatca gatataacca tggccctcct gttccctcta 1920
ctggcagccc tagtgatgac cagctatagc cctgttgat ctctgggctg tgatctgect 1980
cagaaccatg gctacttag caggaacacc ttggtgcttc tgcaccaa at gaggagaatc 2040
tcccccttct tgtgtctcaa ggacagaaga gatttcaggt tccccagga gatggtaaaa 2100
gggagccagt tgcagaaggc ccatgtcatg tctgtctcc atgagatgct gcagcagatc 2160
ttcagcctct tccacacaga gcgtcctct gctgcctgga acatgacct cctagaccaa 2220
ctccacactg gattcatca gcaactgcaa cacctggaga cctgcttgct gcaggtagtg 2280
ggagaaggag aatctgctgg ggcaattagc agcctgcac tgaccttgag gaggtacttc 2340
cagggaatcc gtgtctacct gaaagagaag aaatacagcg actgtgcctg ggaagttgtc 2400
agaatgaaa tcatgaaatc cttgttctta tcaacaaaca tgcaagaaag actgagaagt 2460
aaagatagag acctgggctc atcttgagga tccagatctg ctgtgccttc tagttgccag 2520
ccatctgttg ttgtccctc ccccgctgct tccttgacc tggaagggtc cactccact 2580
gtcctttcct aataaaatga ggaaattgca tgcattgtc tgagtaggtg tcattctatt 2640
ctggggggtg ggggtgggca gcacagcaag ggggaggatt gggaagacaa tagcaggcat 2700
gctggggatg cgggtgggctc tatgggtacc cagggtgctga agaattgacc cggttcctcc 2760
tgggcccagaa agaagcaggc acatccctt ctctgtgaca caccctgtcc acgcccctgg 2820
ttcttagttc cagccccact cataggacac tcatagctca ggagggtcc gccttcaatc 2880
ccaccgcta aagtacttg agcgggtctt cctccctca tcagcccacc aaaccaaacc 2940
tagcctcaa gagtgggaag aaattaaagc aagataggct attaagtga gaggagaga 3000
aaatgcctcc aacatgtgag gaagtaatga gagaaatcat agaatttctt ccgttctc 3060
gtcactgac tgcgtgcgt cggctgttcg gctgcggcga gcggtatcag ctactcaaa 3120
ggcgtaata cggttatcca cagaatcagg ggataacga ggaaagaaca tgtgagcaaa 3180
aggccagcaa aaggccagga accgtaaaaa ggccgcgttg ctggcgttt tccataggct 3240

ccgccccct. gacgagcatc acaaaaatcg acgctcaagt cagaggtggc gaaacccgac 3300
aggactataa agataccagg cgtttcccc tggaagctcc ctcgtgcgt ctcctgttcc 3360
gaccctgccg cttaccggat acctgtccgc ctttctccct tcgggaagcg tggcgctttc 3420
tcaatgctca cgctgtaggt atctcagttc ggtgtaggtc gttegctcca agctgggctg 3480
tgtgcacgaa cccccgttc agcccgaccg ctgcgcctta tccggtaact atcgtcttga 3540
gtccaaccg gtaagacacg acttatcgcc actggcagca gccactggta acaggattag 3600
cagagcgagg tatgtaggcg gtgctacaga gttcttgaag tggtagccta actacggcta 3660
cactagaagg acagtatttg gtatctgcgc tctgctgaag ccagttacct tcggaaaaag 3720
agttggtagc tcttgatccg gcaaaaaac caccgctggg agcggtggtt tttttgtttg 3780
caagcagcag attacgcgca gaaaaaagg atctcaagaa gatcctttga tcttttctac 3840
ggggtctgac gctcagtga acgaaaactc acgttaaggg attttggtca tgagattatc 3900
aaaaaggatc ttcaoctaga tccttttaaa ttaaaaatga agttttaaat caatctaaag 3960
tatatatgag taaacttggg ctgacagtta ccaatgctta atcagtgagg cacctatctc 4020
agcgatctgt ctatttcgtt catccatagt tgcctgactc cggggggggg gggcgctgag 4080
gtctgcctcg tgaagaaggt gttgctgact cataccaggc ctgaatcgcc ccatcatcca 4140
gccagaaagt gagggagcca cggttgatga gagctttggt gtaggtggac cagttggtga 4200
ttttgaactt ttgctttgcc acggaacggt ctgcgttgtc ggaagatgc gtgatctgat 4260
ccttcaactc agcaaaagtt cgatttatc aacaaagccg ccgtcccgtc aagtcagcgt 4320
aatgctctgc cagtgttaca accaattaac caattctgat tagaaaaact catcgagcat 4380
caaatgaaac tgcaatttat tcatatcagg attatcaata ccatattttt gaaaaagccg 4440
tttctgtaat gaaggagaaa actcaccgag gcagttccat aggatggcaa gatcctggta 4500
tcggtctgcg attccgactc gtccaacatc aatacaacct attaatctcc cctcgtcaaa 4560
aataaggtta tcaagtgaga aatcaccatg agtgacgact gaatccggtg agaatggcaa 4620
aagcttatgc atttctttcc agacttggtc aacaggccag ccattacgct cgtcatcaaa 4680
atcactcgca tcaaccaaac cgttattcat tcgtgattgc gcctgagcga gacgaaatac 4740
gcgatcgctg ttaaaaggac aattacaaac aggaatcgaa tgcaaccggc gcaggaacac 4800

tgccagcgca tcaacaatat tttcacctga atcaggatat tcttctaata cctggaatgc 4860
tgttttcccg gggatcgagc tggtagagtaa ccatgcatac tcaggagtag ggataaaatg 4920
cttgatgggc ggaagaggca taaattccgt cagccagttt agtctgacca tctcatctgt 4980
aacatcattg gcaacgctac ctttgccatg tttcagaaac aactctggcg catcgggctt 5040
cccatacaat cgatagattg tcgcacctga ttgcccgaca ttatcgcgag cccatttata 5100
cccatataaa tcagcatcca tgttgaatt taatcgcggc ctgagcaag acgtttcccg 5160
ttgaatatgg ctcataacac cccttgattt actgtttatg taagcagaca gttttattgt 5220
tcatgatgat atatttttat ctgtgcaat gtaacatcag agattttgag acacaacgtg 5280
gctttccccc cccccccatt attgaagcat ttatcagggg tattgtctca tgagcggata 5340
catatttgaa tgtattttaga aaaataaaca aataggggtt ccgcgcacat ttccccgaaa 5400
agtgccacct gacgtctaag aaaccattat tatcatgaca ttaacctata aaaataggcg 5460
tatcacgagg ccctttcgtc 5480

<210> 4
<211> 5322
<212> DNA
<213> Homo sapiens

<400> 4
tcgcgcgttt cggtgatgac ggtgaaaacc tctgacacat gcagctcccg gagacggtca 60
cagcttgtct gtaagcggat gccgggagca gacaagcccg tcaggggcgc tcagcgggtg 120
ttggcgggtg tcggggctgg cttaactatg cggcacaga gcagattgta ctgagagtgc 180
accatagcgc gtgtgaaata ccgcacagat gcgtaaggag aaaataccgc atcagattgg 240
ctattggcca ttgcatacgt tgtatccata tcataatatg tacatttata ttggctcatg 300
tccaacatta ccgcatggt gacattgatt attgactagt tattaatagt aatcaattac 360
ggggtcatta gttcatagcc catatatgga gttecgcggt acataactta cggtaaatgg 420
ccgcctggc tgaccgcca acgacccccg ccattgacg tcaataatga cgtatgttcc 480
catagtaacg ccaataggga ctttcattg acgtcaatgg gtggagtatt tacggtaaac 540
tgcccacttg gcagtacatc aagtgtatca tatgccaaat acgccccta ttgacgtcaa 600

tgacggtaaa tggccgcct ggcattatgc ccagtacatg accttatggg actttcctac 660
ttggcagtac atctacgtat tagtcacgc tattaccatg gtgatgcggg ttgggcagta 720
catcaatggg cgtggatagc ggtttgactc acggggattt ccaagtctcc accccattga 780
cgtcaatggg agtttgtttt ggcacaaaaa tcaacgggac ttccaaaat gtcgtaacaa 840
ctccgccccca ttgacgcaaa tgggcggtag gcgtgtacgg tgggaggtct atataagcag 900
agctcgttta gtgaaccgtc agatgcctg gagacgcat ccacgtgtt ttgacctca 960
tagaagacac cgggaccgat ccagcctcg cgccgggaa cgggtgcattg gaacgcggat 1020
tccccgtgcc aagagtgcg taagtaccgc ctatagactc tataggcaca cccctttggc 1080
tcttatgcat gctatactgt ttttggttg ggccatac acccccgctt ccttatgcta 1140
taggtgatgg tatagcttag cctatagggtg tgggttattg accattattg accactcccc 1200
tattggtgac gatacttcc attactaatc cataacatgg ctctttgcca caactatctc 1260
tattggctat atgccaatac tctgtcctc agagactgac acggactctg ttttttaca 1320
ggatggggtc ccatttatta ttacaaatt cacatataca acaacgccgt cccccgtgcc 1380
cgcagttttt attaaacata gcgtgggac tccacgcgaa tctcgggtac gtgttcgga 1440
catgggctct tctccggtag cggcgagct tccacatccg agccctggtc ccatgcctcc 1500
agcggctcat ggtcgtcgg cagctcctg ctctaacag tggaggccag acttaggcac 1560
agcacaatgc ccaccaccac cagtgtgcg cacaaggccg tggcggtagg gtatgtgtct 1620
gaaaatgagc gtggagattg ggctcgcag gctgacgcag atggaagact taaggcagcg 1680
gcagaagaag atgcaggcag ctgagttgtt gtattctgat aagagtcaga ggtaactccc 1740
gttgcggtgc tgtaacggg ggagggcagt gtagtctgag cagtactcgt tgctgccgcg 1800
cgcgccacca gacataatag ctgacagact aacagactgt tctttccat gggcttttc 1860
tgcagtcacc gtcgtcgaca cgtgtgatca gatatcgcg cgctctaga atggccctcc 1920
tgttccctct actggcagcc ctagtatga ccagctatag cctgttgga tctctgggct 1980
gtgatctgcc tcagaacctt ggcctactta gcaggaacac cttggtgctt ctgcacaaa 2040
tgaggagaat ccccccttc ttgtgtctca aggacagaag agacttcagg ttccccagg 2100
agatggtaaa agggagccag ttgcagaagg ccatgtcat gtctgtcctc catgagatgc 2160

tgcagcagat cttcagcctc ttccacacag agcgctcctc tgctgcctgg aacatgaccc 2220
tcctagacca actccacact ggacttcate agcaactgca acacctggag acctgcttgc 2280
tgcaggtagt gggagaagga gaatctgctg gggcaattag cagccctgca ctgaccttga 2340
ggaggtagt ccaggaatc cgtgtctacc tgaaagagaa gaaatacagc gactgtgcct 2400
gggaagttgt cagaatggaa atcatgaaat ccttgttctt atcaacaaac atgcaagaaa 2460
gactgagaag taaagataga gacctgggct catcttgagg atccagatct acttctggct 2520
aataaaagat cagagctcta gagatctgtg tgttggtttt ttgtgtggtt cccaggtgct 2580
gaagaattga ccggttctt cctgggccag aaagaagcag gcacatcccc ttctctgtga 2640
cacaccctgt ccacgcccct ggttcttagt tccagcccca ctcataggac actcatagct 2700
caggagggct ccgcttcaa tcccaccgc taaagtaett ggagcggctt ctccctccct 2760
catcagccca ccaaaccaaa cctagcctcc aagagtggga agaaattaaa gcaagatagg 2820
ctattaagtg cagagggaga gaaaatgcct ccaacatgtg aggaagtaat gagagaaatc 2880
atagaatttc ttccgcttcc tcgctcactg actcgctgcg ctcggtcgtt cggctgcggc 2940
gagcggatc agctcactca aaggcggtaa tacggttatc cacagaatca ggggataacg 3000
caggaaagaa catgtgagca aaaggccagc aaaaggccag gaaccgtaaa aaggccgcgt 3060
tgctggcggt ttccatagg ctccgcccc ctgacgagca tcacaaaaat cgacgctcaa 3120
gtcagaggtg gcgaaaccg acaggactat aaagatacca ggcgtttccc cctggaagct 3180
ccctcgctgc ctctctgtt ccgacctgc cgttaccgg atacctgtcc gcctttctcc 3240
cttcgggaag cgtggcgctt tctcaatgct cagctgttag gtatctcagt tcggtgtagg 3300
tcgttcgctc caagctgggc tgtgtgcacg aacccccgt tcagcccgac cgctgcgct 3360
tatccggtaa ctatcgtctt ggtccaacc cggtaagaca cgacttatcg cactggcag 3420
cagccactgg taacaggatt agcagagcga ggtatgtagg cggtgctaca gattcttga 3480
agtggggcc taactacggc tacactagaa ggacagtatt tggtatctgc gctctgctga 3540
agccagttac cttcgaaaaa agagttggta gctcttgatc cggcaaaaaa accaccgctg 3600
gtagcggtag ttttttgggt tgcaagcagc agattacgcg cagaaaaaaa ggatctcaag 3660
aagatccttt gatctttct acggggctg acgctcagtg gaacgaaaac tcacgttaag 3720

ggatttttggg catgagatta tcaaaaagga tcttcaccta gatcctttta aattaaaaat 3780
gaagtttttaa atcaatctaa agtatatatg agtaaacttg gtctgacagt taccaatgct 3840
taatcagtga ggcacctatc tcagcgatct gtctatttcg ttcattccata gttgcctgac 3900
tccggggggg gggggcgctg aggtctgcct cgtgaagaag gtgttgctga ctcataccag 3960
gcctgaatcg ccccatcatc cagccagaaa gtgagggagc cacggttgat gagagctttg 4020
ttgtaggtgg accagttggg gattttgaac ttttgctttg ccacggaacg gtctgcgttg 4080
tcgggaagat gcgtgatctg atccttcaac tcagcaaaag ttcgatttat tcaacaaagc 4140
cgccgtcccg tcaagtcagc gtaatgctct gccagtgtta caaccaatta accaattctg 4200
attagaaaaa ctcatcgagc atcaaatgaa actgcaattt attcatatca ggattatcaa 4260
taccatattt ttgaaaaagc cgtttctgta atgaaggaga aaactcaccg aggcagttcc 4320
ataggatggc aagatcctgg tatcggctcg cgattccgac tcgtccaaca tcaatacaac 4380
ctattaattt cccctcgta aaaataaggt tatcaagtga gaaatcacca tgagtgaaga 4440
ctgaatccgg tgagaatggc aaaagcttat gcatttcttt ccagacttgt tcaacaggcc 4500
agccattacg ctogtcatca aaatcactcg catcaaccaa accgttattc attcgtgatt 4560
gcgcctgagc gagacgaaat acgcgatcgc tgttaaaagg acaattacaa acaggaatcg 4620
aatgcaaccg gcgcaggaac actgccagcg catcaacaat attttcacct gaatcaggat 4680
attcttctaa tacctggaat gctgttttcc cggggatcgc agtggtgagt aaccatgcat 4740
catcaggagt acggataaaa tgcttgatgg tcggaagagg cataaattcc gtcagccagt 4800
ttagtctgac catctcatct gtaacatcat tggcaacgct acctttgcca tgtttcagaa 4860
acaactctgg cgcacgggac ttcccataca atcgatagat tgtcgcacct gattgcccga 4920
cattatcgcg agccatttta taccatata aatcagcatc catgttgga tttaatcgcg 4980
gcctcgagca agacgtttcc cgttgaatat ggctcataac accccttgta ttactgttta 5040
tgtaagcaga cagttttatt gttcatgatg atatattttt atcttgtgca atgtaacatc 5100
agagattttg agacacaacg tggctttccc cccccccca ttattgaagc atttatcagg 5160
gttattgtct catgagcgga tacatatitg aatgtattta gaaaaataaa caaatagggg 5220
ttccgcgcac atttcccga aaagtgcac ctgacgtcta agaaaccatt attatcatga 5280

-16-

cattaaccta taaaaatagg cgtatcacga ggccctttcg tc

5322

<210> 5

<211> 5422

<212> DNA

<213> Mus musculus

<400> 5

tcgcgcggtt cggatgatgac ggtgaaaacc tctgacacat gcagctcccg gagacgggtca 60
cagcttgtct gtaagcggat gccgggagca gacaagcccg tcagggcgcg tcagcgggtg 120
ttggcgggtg tcggggctgg cttactatg cggcatcaga gcagattgta ctgagagtgc 180
accatagcg gtgtgaaata ccgcacagat gcgtaaggag aaaataccgc atcagattgg 240
ctattggcca ttgcatacgt tgtatccata tcataatatg tacatttata ttggctcatg 300
tccaacatta ccgccatgtt gacattgatt attgactagt tattaatagt aatcaattac 360
ggggtcatta gttcatagcc catatatgga gttccgcgtt acataactta cggtaaattg 420
cccgcctggc tgaccgccc acgacccccc cccattgacg tcaataatga cgtatgttcc 480
catagtaacg ccaataggga ctttcattg acgtcaatgg gtggagtatt tacggtaaac 540
tgcccacttg gcagtacatc aagtgtatca tatgccagt acgccccta ttgacgtcaa 600
tgacggtaaa tggccgcctt ggcattatgc ccagtacatg accttatggg actttcctac 660
ttggcagtac atctacgtat tagtcatcgc tattaccatg gtgatgcggg tttggcagta 720
catcaatggg cgtggatagc ggtttgactc acggggattt ccaagtctcc accccattga 780
cgtaaatggg agtttgtttt ggacacaaaa tcaacgggac ttccaaaat gtcgtaacaa 840
ctccgcccc ttgacgcaaa tgggcggtag gcgtgtacgg tgggaggtct atataagcag 900
agctcgttta gtgaaccgtc agatcgctg gagacgcat ccacgtgtt ttgacctca 960
tagaagacac cgggaccgat ccagctccg cggccgggaa cgggtgcattg gaacgcggat 1020
tccccgtgcc aagagtgcg taagtaccgc ctatagactc tataggcaca cccctttggc 1080
tcttatgcgt gctatactgt ttttggttg ggccctatac accccgctt ccttatgcta 1140
taggtgatgg tatagcttag cctataggtg tgggttattg accattattg accactcccc 1200
tattggtgac gatactttcc attactaatc cataacatgg ctctttgcca caactatctc 1260
tattggctat atgccaatac tctgtccttc agagactgac acggactctg tatttttaca 1320

ggatggggtc ccatttatta ttacaaatt cacatataca acaacgccgt ccccggtgcc 1380
cgagttttt attaaacata gcgtgggac tccacgcgaa tctcgggtac gtgttccgga 1440
catgggctct tctccggtag cggcggagct tccacatccg agccctggtc ccattgcctcc 1500
agcggctcat ggtcgcctcg cagctccttg ctctaacag tggaggccag acttaggcac 1560
agcacaatgc ccaccaccac cagtgtgccg cacaaggccg tggcggtagg gtatgtgtct 1620
gaaaatgagc gtggagattg ggctcgacg gctgacgcag atggaagact taaggcagcg 1680
gcagaagaag atgcaggcag ctgagttggt gtattctgat aagagtcaga ggtaactccc 1740
gttgcgggtgc tgttaacggt ggagggcagt gtagtctgag cagtactcgt tgctgccgcg 1800
cgcgccacca gacataatag ctgacagact aacagactgt tcttttccat gggctctttc 1860
tgcagatggc taggctctgt gctttcctga tggctcctggc ggtgatgagc tactggccaa 1920
cctgctctct aggatgtgac ctgctcaga ctcataacct caggaacaag agagccttga 1980
cactcctggt acaaatgagg agactctccc ctctctcctg cctgaaggac aggaaggact 2040
ttggattccc gcaggagaag gtggatgcc agcagatcaa gaaggctcaa gccatccctg 2100
tcctgagtga gctgaccag cagatcctga acatcttcac atcaaaggac tcattctgtg 2160
cttggaatgc aacctccta gactattct gcaatgacct ccaccagcag ctcaatgacc 2220
tgcaaggttg tctgatgcag caggtggggg tgcaggaatt tcccctgacc caggaagatg 2280
ccctgctggc tgtgaggaaa tacttcaca ggatcactgt gtacctgaga gagaagaaac 2340
acagcccctg tgctgggag gtgtcagag cagaagtctg gagagccctg tcttctctg 2400
ccaatgtgct ggggaagactg agagaagaga aatgaagatc tgctgtgcct tctagttgcc 2460
agccatctgt tgtttgccc tccccgtgc ctctctgac cctggaaggc gccactccca 2520
ctgtcctttc ctaataaaat gaggaaattg catcgattg tctgagtagg tgtattcta 2580
ttctgggggg tgggtgggg cagcacagca agggggagga ttgggaagac aatagcaggc 2640
atgctgggga tgcgtgggc tctatgggt cccaggtgct gaagaattga cccggttcc 2700
cctgggccag aaagaagcag gcacatcccc ttctctgtga cacacctgt ccacgccct 2760
ggttcttagt tccagcccca ctcataggac actcatagct caggagggt cgccttcaa 2820
tcccaccgc taaagtactt ggagcgtct ctccctcct catcagccca ccaaaccaaa 2880

cttagcctcc aagagtggga agaaattaaa gcaagatagg ctattaagtg cagagggaga 2940
gaaaatgcct ccaacatgtg aggaagtaat gagagaaatc atagaatttc ttccgcttcc 3000
tcgctcactg actcgtcgcg ctcggtcggt cggtgcggc gagcgggtatc agctcactca 3060
aaggcggtaa tacggttatt cacagaatca ggggataacg caggaaagaa catgtgagca 3120
aaaggccagc aaaaggccag gaaccgtaaa aaggccgctg tgctggcggt tttccatagg 3180
ctccgcccc ctgacgagca tcacaaaaat cgacgctcaa gtcagagggt gcgaaacccg 3240
acaggactat aaagatacca ggcgtttccc cctggaagct ccctcgtcg ctctcctgtt 3300
ccgaccctgc cgttaccgg atacctgtcc gcctttctcc ctccgggaag cgtggcgctt 3360
tctcaatgct cagctgtag gtatctcagt tcggtgtagg tcgttcgctc caagctgggc 3420
tgtgtgcacg aacccccgt tcagcccgac cgctgcgct tatccggtaa ctatcgtctt 3480
gagtccaacc cggtaaagaca cgacttatcg ccactggcag cagccactgg taacaggatt 3540
agcagagcga ggtatgtagg cgggtctaca gagttettga agtgggtggc taactacggc 3600
tacactagaa ggacagtatt tggatctgc gctctgctga agccagttac ctccgaaaa 3660
agagttggta gctcttgatc cggcaaaaa accaccgctg gtagcgggtg tttttttgtt 3720
tgcaagcagc agattacgcg cagaaaaaaa ggatctcaag aagatccttt gatcttttct 3780
acggggtctg acgctcagtg gaacgaaaac tcacgttaag ggattttgt catgagatta 3840
tcaaaaagga tcttcaccta gatccttta aattaaaaat gaagtttta atcaatctaa 3900
agtatatatg agtaaacttg gtctgacagt taccaatgct taatcagtga ggcacctatc 3960
tcagcgatct gtctatttcg ttcattcata gttgcctgac tccggggggg gggggcgctg 4020
aggctcgct cgtgaagaag gtgttgctga ctcataccag gcctgaatcg ccccatcatc 4080
cagccagaaa gtgagggagc cacggttgat gagagctttg ttgtagggtg accagttgg 4140
gattttgaac ttttgctttg ccacggaacg gtctgcgttg tcgggaagat gcgtgatctg 4200
atccttcaac tcagcaaaag ttcgatttat tcaacaaagc cgccgtcccg tcaagtcagc 4260
gtaatgctct gcagtgta caaccaatta accaattctg attagaaaaa ctcatcgagc 4320
atcaaatgaa actgcaattt attcatatca ggattatcaa taccatattt ttgaaaaagc 4380
cgtttctgta atgaaggaga aaactcaccg aggcagttcc ataggatggc aagatcctgg 4440

tatcggctctg cgattccgac tcgtccaaca tcaatacaac ctattaattt cccctcgtca 4500
aaaaataaggt tatcaagtga gaaatcacca tgagtgcga ctgaatccgg tgagaatggc 4560
aaaagcttat gcatttcttt ccagacttgt tcaacaggcc agccattacg ctgcgtcatca 4620
aaatcactcg catcaaccaa accgttattc attcgtgatt gcgcctgagc gagacgaaat 4680
acgcgatcgc tgttaaaagg acaattacaa acaggaatcg aatgcaaccg gcgcaggaaac 4740
actgccagcg catcaacaat attttcacct gaatcaggat attcttctaa tacctggaat 4800
gctgttttcc cggggatcgc agtggtagt aaccatgcat catcaggagt acggataaaa 4860
tgcttgatgg tcggaagagg cataaattcc gtcagccagt ttagtctgac catctcatct 4920
gtaacatcat tggcaacgct acctttgcca tgtttcagaa acaactctgg cgcacgggc 4980
ttcccataca atcgatagat tgtcgacct gattgccga cattatcggc agcccattta 5040
taccatata aatcagcacc catgttgaa ttaatcggc gctcagaca agacgtttcc 5100
cgttgaatat ggtcataac accccttgta ttactgttta tgtaagcaga cagttttatt 5160
gttcatgatg atatattttt atcttgigca atgtaacatc agagattttg agacacaacg 5220
tggctttccc ccccccca ttattgaagc atttatcagg gttattgtct catgagcgga 5280
tacatatttg aatgtattta gaaaaataaa caaatagggg ttccggcac atttccccga 5340
aaagtgcac ctgacgtcta agaaaccatt attatcatga cattaacctt taaaaatagg 5400
cgtatcacga ggccctttcg tc 5422

<210> 6

<211> 5259

<212> DNA

<213> Mus musculus

<400> 6

tcgcgcgttt cggatgatgac ggtgaaaacc tctgacacat gcagtcocg gagacggtea 60
cagcttgtct gtaagcggat gccgggagca gacaagcccg tcagggcgcg tcagcgggtg 120
ttggcgggtg tcggggctgg cttaaactatg cggcatcaga gcagattgta ctgagagtgc 180
accatatgcg gtgtgaaata ccgcacagat gcgtaaggag aaaataccgc atcagattgg 240
ctattggcca ttgcatacgt tgtatccata tcataatatg tacatttata ttggctcatg 300

tccaacatta ccgccatggt gacattgatt attgactagt tattaatagt aatcaattac 360
ggggtcatta gttcatagcc catatatgga gttccgcgtt acataactta cggtaaatgg 420
cccgcctggc tgaccgccc aacgaccccg ccattgacg tcaataatga cgtatgttcc 480
catagtaacg ccaataggga ctttccattg acgtcaatgg gtggagtatt tacggtaaac 540
tgcccacttg gcagtacatc aagtgtatca tatgccagt acgcccccta ttgacgtcaa 600
tgacggtaaa tgcccgcct ggcattatgc ccagtacatg accttatggg actttcctac 660
ttggcagtag atctacgtat tagtcatgcg tattaccatg gtgatgcggg tttggcagta 720
catcaatggg cgtggatagc ggtttgactc acggggattt ccaagtctcc accccattga 780
cgtcaatggg agtttgtttt ggcacaaaaa tcaacgggac ttccaaaat gtcgtaacaa 840
ctccgcccc a ttgacgcaa tgggcggtag gcgtgtacgg tgggaggtct atataagcag 900
agtcgttta gtgaaccgtc agatgcctg gagacgcat ccacgtgtt ttgacctca 960
tagaagacac cgggaccgat ccagcctccg cggccgggaa cgggtgcattg gaacgggat 1020
tccccgtgc aagagtgcg taagtaccgc ctatagactc tataggcaca ccccttggc 1080
tcttatgcat gctatactgt ttttggcttg gggcctatac accccgctt ccttatgcta 1140
taggtgatgg tatagcttag cctatagggtg tgggttattg accattattg accactcccc 1200
tattggtgac gatactttcc attactaate cataacatgg ctctttgcca caactatctc 1260
tattggctat atgccaatac tctgtcctc agagactgac acggactctg tatttttaca 1320
ggatggggtc ccatttatta ttacaaatt cacatataca acaacgccgt cccccgtgc 1380
cgcagttttt attaaacata gcggtggatc tccacgcgaa tctcgggtac gtgttccgga 1440
catgggctct tctccgtag cggcggagct tccacatccg agcctggtc ccagtcctcc 1500
agcggctcat ggtcgtcgg cagctccttg ctctaacag tggaggccag acttaggcac 1560
agcacaatgc ccaccaccac cagtgtgccg cacaaggccg tggcggtagg gtatgtgtct 1620
gaaaatgagc gtggagattg ggctcgacg gctgacgcag atggaagact taaggcagcg 1680
gcagaagaag atgcaggcag ctgagttgtt gtattctgat aagagtcaga ggtaactccc 1740
gttgcggtgc tgtaacggg ggaggcaggt gtagtctgag cagtactcgt tgctgccgcg 1800
cgcgccacca gacataatag ctgacagact aacagactgt tcctttccat gggctctttc 1860

tgcagatggc taggctctgt gctttcctga tggtcctggc ggtgatgagc tactcgccaa 1920
cctgctctct aggatgtgac ctgcctcaga ctcataacct caggaacaag agagccttga 1980
cactcctggc aaaaatgagg agactctccc ctctctcctg cctgaaggac aggaaggact 2040
ttggattccc gcaggagaag gtggatgcc agcagatcaa gaaggctcaa gccatccctg 2100
tcctgagtga gctgaccag cagatcctga acatcttcac atcaaaggac tcctctgctg 2160
cttggaatgc aacctccta gactcattct gcaatgacct ccaccagcag ctcaatgacc 2220
tgcaagggtg tctgatgcag cagggtgggg tgcaaggaatt tccctgacc caggaagatg 2280
ccctgctggc tgtgaggaaa tacttccaca ggatcactgt gtacctgaga gagaagaaac 2340
acagccctg tgctgggag gtggtcagag cagaagtctg gagagccctg tcttctctg 2400
ccaatgtgct gggaagactg agagaagaga aatgaagatc cagatctact tctggctaata 2460
aaaagatcag agctctagag atctgtgtgt tggttttttg tgtggtaccc aggtgctgaa 2520
gaattgacct ggttctctct gggccagaaa gaagcaggca catcccttc tctgtgacac 2580
accctgtcca cggccctggc tcttagttcc agccctactc ataggacact catagctcag 2640
gagggctccg cttcaatcc caccgctaa agtacttga gcggtctctc cctccctcat 2700
cagcccacca aaccaaact agcctccaag agtggaaga aattaaagca agataggcta 2760
ttaagtgcag agggagagaa aatgcctcca acatgtgagg aagtaatgag agaaatcata 2820
gaatttcttc cgcttctctg ctactgact cgctgcgctc ggtcgttcgg ctgcggcgag 2880
cggatcagc tactcaaag gcggaatac ggttatccac agaatacagg gataacgcag 2940
gaaagaacat gtgagcaaaa ggccagcaaa aggccaggaa ccgtaaaaag gccgcgttg 3000
tggcgttttt ccataggctc cgccccctg acgagcatca caaaaatcga cgctcaagtc 3060
agaggtggcg aaaccgaca ggactataaa gataccaggc gtttccccct ggaagctccc 3120
tcgtgcgctc tctgttccg acctgccgc ttaccggata cctgtccgc tttctccctt 3180
cgggaagcgt ggcgctttct caatgctcac gctgtaggta tctcagttcg gtgtaggtcg 3240
ttcgctccaa gctgggctgt gtgcacgaac cccccgttca gccgaaccgc tgcgccttat 3300
ccggttaacta tcgtcttgag tccaaccgg taagacacga cttatcgcca ctggcagcag 3360
ccactggtaa caggattagc agagcgagg atgtaggcg tgctacagag ttcttgaagt 3420

ggtaggctaa ctacggctac actagaagga cagtatttgg tatctgcgct ctgctgaagc 3480
cagttacctt cggaaaaaga gttggtagct cttgatccgg caaacaacc accgctggta 3540
gcggtggttt ttttgtttgc aagcagcaga ttacgcgcag aaaaaagga tctcaagaag 3600
atcctttgat cttttctacg gggctcgacg ctcaagtggaa cgaaaactca cgtaaggga 3660
ttttggtcat gagattatca aaaaggatct tcacctagat ctttttaaat taaaaatgaa 3720
gttttaaatc aatctaaagt atatatgagt aaacttggtc tgacagttaac caatgcttaa 3780
tcagtgaggc acctatctca gcgatctgtc tatttcgttc atccatagtt gcctgactcc 3840
gggggggggg ggcgctgagg tctgcctcgt gaagaagggtg ttgctgactc ataccaggcc 3900
tgaatcgccc catcatccag ccagaaagtg agggagccac ggttgatgag agctttgttg 3960
taggtggacc agttggtgat ttgaacttt tgctttgccg cggaacggtc tgcgttgctg 4020
ggaagatgag tgatctgac cttcaactca gcaaaagttc gatttattca acaaagccgc 4080
cgtcccgta agtcagcgta atgctcggc agtggtacaa ccaattaacc aattctgatt 4140
agaaaaactc atcgagcatc aaatgaaact gcaatttatt catatcagga ttatcaatac 4200
catatttttg aaaaagccgt ttctgtaatg aaggagaaaa ctcaccgagg cagttccata 4260
ggatggcaag atcctggtat cggctcgcga ttccgactcg tccaacatca atacaacct 4320
ttaatttccc ctgctcaaaa ataagggtat caagtgagaa atcaccatga gtgacgactg 4380
aatccggtga gaatggcaaa agcttatgca tttctttcca gaattgttca acaggccagc 4440
cattacgctc gtcataaaaa tctctgcat caaccaaacc gtattcatt cgtgattgag 4500
cctgagcgag acgaaatacg cgatcgctgt taaaaggaca attacaaaca ggaatcgaat 4560
gcaaccggcg caggaacact gccagcgcat caacaatatt ttcacctgaa tcaggatatt 4620
cttctaatac ctggaatgct gttttcccg ggatcgaggt ggtgagtaac catgcatcat 4680
caggagtacg gataaaatgc ttgatggctg gaagaggcat aaattccgct agccagttta 4740
gtctgaccat ctcatctgta acatcattg caacgctacc tttgccatgt ttcagaaaca 4800
actctggcgc atcgggcttc ccatacaatc gatagattgt cgcacctgat tgcccgacat 4860
tatcgcgagc ccattttatac ccataataat cagcatccat gttggaattt aatcgcggcc 4920
tcgagcaaga cgtttccgt tgaatatggc tcataacacc cttgtatta ctgtttatgt 4980

-23-

aagcagacag ttttattgtt catgatgata tattttttatc ttgtgcaatg taacatcaga 5040
 gatttttgaga cacaacgtgg ctttcccccc cccccatta ttgaagcatt tatcagggtt 5100
 attgtctcat gagcggatac atatttgaat gtatttagaa aaataaacia ataggggttc 5160
 cgcgcacatt tccccgaaaa gtgccacctg acgtctaaga aaccattatt atcatgacat 5220
 taacctataa aaataggcgt atcacgaggc cctttcgtc 5259

<210> 7
 <211> 585
 <212> DNA
 <213> Homo sapiens

<220>
 <221> CDS
 <222> (1)..(585)

<220>
 <221> sig_peptide
 <222> (1)..(69)

<220>
 <221> mat_peptide
 <222> (70)..(585)

<400> 7
 atg gcc ctc ctg ttc cct cta ctg gca gcc cta gtg atg acc agc tat 48
 Met Ala Leu Leu Phe Pro Leu Leu Ala Ala Leu Val Met Thr Ser Tyr
 -20 -15 -10
 agc cct gtt gga tct ctg ggc tgt gat ctg cct cag aac cat ggc cta 96
 Ser Pro Val Gly Ser Leu Gly Cys Asp Leu Pro Gln Asn His Gly Leu
 -5 -1 1 5
 ctt agc agg aac acc ttg gtg ctt ctg cac caa atg agg aga atc tcc 144
 Leu Ser Arg Asn Thr Leu Val Leu Leu His Gln Met Arg Arg Ile Ser
 10 15 20 25
 cct ttc ttg tgt ctc aag gac aga aga gac ttc agg ttc ccc cag gag 192
 Pro Phe Leu Cys Leu Lys Asp Arg Arg Asp Phe Arg Phe Pro Gln Glu
 30 35 40
 atg gta aaa ggg agc cag ttg cag aag gcc cat gtc atg tct gtc ctc 240
 Met Val Lys Gly Ser Gln Leu Gln Lys Ala His Val Met Ser Val Leu
 45 50 55
 cat gag atg ctg cag cag atc ttc agc ctc ttc cac aca gag cgc tcc 288
 His Glu Met Leu Gln Gln Ile Phe Ser Leu Phe His Thr Glu Arg Ser
 60 65 70

-24-

tct gct gcc tgg aac atg acc ctc cta gac caa ctc cac act gga ctt 336
 Ser Ala Ala Trp Asn Met Thr Leu Leu Asp Gln Leu His Thr Gly Leu
 75 80 85

cat cag caa ctg caa cac ctg gag acc tgc ttg ctg cag gta gtg gga 384
 His Gln Gln Leu Gln His Leu Glu Thr Cys Leu Leu Gln Val Val Gly
 90 95 100 105

gaa gga gaa tct gct ggg gca att agc agc cct gca ctg acc ttg agg 432
 Glu Gly Glu Ser Ala Gly Ala Ile Ser Ser Pro Ala Leu Thr Leu Arg
 110 115 120

agg tac ttc cag gga atc cgt gtc tac ctg aaa gag aag aaa tac agc 480
 Arg Tyr Phe Gln Gly Ile Arg Val Tyr Leu Lys Glu Lys Lys Tyr Ser
 125 130 135

gac tgt gcc tgg gaa gtt gtc aga atg gaa atc atg aaa tcc ttg ttc 528
 Asp Cys Ala Trp Glu Val Val Arg Met Glu Ile Met Lys Ser Leu Phe
 140 145 150

tta tca aca aac atg caa gaa aga ctg aga agt aaa gat aga gac ctg 576
 Leu Ser Thr Asn Met Gln Glu Arg Leu Arg Ser Lys Asp Arg Asp Leu
 155 160 165

ggc tca tct 585
 Gly Ser Ser
 170

<210> 8
 <211> 195
 <212> PRT
 <213> Homo sapiens

<400> 8
 Met Ala Leu Leu Phe Pro Leu Leu Ala Ala Leu Val Met Thr Ser Tyr
 -20 -15 -10

Ser Pro Val Gly Ser Leu Gly Cys Asp Leu Pro Gln Asn His Gly Leu
 -5 -1 1 5

Leu Ser Arg Asn Thr Leu Val Leu Leu His Gln Met Arg Arg Ile Ser
 10 15 20 25

Pro Phe Leu Cys Leu Lys Asp Arg Arg Asp Phe Arg Phe Pro Gln Glu
 30 35 40

Met Val Lys Gly Ser Gln Leu Gln Lys Ala His Val Met Ser Val Leu
 45 50 55

His Glu Met Leu Gln Gln Ile Phe Ser Leu Phe His Thr Glu Arg Ser

-25-

60 65 70
 Ser Ala Ala Trp Asn Met Thr Leu Leu Asp Gln Leu His Thr Gly Leu
 75 80 85
 His Gln Gln Leu Gln His Leu Glu Thr Cys Leu Leu Gln Val Val Gly
 90 95 100 105
 Glu Gly Glu Ser Ala Gly Ala Ile Ser Ser Pro Ala Leu Thr Leu Arg
 110 115 120
 Arg Tyr Phe Gln Gly Ile Arg Val Tyr Leu Lys Glu Lys Lys Tyr Ser
 125 130 135
 Asp Cys Ala Trp Glu Val Val Arg Met Glu Ile Met Lys Ser Leu Phe
 140 145 150
 Leu Ser Thr Asn Met Gln Glu Arg Leu Arg Ser Lys Asp Arg Asp Leu
 155 160 165
 Gly Ser Ser
 170

<210> 9

<211> 567

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1)..(567)

<220>

<221> sig_peptide

<222> (1)..(69)

<220>

<221> mat_peptide

<222> (70)..(567)

<400> 9

atg gcc tcg ccc ttt gct tta ctg atg gtc ctg gtg gtg ctc agc tgc 48
 Met Ala Ser Pro Phe Ala Leu Leu Met Val Leu Val Val Leu Ser Cys
 -20 -15 -10

aag tca agc tgc tct ctg ggc tgt gat ctc cct gag acc cac agc ctg 96
 Lys Ser Ser Cys Ser Leu Gly Cys Asp Leu Pro Glu Thr His Ser Leu
 -5 -1 1 5

gat aac agg agg acc ttg atg ctc ctg gca caa atg agc aga atc tct 144
 Asp Asn Arg Arg Thr Leu Met Leu Leu Ala Gln Met Ser Arg Ile Ser

-26-

10	15	20	25	
cct tcc tcc tgt ctg atg gac aga cat gac ttt gga ttt ccc cag gag	192			
Pro Ser Ser Cys Leu Met Asp Arg His Asp Phe Gly Phe Pro Gln Glu				
30	35	40		
gag ttt gat ggc aac cag ttc cag aag gct cca gcc atc tct gtc ctc	240			
Glu Phe Asp Gly Asn Gln Phe Gln Lys Ala Pro Ala Ile Ser Val Leu				
45	50	55		
cat gag ctg atc cag cag atc ttc aac ctc ttt acc aca aaa gat tca	288			
His Glu Leu Ile Gln Gln Ile Phe Asn Leu Phe Thr Thr Lys Asp Ser				
60	65	70		
tct gct gct tgg gat gag gac ctc cta gac aaa ttc tgc acc gaa ctc	336			
Ser Ala Ala Trp Asp Glu Asp Leu Leu Asp Lys Phe Cys Thr Glu Leu				
75	80	85		
tac cag cag ctg aat gac ttg gaa gcc tgt gtg atg cag gag gag agg	384			
Tyr Gln Gln Leu Asn Asp Leu Glu Ala Cys Val Met Gln Glu Glu Arg				
90	95	100	105	
gtg gga gaa act ccc ctg atg aat gcg gac tcc atc ttg gct gtg aag	432			
Val Gly Glu Thr Pro Leu Met Asn Ala Asp Ser Ile Leu Ala Val Lys				
110	115	120		
aaa tac ttc cga aga atc act ctc tat ctg aca gag aag aaa tac agc	480			
Lys Tyr Phe Arg Arg Ile Thr Leu Tyr Leu Thr Glu Lys Lys Tyr Ser				
125	130	135		
cct tgt gcc tgg gag gtt gtc aga gca gaa atc atg aga tcc ctc tct	528			
Pro Cys Ala Trp Glu Val Val Arg Ala Glu Ile Met Arg Ser Leu Ser				
140	145	150		
tta tca aca aac ttg caa gaa aga tta agg agg aag gaa	567			
Leu Ser Thr Asn Leu Gln Glu Arg Leu Arg Arg Lys Glu				
155	160	165		

<210> 10

<211> 189

<212> PRT

<213> Homo sapiens

<400> 10

Met Ala Ser Pro Phe Ala Leu Leu Met Val Leu Val Val Leu Ser Cys
-20 -15 -10

Lys Ser Ser Cys Ser Leu Gly Cys Asp Leu Pro Glu Thr His Ser Leu
-5 -1 1 5

Asp Asn Arg Arg Thr Leu Met Leu Leu Ala Gln Met Ser Arg Ile Ser

-27-

10 15 20 25
 Pro Ser Ser Cys Leu Met Asp Arg His Asp Phe Gly Phe Pro Gln Glu
 30 35 40
 Glu Phe Asp Gly Asn Gln Phe Gln Lys Ala Pro Ala Ile Ser Val Leu
 45 50 55
 His Glu Leu Ile Gln Gln Ile Phe Asn Leu Phe Thr Thr Lys Asp Ser
 60 65 70
 Ser Ala Ala Trp Asp Glu Asp Leu Leu Asp Lys Phe Cys Thr Glu Leu
 75 80 85
 Tyr Gln Gln Leu Asn Asp Leu Glu Ala Cys Val Met Gln Glu Glu Arg
 90 95 100 105
 Val Gly Glu Thr Pro Leu Met Asn Ala Asp Ser Ile Leu Ala Val Lys
 110 115 120
 Lys Tyr Phe Arg Arg Ile Thr Leu Tyr Leu Thr Glu Lys Lys Tyr Ser
 125 130 135
 Pro Cys Ala Trp Glu Val Val Arg Ala Glu Ile Met Arg Ser Leu Ser
 140 145 150
 Leu Ser Thr Asn Leu Gln Glu Arg Leu Arg Arg Lys Glu
 155 160 165

<210> 11

<211> 567

<212> DNA

<213> Mus musculus

<220>

<221> CDS

<222> (1)..(567)

<220>

<221> sig_peptide

<222> (1)..(69)

<220>

<221> mat_peptide

<222> (70)..(567)

<400> 11

atg gct agg ctc tgt gct ttc ctg atg gtc ctg gcg gtg atg agc tac 48
 Met Ala Arg Leu Cys Ala Phe Leu Met Val Leu Ala Val Met Ser Tyr
 -20 -15 -10

-28-

tgg cca acc tgc tct cta gga tgt gac ctg cct cag act cat aac ctc 96
 Trp Pro Thr Cys Ser Leu Gly Cys Asp Leu Pro Gln Thr His Asn Leu
 -5 -1 1 5

agg aac aag aga gcc ttg aca ctc ctg gta caa atg agg aga ctc tcc 144
 Arg Asn Lys Arg Ala Leu Thr Leu Leu Val Gln Met Arg Arg Leu Ser
 10 15 20 25

cct ctc tcc tgc ctg aag gac agg aag gac ttt gga ttc ccg cag gag 192
 Pro Leu Ser Cys Leu Lys Asp Arg Lys Asp Phe Gly Phe Pro Gln Glu
 30 35 40

aag gtg gat gcc cag cag atc aag aag gct caa gcc atc cct gtc ctg 240
 Lys Val Asp Ala Gln Gln Ile Lys Lys Ala Gln Ala Ile Pro Val Leu
 45 50 55

agt gag ctg acc cag cag atc ctg aac atc ttc aca tca aag gac tca 288
 Ser Glu Leu Thr Gln Gln Ile Leu Asn Ile Phe Thr Ser Lys Asp Ser
 60 65 70

tct gct gct tgg aat gca acc ctc cta gac tca ttc tgc aat gac ctc 336
 Ser Ala Ala Trp Asn Ala Thr Leu Leu Asp Ser Phe Cys Asn Asp Leu
 75 80 85

cac cag cag ctc aat gac ctg caa ggt tgt ctg atg cag cag gtg ggg 384
 His Gln Gln Leu Asn Asp Leu Gln Gly Cys Leu Met Gln Gln Val Gly
 90 95 100 105

gtg cag gaa ttt ccc ctg acc cag gaa gat gcc ctg ctg gct gtg agg 432
 Val Gln Glu Phe Pro Leu Thr Gln Glu Asp Ala Leu Leu Ala Val Arg
 110 115 120

aaa tac ttc cac agg atc act gtg tac ctg aga gag aag aaa cac agc 480
 Lys Tyr Phe His Arg Ile Thr Val Tyr Leu Arg Glu Lys Lys His Ser
 125 130 135

ccc tgt gcc tgg gag gtg gtc aga gca gaa gtc tgg aga gcc ctg tct 528
 Pro Cys Ala Trp Glu Val Val Arg Ala Glu Val Trp Arg Ala Leu Ser
 140 145 150

tcc tct gcc aat gtg ctg gga aga ctg aga gaa gag aaa 567
 Ser Ser Ala Asn Val Leu Gly Arg Leu Arg Glu Glu Lys
 155 160 165

<210> 12

<211> 189

<212> PRT

<213> Mus musculus

<400> 12

Met Ala Arg Leu Cys Ala Phe Leu Met Val Leu Ala Val Met Ser Tyr

-29-

-20 -15 -10
 Trp Pro Thr Cys Ser Leu Gly Cys Asp Leu Pro Gln Thr His Asn Leu
 -5 -1 1 5
 Arg Asn Lys Arg Ala Leu Thr Leu Leu Val Gln Met Arg Arg Leu Ser
 10 15 20 25
 Pro Leu Ser Cys Leu Lys Asp Arg Lys Asp Phe Gly Phe Pro Gln Glu
 30 35 40
 Lys Val Asp Ala Gln Gln Ile Lys Lys Ala Gln Ala Ile Pro Val Leu
 45 50 55
 Ser Glu Leu Thr Gln Gln Ile Leu Asn Ile Phe Thr Ser Lys Asp Ser
 60 65 70
 Ser Ala Ala Trp Asn Ala Thr Leu Leu Asp Ser Phe Cys Asn Asp Leu
 75 80 85
 His Gln Gln Leu Asn Asp Leu Gln Gly Cys Leu Met Gln Gln Val Gly
 90 95 100 105
 Val Gln Glu Phe Pro Leu Thr Gln Glu Asp Ala Leu Leu Ala Val Arg
 110 115 120
 Lys Tyr Phe His Arg Ile Thr Val Tyr Leu Arg Glu Lys Lys His Ser
 125 130 135
 Pro Cys Ala Trp Glu Val Val Arg Ala Glu Val Trp Arg Ala Leu Ser
 140 145 150
 Ser Ser Ala Asn Val Leu Gly Arg Leu Arg Glu Glu Lys
 155 160 165

<210> 13

<211> 459

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1)..(459)

<220>

<221> sig_peptide

<222> (1)..(60)

<220>

<221> mat_peptide

<222> (61)..(459)

-30-

<400> 13

atg tac agg atg caa ctc ctg tct tgc att gca cta agt ctt gca ctt 48
 Met Tyr Arg Met Gln Leu Leu Ser Cys Ile Ala Leu Ser Leu Ala Leu
 -20 -15 -10 -5

 gtc aca aac agt gca cct act tca agt tct aca aag aaa aca cag cta 96
 Val Thr Asn Ser Ala Pro Thr Ser Ser Ser Thr Lys Lys Thr Gln Leu
 -1 1 5 10

 caa ctg gag cat tta ctt ctg gat tta cag atg att ttg aat gga att 144
 Gln Leu Glu His Leu Leu Leu Asp Leu Gln Met Ile Leu Asn Gly Ile
 15 20 25

 aat aat tac aag aat ccc aaa ctc acc agg atg ctc aca ttt aag ttt 192
 Asn Asn Tyr Lys Asn Pro Lys Leu Thr Arg Met Leu Thr Phe Lys Phe
 30 35 40

 tac atg ccc aag aag gcc aca gaa ctg aaa cat ctt cag tgt cta gaa 240
 Tyr Met Pro Lys Lys Ala Thr Glu Leu Lys His Leu Gln Cys Leu Glu
 45 50 55 60

 gaa gaa ctc aaa cct ctg gag gaa gtg cta aat tta gct caa agc aaa 288
 Glu Glu Leu Lys Pro Leu Glu Glu Val Leu Asn Leu Ala Gln Ser Lys
 65 70 75

 aac ttt cac tta aga ccc agg gac tta atc agc aat atc aac gta ata 336
 Asn Phe His Leu Arg Pro Arg Asp Leu Ile Ser Asn Ile Asn Val Ile
 80 85 90

 gtt ctg gaa cta aag gga tct gaa aca aca ttc atg tgt gaa tat gct 384
 Val Leu Glu Leu Lys Gly Ser Glu Thr Thr Phe Met Cys Glu Tyr Ala
 95 100 105

 gat gag aca gca acc att gta gaa ttt ctg aac aga tgg att acc ttt 432
 Asp Glu Thr Ala Thr Ile Val Glu Phe Leu Asn Arg Trp Ile Thr Phe
 110 115 120

 tgt caa agc atc atc tca aca ctg act 459
 Cys Gln Ser Ile Ile Ser Thr Leu Thr
 125 130

<210> 14

<211> 153

<212> PRT

<213> Homo sapiens

<400> 14

Met Tyr Arg Met Gln Leu Leu Ser Cys Ile Ala Leu Ser Leu Ala Leu
 -20 -15 -10 -5

-31-

Val Thr Asn Ser Ala Pro Thr Ser Ser Ser Thr Lys Lys Thr Gln Leu
 -1 1 5 10

Gln Leu Glu His Leu Leu Leu Asp Leu Gln Met Ile Leu Asn Gly Ile
 15 20 25

Asn Asn Tyr Lys Asn Pro Lys Leu Thr Arg Met Leu Thr Phe Lys Phe
 30 35 40

Tyr Met Pro Lys Lys Ala Thr Glu Leu Lys His Leu Gln Cys Leu Glu
 45 50 55 60

Glu Glu Leu Lys Pro Leu Glu Glu Val Leu Asn Leu Ala Gln Ser Lys
 65 70 75

Asn Phe His Leu Arg Pro Arg Asp Leu Ile Ser Asn Ile Asn Val Ile
 80 85 90

Val Leu Glu Leu Lys Gly Ser Glu Thr Thr Phe Met Cys Glu Tyr Ala
 95 100 105

Asp Glu Thr Ala Thr Ile Val Glu Phe Leu Asn Arg Trp Ile Thr Phe
 110 115 120

Cys Gln Ser Ile Ile Ser Thr Leu Thr
 125 130

<210> 15

<211> 26

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:PCR oligo

<400> 15

aactgcagat ggctaggctc tgtgct

26

<210> 16

<211> 26

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:PCR oligo

<400> 16

gaagatcttc atttctcttc tctcag

26

-32-

<210> 17
<211> 26
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:PCR oligo

<400> 17
aactgcagat ggcctcgccc ttgtct

26

<210> 18
<211> 28
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:PCR oligo

<400> 18
cgggatcctt attccttcct ccttaatc

28

<210> 19
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:PCR oligo

<400> 19
gctctagatg gccctcctgt tccct

25

<210> 20
<211> 26
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:PCR oligo

<400> 20
gcggatcctc aagatgagcc caggtc

26

<210> 21
<211> 36
<212> DNA
<213> Artificial Sequence

-33-

<220>

<223> Description of Artificial Sequence: PCR oligo

<400> 21

acgcgtcgac atgtgtcctc agaagctaac catctc

36

<210> 22

<211> 34

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: PCR oligo

<400> 22

gcggatcctt aggatcggac cctgcaggga acac

34

<210> 23

<211> 37

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: PCR oligo

<400> 23

catgccatgg gtcaatcacg ctacctctc tttttgg

37

<210> 24

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: PCR oligo

<400> 24

gcggatcctc aggcggagct cagatagccc

30

<210> 25

<211> 5469

<212> DNA

<213> Mus musculus

<220>

<221> CDS

<222> (1923)..(2393)

<220>

<221> sig_peptide

<222> (1923) .. (1994)

<220>

<221> mat_peptide

<222> (1995) .. (2393)

<400> 25

tcgcgcgttt cggatgatgac ggtgaaaacc tctgacacat gcagctcccg gagacgggtca 60
cagcttgtct gtaagcggat gccggggagca gacaagcccc tcagggcgcg tcagcgggtg 120
ttggcgggtg tcggggctgg ctttaactatg cggcatcaga gcagattgta ctgagagtgc 180
accatatgcg gtgtgaaata ccgcacagat gcgtaaggag aaaataccgc atcagattgg 240
ctattggcca ttgcatacgt tgtatccata tcataatatg tacatttata ttggctcatg 300
tccaacatta ccgcatggt gacattgatt attgactagt tattaatagt aatcaattac 360
ggggtcatta gttcatagcc catatatgga gttccgcgtt acataactta cggtaaatgg 420
cccgcctggc tgaccgccc aagaccccc cccattgacg tcaataatga cgtatgttcc 480
catagtaacg ccaataggga ctttccattg acgtcaatgg gtggagtatt tacggtaaac 540
tgcccacttg gcagtacatc aagtgtatca tatgccaagt acgcccccta ttgacgtcaa 600
tgacggtaaa tggcccgctt ggcattatgc ccagtacatg accttatggg actttcctac 660
ttggcagtac atctacgtat tagtcatgcg tattaccatg gtgatgcggg tttggcagta 720
catcaatggg cgtggatagc ggtttgactc acggggattt ccaagtctcc accccattga 780
cgtcaatggg agtttgtttt ggcacaaaaa tcaacgggac tttccaaaat gtcgtaacaa 840
ctccgcccc ttgacgcaaa tgggcggtag gcgtgtacgg tgggaggtct atataagcag 900
agctcgttta gtgaaccgtc agatcgctg gagacgcat ccacgctgtt ttgacctcca 960
tagaagacac cgggaccgat ccagcctccg cggccgggaa cgtgcattg gaacgcgat 1020
tccccgtgcc aagagtgacg taagtaccgc ctatagagtc tataggccca cccccttggc 1080
ttcttatgca tgctatactg tttttggctt ggggtctata ccccccgct tcctcatgtt 1140
ataggatgat gtatagetta gcctataggt gtgggttatt gaccattatt gaccactccc 1200
ctattggtga cgatactttc cattactaat ccataacatg gctctttgcc acaactctct 1260
ttattggcta tatgccaata cactgtcctt cagagactga cacggactct gtatttttac 1320
aggatggggt ctcatattt atttacaaat tcacatatac aacaccaccg tcccagtg 1380

ccgcagtttt tattaacat aacgtgggat ctccacgcga atctcgggta cgtgttccgg 1440
 acatgggctc ttctccggtg gcggcggagc ttctacatcc gagccctgct cccatgcctc 1500
 cagcgactca tggtcgctcg gcagctcctt gctcctaaca gtggaggcca gacttaggca 1560
 cagcacgatg cccaccacca ccagtgtgcc gcacaaggcc gtggcggtag ggtatgtgtc 1620
 tgaaaatgag ctccggggagc gggcttgca cgcgtgacgca tttggaagac ttaaggcagc 1680
 ggcaagaaga gatgcaggca gctgagttgt tgtgttctga taagagtcag aggttaactcc 1740
 cgttgccggtg ctgttaacgg tggagggcag tgtagtctga gcagtactcg ttgctgccgc 1800
 gcgcgccacc agacataata gctgacagac taacagactg ttcctttcca tgggtctttt 1860
 ctgcagtcac cgtcctraga tctagcctca accctgacta tcttcaggt cattgttcca 1920
)
 ac atg gcc ctg tgg atc gac agg atg caa ctc ctg tct tgc att gca 1967
 Met Ala Leu Trp Ile Asp Arg Met Gln Leu Leu Ser Cys Ile Ala
 -20 -15 -10
 cta agt ctt gca ctt gtc aca aac agt gca oct act tca agt tct aca 2015
 Leu Ser Leu Ala Leu Val Thr Asn Ser Ala Pro Thr Ser Ser Ser Thr
 -5 -1 1 5
 aag aaa aca cag cta caa ctg gag cat tta ctg ctg gat tta cag atg 2063
 Lys Lys Thr Gln Leu Gln Leu Glu His Leu Leu Leu Asp Leu Gln Met
 10 15 20
 att ttg aat gga att aat aat tac aag aat ccc aaa ctc acc agg atg 2111
 Ile Leu Asn Gly Ile Asn Asn Tyr Lys Asn Pro Lys Leu Thr Arg Met
 25 30 35
 ctc aca ttt aag ttt tac atg ccc aag aag gcc aca gaa ctg aaa cat 2159
 Leu Thr Phe Lys Phe Tyr Met Pro Lys Lys Ala Thr Glu Leu Lys His
 40 45 50 55
 ctt cag tgt cta gaa gaa gaa ctc aaa cct ctg gag gaa gtg cta aat 2207
 Leu Gln Cys Leu Glu Glu Glu Leu Lys Pro Leu Glu Glu Val Leu Asn
 60 65 70
 tta gct caa agc aaa aac ttt cac tta aga ccc agg gac tta atc agc 2255
 Leu Ala Gln Ser Lys Asn Phe His Leu Arg Pro Arg Asp Leu Ile Ser
 75 80 85
 aat atc aac gta ata gtt ctg gaa cta aag gga tct gaa aca aca ttc 2303
 Asn Ile Asn Val Ile Val Leu Glu Leu Lys Gly Ser Glu Thr Thr Phe
 90 95 100
 atg tgt gaa tat gct gat gag aca gca acc att gta gaa ttt ctg aac 2351

-36-

Met Cys Glu Tyr Ala Asp Glu Thr Ala Thr Ile Val Glu Phe Leu Asn
 105 110 115

aga tgg att acc ttt tgt caa agc atc atc tca aca cta act 2393

Arg Trp Ile Thr Phe Cys Gln Ser Ile Ile Ser Thr Leu Thr
 120 125 130

tgataattaa gtgcttccca cttaaaacat atcagggatc tcgactctag aggatcatcg 2453

cggccgctct agaccaggcg cctggatcca gatctgctgt gccttctagt tgccagccat 2513

ctgttgtttg cccctcccc gtgccttctt tgaccctgga aggtgccact cccactgtcc 2573

tttcctaata aaatgaggaa attgcatcgc attgtctgag taggtgtcat tctattctgg 2633

ggggtggggt ggggcagcac agcaaggggg aggattggga agacaatagc aggcattgtg 2693

gggatgcggt gggctctatg ggtaccagg tgctgaagaa ttgaccgggt tctctctggg 2753

ccagaaagaa gcaggcacat ccccttctct gtgacacacc ctgtccacgc ccttggttct 2813

tagttccagc cccactcata ggacactcat agctcaggag ggctccgcct tcaatccac 2873

ccgctaaagt acttgagcgg gtctctccct cctcatcag cccaccaaac caaacctagc 2933

ctccaagagt gggagaaat taaagcaaga taggctatta agtgacagagg gagagaaaat 2993

gcctccaaca tgtgaggaag taatgagaga aatcatagaa tttcttcgc ttcctcgctc 3053

actgactcgc tgcgctcggg cgttcggctg cggcgagcgg tatcagctca ctcaaaggcg 3113

gtaatacggg tatccacaga atcaggggat aacgcaggaa agaacatgtg agcaaaaggc 3173

cagcaaaagg ccaggaaccg taaaaaggcc gcgttgctgg cgtttttcca taggctccgc 3233

cccctgacg agcatcaca aaatcgacgc tcaagtcaga ggtggcgaaa cccgacagga 3293

ctataaagat accaggcgtt tccccctgga agtccctcg tgcgctctcc tgttccgacc 3353

ctgcgcgtta ccgatacct gtccgccttt ctcccttcgg gaagcgtggc gctttctcaa 3413

tgctcacgct gtaggtatct cagttcgggt taggtcgttc gctccaagct gggctgtgtg 3473

cacgaacccc ccgttcagcc cgaccgctgc gccttatccg gtaactatcg tcttgagtcc 3533

aaccgggtaa gacacgactt atcgccactg gcagcagcca ctggtaacag gattagcaga 3593

gcgaggatat taggcggtgc tacagagttc ttgaagtggg gccctaacta cggctacact 3653

agaaggacag tatttggtat ctgcgctctg ctgaagccag ttaccttcgg aaaaagagtt 3713

ggtagctctt gatccggcaa acaaacacc gctggtagcg gtggtttttt tgtttgcaag 3773

cagcagatta cgcgagaaa aaaaggatct caagaagatc ctttgatctt ttctacggg 3833
tctgacgctc agtggaacga aaactcacgt taagggattt tggcatgag attatcaaaa 3893
aggatcttca cctagatcct tttaaattaa aatgaagt tttaatcaat ctaaagtata 3953
tatgagtaaa cttggtctga cagttacca tgettaatca gtgaggcacc tatctcagcg 4013
atctgtctat ttcgttcac catagttgcc tgactccggg gggggggggc gctgaggtct 4073
gectcgtgaa gaaggtgttg ctgactcata ccaggcctga atcgcccat catccagcca 4133
gaaagtgagg gagccacggt tgatgagagc ttgtttgtag gtggaccagt tggtgatttt 4193
gaacttttgc ttggccacgg aacggtctgc gttgtcggga agatgcgtga tctgatcctt 4253
caactcagca aaagtccgat ttattcaaca aagccgcgt cccgtcaagt cagcgtaatg 4313
ctctgccagt gttacaacca attaaccaat tctgattaga aaaactcatc gagcatcaaa 4373
tgaaactgca atttattcat atcaggatta tcaataccat atttttgaaa aagccgtttc 4433
tgtaatgaag gagaaaactc accgaggcag ttccatagga tggcaagatc ctggtatcgg 4493
tctgcgattc cgactcgtcc aacatcaata caacctatta atttccctc gtcaaaaata 4553
aggttatcaa gtgagaaatc accatgagtg acgactgaat ccggtgagaa tggcaaaagc 4613
ttatgcattt ctttcagac ttgttcaaca ggccagccat tacgctcgtc atcaaaatca 4673
ctcgcacaa ccaaaccgtt attcattcgt gattgcgcct gagcgagacg aaatacgcga 4733
tcgctgttaa aaggacaatt acaaacagga atcgaatgca accggcgag gaacactgcc 4793
agcgcacaa caatattttc acctgaatca ggatattctt ctaatacctg gaatgctgtt 4853
ttcccgggga tcgcagtggg gagtaaccat gcatcatcag gactacggat aaaatgcttg 4913
atggtcggaa gaggcataaa ttccgtcagc cagtttagtc tgaccatctc atctgtaaca 4973
tcattggcaa cgctacctt gccatgtttc agaaacaact ctggcgcatc gggcttccca 5033
tacaatcgat agattgtcgc acctgattgc ccgacattat cgcgagccca ttatatacca 5093
tataaatcag catccatgtt ggaatttaac cgcggcctcg agcaagacgt ttcccgttga 5153
atatggctca taacaccct tgtattactg ttatgtgtaag cagacagttt tattgttcat 5213
gatgatatat ttttatcttg tgcaatgtaa catcagagat ttgagacac aacgtggctt 5273
tcccccccc cccattattg aagcatttat cagggttatt gtctcatgag cggatacata 5333

-38-

tttgaatgta tttagaaaaa taaacaaata ggggttcgc gcacatttcc ccgaaaagtg 5393

ccacctgacg tctaagaaac cattattatc atgacattaa cctataaaaa taggcgtatc 5453

acgaggccct ttctgc 5469

<210> 26

<211> 157

<212> PRT

<213> Mus musculus

<400> 26

Met Ala Leu Trp Ile Asp Arg Met Gln Leu Leu Ser Cys Ile Ala Leu
-20 -15 -10

Ser Leu Ala Leu Val Thr Asn Ser Ala Pro Thr Ser Ser Ser Thr Lys
-5 -1 1 5

Lys Thr Gln Leu Gln Leu Glu His Leu Leu Leu Asp Leu Gln Met Ile
10 15 20

Leu Asn Gly Ile Asn Asn Tyr Lys Asn Pro Lys Leu Thr Arg Met Leu
25 30 35 40

Thr Phe Lys Phe Tyr Met Pro Lys Lys Ala Thr Glu Leu Lys His Leu
45 50 55

Gln Cys Leu Glu Glu Glu Leu Lys Pro Leu Glu Glu Val Leu Asn Leu
60 65 70

Ala Gln Ser Lys Asn Phe His Leu Arg Pro Arg Asp Leu Ile Ser Asn
75 80 85

Ile Asn Val Ile Val Leu Glu Leu Lys Gly Ser Glu Thr Thr Phe Met
90 95 100

Cys Glu Tyr Ala Asp Glu Thr Ala Thr Ile Val Glu Phe Leu Asn Arg
105 110 115 120

Trp Ile Thr Phe Cys Gln Ser Ile Ile Ser Thr Leu Thr
125 130